as a symptom. The division does not seem to me to be justified.

In a very few cases I have seen herpes labialis develop, but in so few that I can not state whether this is more likely to occur in childhood. At any age it is certainly very rare. Desquamation has been described by French observers as occurring frequently. Various accidental rashes have also been described, such as a rubeoloid and a scarlatinal eruption. They offer, however, no peculiarities in childhood.

Enlargement of the spleen is probably always present, although often, as in adult life, not discoverable. In the majority of cases this enlargement can be detected if sought for at the proper time and when the abdomen is not too much distended.

Respiration offers nothing peculiar, nor do affections of the respiratory apparatus. The statements of writers regarding the frequency of epistaxis in childhood vary very considerably. It is probably nearly as frequent as in adult life. The pulse is often unusually slow as compared with the rate which one would expect for the same elevation of temperature in other diseases. This is, as we know, also true of typhoid fever in adults.

The temperature in typhoid fever in children runs a course similar to that in the adult disease. It is, however, subject to still greater variations in early life. Not infrequently it rises rapidly without the usual step-like ascent. Often it remains high at the acme of the disease with little tendency to morning fall. Very often it falls toward the end of the attack with much greater rapidity than in adults. Indeed, the fall may be almost critical at times. The remittent character of the last part of an attack of adult typhoid is very liable to be absent in children. Quite certainly, as already pointed out, the temperature continues elevated for a decidedly shorter time than in adult life.

The tongue is not as liable to become dry as in adult life. In some instances, however, especially in later childhood, this symptom is not infrequently seen. Vomiting is a very common initial symptom, much more so than in adult life. Even later in the disease it is sometimes a most troublesome symptom. I have seen it the direct cause of death. Diarrhea is, in my experience, much more liable to be absent than present. This is the experience also of many others, especially in the cases occurring in early childhood. Certainly diarrhea is not more common than constipation at any period of childhood. Abdominal distension is not often a troublesome symptom, although often present to a considerable extent. This is in sharp contrast to the tympanites so common in adults. There are exceptions to this rule, and in these the distension may be excessive. Abdominal tenderness is seldom marked. Hemorrhage is of very rare occurrence. When it does take place it is nearly always in cases in later childhood, and especially in those over 9 or 10 years of age. I have seen it in one or two cases only, and then not severe. However, severe and even fatal hemorrhage has been reported. The infrequency of hemorrhage doubtless depends on the much less pronounced development of intestinal lesions, generally, but not always, characteristic of childhood. The same may be said of perforation, a very uncommon occurrence in typhoid fever in childhood, and almost never seen except toward the end of this period. I can recall perforation in only three or four cases in my experience; the youngest, I think, was 9, the oldest 12 years. Yet it would appear from

medical literature that perforation may take place under the age of 2 years.

One of the characteristics of typhoid fever in children is the tendency for the nervous symptoms to predominate over the intestinal. This by no means implies that nervous symptoms are more common and more severe in the disease in childhood than in adult life. On the contrary, in the majority of cases in childhood they are trifling or absent. Extreme torpor and the tendency to coma, coma vigil and carphology occur much less often than in adult life. They are especially rare in young children. Headache is a common symptom, especially at the beginning, but no more so than in adults. Slight delirium is apt to occur, especially at night. A certain degree of apathy is very common and often one of the most characteristic symptoms, especially in young children.

A condition not infrequently seen at the beginning, and liable to cause confusion in diagnosis, is that of pseudomeningitis. Meningitic symptoms do, it is true, occur in adult typhoid also, but certainly less characteristically. As a rule, the diagnosis can be made easily, but I have seen typhoid fever begin in children with all appearances of acute leptomeningitis, and lead for a time to an entirely erroneous diagnosis. Aphasia is a nervous symptom rather to be considered as a complication. It is generally said to occur more frequently in childhood.

The discussion of the complications and sequels, as of the diagnosis, is beyond the province of this paper.

REFERENCES.

1. N. Y. Med. Record, December, 1895, p. 803.
2. Am. Jour. of Obstet., February, 1895, p. 175.
3. Gaz. Hebdom., May 6, 1900.
4. Ibid., Feb. 8, 1900, p. 126.
5. Rev. Mens. des Malad. de l'Enf., Jan. 31, 1900.
6. Ibid., 1899, 555.
7. Pediatrics, viii, 1899, 527.
8. Jahrb. f. Kinderheilk., xlviii, 1898, 198.
9. Boston Med. and Surg. Jour., Feb. 27, 1896.
10. Arch. of Pediatrics, April, 1901.
11. Phila. Med. Jour., Oct. 15, 1898.

THE DIAGNOSIS OF TYPHOID FEVER IN THE LABORATORY.

JOHN LOVETT MORSE, A.M., M.D.

Instructor in Diseases of Children, Harvard Medical School; Assistant Visiting Physician at the City Hospital and at the Infants' Hospital.
BOSTON, MASS.

Before taking up the subject which has been assigned to me, I wish to protest against the separation of laboratory and clinical methods in the diagnosis of this disease. They should not be thought of separately, but together. Both are fallible, and both should be used understandingly and with due recognition of their limitations. Otherwise they may well lead to erroneous conclusions.

The various laboratory tests which are used for the diagnosis of typhoid fever are, briefly, as follows: The Widal reaction; examination of the urine and feces for typhoid bacilli; examination of the blood for typhoid bacilli; examination of rose spots for typhoid bacilli; the diazo-reaction in the urine; the leucocyte count; tests which show the presence or absence of other diseases. The value of these tests is very unequal. Some are easy to carry out, some difficult. All have their limitations. As this, however, does not seem either the time or place to go into the details of the performance of these tests, I shall confine

myself entirely to a consideration of their value and limitations.

THE WIDAL REACTION.

The Widal reaction is present in at least 95 per cent. of all cases of typhoid fever. It seldom appears, however, before the second week of fever, and may be delayed even until convalescence. In a small number of cases it may never be present; in others it may be intermittent. As it seldom appears before the beginning of the second week, the test is of little value up to this time. As it often appears late in the disease, and as it may be present only intermittently, typhoid fever can not be excluded by a single or even by repeated negative tests. Repeated negative tests, however, are very strong evidence against the existence of typhoid. The reaction may persist in the blood as long as twenty-one years after recovery from typhoid. Hence, a positive reaction in the absence of other symptoms of disease is not diagnostic of typhoid. A positive reaction, if the patient has not previously had typhoid, is almost certain proof of typhoid. A positive reaction may occur, however, in other diseases than typhoid. This error does not exceed 2 per cent. A negative reaction followed by a positive reaction in a dilution of 1 to 50 is absolute proof of typhoid.

The Widal reaction occurs under the same conditions and with the same limitations in children as in adults. There is some evidence to show, however, that in them the reaction appears earlier, is feebler, and persists for a shorter time than in adults. Owing to the comparative mildness and to the large number of atypical cases of typhoid in children, the Widal test is exceptionally important in them as an aid to the diagnosis of this disease. It is of especial value in two ways: 1, in ruling out many cases of gastro-intestinal disorders which might otherwise be mistaken for typhoid, and 2, in making a positive diagnosis possible in many mild cases which might otherwise pass unrecognized.

The Widal reaction appears in infantile typhoid as in that of older children and adults. The presence of the Widal reaction in infancy, especially in early infancy, is of less diagnostic value, however, than in adult life, as the reaction may be transmitted from the mother through the placenta or through the milk. The agglutinating power may be transmitted to the infant through the placenta, not only during the course of or convalescence from typhoid, but even when pregnancy takes place years after recovery. It is probable that it may also be transmitted through the milk after many years. When transmitted through the milk, however, it does not persist more than a week after the cessation of nursing. In early infancy, therefore, a positive Widal reaction is of somewhat less diagnostic value than in older children and adults. If the mother has had typhoid, and especially if she is nursing the infant, it should be looked on with some suspicion unless associated with other characteristic signs of typhoid. Examination of the mother's blood and milk and the cessation of breast-feeding will then assist in estimating the true value of the reaction in the infant.

This test, requiring, as it does, the maintenance and preparation of a fresh bouillon culture of the typhoid bacillus every twenty-four hours, is not a practicable one for the general practitioner. The boards of health of many states and of most of the large cities now make this test for the practitioner. All that is required of him is to send them a few drops of dried blood. This certainly does not appear impracticable,

or even difficult. The laboratory can report on the blood in less than an hour. That certainly seems rapid enough.

EXAMINATION OF THE FECES FOR TYPHOID BACILLI.

Typhoid bacilli have been found in the stools of typhoid fever patients by many observers. They have been found in some cases as early as the third or fourth day after taking to bed. They have not infrequently been found in the stools before the appearance of the Widal reaction in the blood. At least three days, and more often four or five days, are required to prove their presence, however. Although they are as a rule present in the stools before the Widal reaction appears in the blood, yet, so much more time is required for their demonstration that but little time is gained. The two tests, therefore, are of nearly equal value as regards early diagnosis. In cases in which the appearance of the Widal reaction is delayed, examination of the stools may render an earlier diagnosis possible. The examination of the feces for typhoid bacilli requires a complete laboratory, however, and much expert knowledge. It is entirely impracticable for the practicing physician and is not, so far as I know, carried on as a routine practice by any laboratory in this country.

EXAMINATION OF THE URINE FOR TYPHOID BACILLI.

Typhoid bacilli can be found in the urine in from 20 to 30 per cent. of all cases of typhoid at some time during their course. As a rule, they do not appear in the urine until late in the disease and after the appearance of the Widal reaction in the blood. In a few cases, however, they have been demonstrated in the urine before the appearance of the Widal reaction. The isolation of the bacilli is less difficult than from the stools, requiring as a rule but two days. The limitations to the use of this test are otherwise the same as in the case of the stools.

EXAMINATION OF THE BLOOD FOR TYPHOID BACILLI.

Typhoid bacilli have been cultivated from the splenic blood, obtained by puncture, by several observers. This test is deserving of no consideration, however, as puncture of the spleen is an entirely unjustifiable procedure.

Recent observers have cultivated the typhoid bacillus from blood taken from the veins of the arm in 80 per cent. of cases of typhoid. They used from 15 to 20 cubic centimeters of blood. The results of other observers have been much less satisfactory. There are no data as to the time at which the bacilli first appear in the blood, and hence no means of comparing the value of this method with that of the Widal reaction. This test requires a complete laboratory, technical knowledge, and at least two days. At best it is positive in only 80 per cent. of the cases, compared with 95 per cent. of positive reactions with the Widal test. Tapping a vein is a much more serious and delicate procedure than pricking the ear. The results, therefore, hardly justify the employment of this more complicated method in preference to the Widal reaction except in exceptional cases, even if it were not impracticable for the practitioner.

EXAMINATION OF THE ROSE SPOTS.

Several observers have recently succeeded in cultivating the typhoid bacilli from the rose spots. If five or six spots were excised positive cultures were obtained in almost every case. This method is not applicable before the appearance of the rose spots or in those rare cases in which there is no eruption. In a series of com-

parative tests the bacilli were almost invariably obtained from the rose spots before the appearance of the Widal reaction. They were found, on the average, six days before the Widal reaction. The bacteriologic examination of the rose spots seems, therefore, to be almost, if not quite, as accurate a method of diagnosis as the Widal test and, moreover, to afford a positive diagnosis earlier in the course of the disease. The excision of several rose spots under aseptic precautions is, however, a somewhat complicated procedure. Moreover, the cultures must be made at once and must be examined by an expert with a laboratory at his command. These facts render this method, accurate as it may be, of but little value to the practitioner.

THE DIAZO-REACTION.

This test is a simple one and can be easily performed by any one with but little expenditure of time. The diazo-reaction can be obtained in the urine of most cases of typhoid, often appearing early. As it is also often present in tuberculosis, septicemia and other febrile diseases, the very conditions with which typhoid is most easily confounded, it is of but comparatively little value.

THE LEUCOCYTE COUNT.

The number of white corpuscles in the peripheral circulation is subnormal throughout the course of typhoid. The average number per cubic millimeter is 5000 or even less. Any increase above the normal number means some complication, such as hemorrhage or inflammation, or some outside influence, such as cold bathing. Together with the diminution in the number of white corpuscles there is a relative increase in the proportion of mononuclear cells.

The estimation of the number of white corpuscles is a simple procedure, requiring but little time or experience, and is a perfectly practicable method of diagnosis for every one. It affords information of but little less value than the Widal reaction. The differential count of the white corpuscles is hardly more difficult but is rarely necessary.

In using the leucocyte count in the diagnosis of typhoid fever from other conditions, it must be remembered that the general rule applies here as elsewhere, namely, that the white count is only of value when the diagnosis lies between two diseases, one of which has a leucocytosis and one of which has not. For example, the white count is of no value in the differential diagnosis between typhoid and malaria, neither of which has a leucocytosis. On the other hand it is of great value in the diagnosis between typhoid and septicemia, the latter having a leucocytosis while typhoid does not. The diagnosis of typhoid, in the absence of any inflammatory complication or hemorrhage, must be viewed with much suspicion if the white count is above 10,000. The diagnosis of typhoid, made in the face of an inexplicable leucocytosis, is almost invariably wrong. An increase in the number of white corpuscles, in a case known to be typhoid, shows the presence of some complication, usually inflammatory. The information thus afforded should never be disregarded. The white count promises to afford a means of early diagnosis of perforation or even of threatened perforation. If, in addition to the pain in the abdomen, there is a progressive increase in the number of white corpuscles an actual or threatened perforation is almost certainly present.

TESTS WHICH SHOW THE PRESENCE OR ABSENCE OF OTHER DISEASES.

Although subordinate to the tests which demonstrate the existence of typhoid fever, those which reveal the presence or absence of other diseases are of great value in differential diagnosis. The examination of the blood for the plasmodium malarie when there is a question of malaria and lumbar puncture when there is a question of meningitis are among the most important. The value of the leucocyte count has been already mentioned. Other tests will suggest themselves in special cases.

CONCLUSIONS.

Laboratory and clinical methods of diagnosis should not be separated. Both are important and both fallible. Neither is complete without the other. Laboratory methods must be used with due recognition of their value and of their limitations. Otherwise they may lead to erroneous conclusions. The Widal reaction and the leucocyte count are the most useful of the laboratory tests for the diagnosis of typhoid fever. They are easily carried out and within the reach of all. Certain other methods are of equal or even greater value but are impracticable for everyday use.

REFERENCES.

- Curschmann: München. Med. Woch., Nov. 28, 1899.
 Elsner: Zeitschr. f. Hygien., 1895, xxi, p. 25.
 Gwyn: Phila. Med. Jour., 1900, v, p. 519.
 Morse: Boston Med. and Surg. Jour., 1899, cxi, p. 36; Archives of Pediatrics, 1900, xviii, 881; Ibid., 1901, xix, 338.
 Neufeld: Zeitschr. f. Hygiene, 1899, xxx, 3, p. 499.
 Piorkowski: Centralbl. f. innere Med., April 29, 1899.
 Remy: Annales de l'Institut Pasteur, 1900, xiv, p. 555.
 Richardson: Boston Med. and Surg. Jour., 1897, cxxvii, p. 133; Phila. Med. Jour., March 3, 1900.
 Schottmüller: Deutsche Med. Woch., 1900, xxvi, p. 511.
 Thayer: Jour. of the Boston Soc. of Med. Sci., 1900, v, p. 23.

THE TREATMENT OF TEMPERATURE BY DRUGS.

EDWIN ROSENTHAL, M.D.
 PHILADELPHIA.

Collective statistics of typhoid fever in children have placed the mortality rate at or near 5 per cent. Regarding treatment, no especial records have ever been presented, except those that may be questioned by reason of their representing certain plans of treatment, concerning which the profession has but little knowledge. Hence, as now constituted, it is impossible to substantiate certain classic methods by unquestioned results, and to arrive at some accurate conclusion, the statistical records of this disease in the adult should be considered for comparison.

As a diagnostic implement in the hands of the physician, the thermometer retains the first place, and, by its constant use, much reliance has been placed on what it shows, and its value remains unquestioned. By its use alone, we are very often enabled to diagnose typhoid fever, when many of the specific signs are late, or are not marked so as to be easily recognized.

THE AVERAGE DURATION OF THE FEBRILE MOVEMENT IN CHILDREN.

When using drugs in typhoid fever it is always well to know for what purpose. Hence, when giving drugs for fever this is a pertinent question: Is it given for the disease pure and simple, or for a complication? In young children the fever often lasts only from 8 to 14 days, according to Morse. The average course under 10 years is 19.3 days, and from 10 to 15 years, 22.6