perineal prostatectomy after removing the mucous membrane, but it is an extraordinary thing to open the bladder, pass in the hand and rip out everything and then expect nothing wrong to happen. He saw one case operated on suprapubically, and there was absolute stricture at the neck of the bladder, admitting only a filiform bougie.

The mortality is the point on which the whole thing should hinge. These old people should not be treated for the sake of usefulness to the surgeon, but because he wants to make them bless him as long as they live. Dr. Ferguson said he would rather have 5 per cent. of perineal fistulas than have 5 per cent. of his patients in the grave. Tuffier collected 2,227 cases with only 6 per cent. of mortality by the perineal route. Tenney and Chase of Boston collected 2,342 cases operated on by the perineal route and 667 operated on by the suprapubic route, with a mortality of 7 per cent. on the one hand and 13.2 per cent. on the other.

GANGOSA.

O. J. MINK, M.D. AND N. T. McLEAN, M.D.

Assistant Surgeons, U. S. Army.

U. S. NAVAL STATION, GUAM.

DEFINITION.—Gangosa, a Spanish word meaning muffled voice, is the name employed by the Spaniards in the Ladrone and Caroline Islands to describe a disease characterized by a destructive ulceration, usually beginning on the soft palate, pillars or uvula, and extending by continuity to the hard palate and nasal cavity, larynx, and even to the face. Active ulceration is followed after a variable period by cicatrization or chronic ulceration. Mutilation always results. Constitutional symptoms are either slight or absent.

SYNONYMS.—*Rhinopharyngitis mutilans* (Leys). *Ogo* (Chamorro).

HISTORY AND GEOGRAPHY.

There is absolute proof that gangosa has existed in the Island of Guam for at least 150 years, and probably much longer. In 1828 the Spanish Royal Commission sent to investigate conditions in the Ladrone Islands reported gangosa as being prevalent and recommended that these cases and cases of leprosy and large ulcerations be isolated on one of the smaller islands of the group. This is the earliest official recognition of the existence of the disease and the name gangosa. From these sources it is evident that the disease showed the same characteristics as at the present time. The known distribution of gangosa is somewhat indefinite. It exists throughout the Ladrone and Caroline islands. The case from Panama, recently reported by Fordyce and confirmed by Arnold, is undoubtedly true gangosa.¹

Surgeon J. F. Leys,² United States Navy, has carefully studied the bibliography and mentions reports of rhinopharyngeal ulcerations in Fiji, British Guiana, Jamaica, Italy, Dominica and Nevis. Inasmuch as Dr. Leys, who is familiar with the disease as it exists in Guam, has recently discussed the cases found in Dominica and Nevis with Dr. J. Numa Rat, the recorder of these cases, the disease undoubtedly exists in the West Indies as well as in Polynesia.

ETIOLOGY.

General Prevalence.—During the past year 125 cases have been examined. Many of these were brought under compulsion by the authorities. It is certain that the work of the past year did not include all the cases, as many not yet apprehended are known to exist in various parts of the island. A conservative estimate places the total number between 200 and 250. The population of the island being about 11,000, the ratio is 18 to 22 per 1,000. As the natives of the various parts of the island are intimately associated by religious, marital and com-

mercial relations, no topographic distinctions can be made.

Season.—The limited number of cases seen during the period of invasion makes it impossible to give any definite statement concerning the effect of season.

Age.—In a series of 80 cases the ages at the time of invasion were as follows:

First decade	2
Second decade	38
Third decade	23
Fourth decade	13
Fifth decade	2
Sixth decade	2

The youngest ages at invasion were 3 and 9 years; the oldest were 54 and 59 years. From these figures it appears that the majority of the cases start during the second and third decade.

Sex.—In the above series of 80 cases, 49 were in females and 31 in males.

Immunity.—No cases have been observed or recorded in whites. Cases in persons of mixed white and native blood are infrequent. The great majority of all cases appear in the pure-blood native. The absolute and relative immunity in the white and mixed races respectively may be explained by the superior hygienic and dietetic surroundings of these races.

Heredity.—There is no evidence showing that heredity plays any part in the transmission of the disease.

Food.—The native theory and that held by the Spaniards gives food a prominent place in the causation of the disease. They believe that such foods as decomposed or very salty fish, especially if eaten raw, uncooked taro and native pepper are the direct causes of the disease. This theory seems improbable. The eating of tainted fish is not more frequent than in any other fish-eating community, and when eating the usual symptoms of ptomain poisoning occur. Cases of gangosa rarely give a history of such poisoning. The eating of very salty fish, raw taro and native pepper is not sufficiently frequent to explain the large number of cases. Furthermore, cases are seen in natives eating the same food as the whites.

A Manifestation of Syphilis.—The disease has been considered a late manifestation of syphilis. During the past year we have not seen syphilis of any type among the native population and do not believe it exists. Our observations agree with those of Leys,³ who states:

It (gangosa) appears in healthy and well-developed persons of all ages, with no signs of hereditary syphilis in their own persons or in their brothers and sisters; and with no signs of syphilis in their parents. . . . Acquired syphilis is a common disease in most races over nearly the whole world. It is an extremely rare disease here, and neither primary nor secondary syphilis has been seen in a native during the past year, among thousands of persons treated for other diseases, including several prostitutes. This disease is common here, and rare or unknown where syphilis is common. Dr. Daniels, who was in Fiji for years, states that there was no syphilis in Fiji at a time when the lesions in this disease were common. The appearance of the primary lesion of this disease in otherwise healthy children of healthy parents, at 3, 4 and 9 years of age, excludes acquired syphilis.

A Sequel of Yaws.—Gangosa has been described as a sequel of yaws. Only one of our cases gave a history of this disease. While gangosa is limited to a few localities, yaws is a common disease in most tropical countries. We see no reason to consider it as an etiologic factor.

Specific Infecting Agent.—Our observations lead us to believe that there is a specific infecting agent, the nature of which is as yet undetermined. It is believed

1. Fordyce: Jour. Cut. Dis., January, 1906.

2. Leys: Jour. Trop. Med., London, Feb. 15, 1906.

3. Leys: Report of the Surgeon-General, U. S. Navy, 1905, p. 93.

that in the transmission of the disease flies are an important factor. The lesions of gangosa are so exposed that flies are attracted in great numbers, and, while feeding on the exudations, their legs, wings and bodies are contaminated. They in turn may infect the patient directly or indirectly through the food. The use in common by all members of the family of such articles as towels, clothing, pipes, cigars and betel-nut, with absence of dietetic hygiene, renders direct infection probable. The individual vital resistance is lowered by overcrowding, 15 to 20 persons, as a rule, sleeping in a small, unventilated hut. The resistance being lowered by such environment, the disease is easily transferred.

Inoculation.—No experiments have been made or cases seen which would indicate that infection by inoculation occurs.

MORBID ANATOMY.

As the disease is not fatal, autopsy is only possible when death results from intercurrent diseases. We have had but one case for autopsy:

Autopsy.—A male of 18 years. Clinical diagnosis; native epidemic asthma. Well developed and nourished, small muscular build. Nose and upper lip were entirely destroyed and replaced by triangular opening leading to the pharynx. Bony parts forming roof of mouth absent. Cuticle between eyes and inferior maxilla was a mass of superficial scar tissue. Tongue was normal. Tonsils and pillars were replaced by scar tissue. Larynx was normal. Eyes were normal. Brain was not examined. Heart was hypertrophied. Pericardium was normal. Pleura firm, adhesions right apex and left base and apex.

Lungs: Right upper lobe firmly adherent, middle and lower lobes dark in color; bronchioles contain thin purulent fluid. Left lung firmly adherent at apex and base, lower lobe dark in color, bronchioles contained thin purulent fluid. Left pneumothorax. Liver: Large, pale and mottled in appearance (a condition frequently seen at autopsy here). Gall bladder filled with dark bile. Kidneys normal. Stomach distended but normal. Intestines normal, contained large numbers of ascaris, hook and whip worms. Pancreas in consistency increased.

Cause of death: Native epidemic asthma. Sections of gangotic area were sent to the United States for examination.

PATHOLOGIC HISTOLOGY.

Fordyce states that the lesion was a granuloma, the nature of which could not be determined by the methods employed. It could be differentiated from others of this class, like blastomycosis, actinomycosis, rhinoscleroma and lepra, by the absence of their specific microorganisms; from mycosis fungoides by the character of the infiltrate and the absence of fragmentation. The histologic picture could be readily mistaken for tuberculosis, as the giant cells were numerous, with nuclei arranged peripherally. Many independent foci containing three or four cells were encountered deep in the corium.

SYMPTOMS.

Onset.—Only three cases have been observed from the earliest stages of the disease. The majority of patients present themselves for treatment only after mutilation is marked. In the three cases above mentioned the initial symptoms led to a diagnosis of tonsillitis, pharyngitis and laryngitis of mild degree. The patients were between the ages of 12 and 15 years and were in good physical condition. Prostration was slight or absent. All showed a slight rise of temperature and complained of soreness in the posterior nares and pharynx, with stiffness of the muscles of deglutition. In one case a typical acute coryza was present. Inspection at this time showed mild congestion of tonsils, pillars, soft palate and uvula.

During the first week the general condition was practically unchanged.

Local Symptoms.—The throat symptoms became localized, and on the third day a patch of yellowish-gray membrane was observed on the soft palate in the first case, the uvula in the second and the right pillar in the third case. The membrane was elevated, thick and extremely tenacious. On removal the denuded surfaces bled freely. Within twenty-four hours of the appearance of the membrane the typical ulceration was established. On the area covered by the membrane a number of small depressions appeared. The ridges between the depressions and the membrane were rapidly absorbed. The ulcer, now about one-half inch in diameter, had a punched-out appearance, with undermined edges and a deep, uneven floor, covered with a yellowish-white, very offensive discharge. It was surrounded by a zone of inflammation about one-fourth inch in width. The depth of the ulcer rapidly increased until, in the case of the uvula and soft palate, the tissues were perforated by the seventh day. These cases are still under observation.

Course of the Disease.—After the first week no marked changes occur in the character of the ulcer. It progresses steadily, the rapidity varying with the individual case, destroying the bony and soft parts with equal ease. This active stage may continue indefinitely or may become arrested at any period. The average duration in those cases of our series, which later became quiescent, was about two years. The active stage in those cases varied from one to seven years in duration.

In seven cases, which have never been quiescent, the disease has progressed slowly, but steadily, the shortest having continued for ten and the longest for thirty-five years. At any time during quiescence the disease may again become active. This activity may continue indefinitely or pass into a second stage of quiescence. These alternate periods of activity or quiescence may occur repeatedly. When inactive the zone of inflammation is replaced by a more or less perfectly scarred area, from which an abundant and very offensive discharge pours. At no time during either stage is the patient's general health materially affected.

Fulminating Gangosa.—This type occurs in children under 5 years of age, and all cases seen have been in gangotic families. In some instances two or three children in the same family have been attacked. The sudden onset, extreme prostration, extensive membrane, marked cervical adenitis and rapidly developing toxemia and dyspnea give a picture closely resembling diphtheria. Surgeon C. P. Bagg, United States Navy, states that cultures from the throats were negative for Klebs-Loeffler bacilli. Death occurs from toxemia rather than from dyspnea. The great majority of these cases prove fatal within forty-eight hours, but if the patient survives this period the typical mutilation rapidly develops and the case follows the usual course.

COMPLICATIONS AND SEQUELÆ.

All complications result from the direct extension of the disease and vary in severity with the extent of ulceration. The principal parts which may be involved are the nose, hard palate and superior maxillæ, larynx, eyes, face and teeth.

Nose.—The extensive destruction of the deeper parts may totally destroy this organ. The septum and turbinates are early attacked. The supporting structure is destroyed, the external soft parts sink and later may be

TABLE OF CASES.

General History.			History of Disease. (Years.)		Present Condition.						
Age and Sex.	Family History.	Duration.	Active.	Quiescent.	Tonsils and Pillars.	Uvula and Soft Palate.	Hard Palate.	Larynx.	Nose.	Eyes.	General Condition.
1 35F.	Good. Only case.	30	20	10	Destroyed.	Destroyed.	Perforated.	Unaffected.	Flattened and septum perforated.	Unaffected.	Excellent.
2 83F.	Good. Only case.	2	1 1/2	1/4	Unaffected.	Very slightly affected.	Unaffected.	Unaffected.	Flattened and septum perforated.	Unaffected.	Excellent.
3 46F.	Good. Only case.	13	13		Slightly affected.	Unaffected.	Unaffected.	Unaffected.	Destroyed.	Unaffected.	Excellent. Deaf 13 years.
4 45M.	Good. Only case.	8	8		Destroyed.	Destroyed.	Badly necrosed.	Badly ulcerated.	Destroyed.	Destroyed.	Excellent. Voice hoarse and totally blind.
5 36M.	Good. Only case.	10	1 1/2	9 1/2	Slightly affected.	Destroyed.	Perforated.	Badly ulcerated.	Septum destroyed.	Unaffected.	Excellent. Voice hoarse.
6 42M.	Good. Only case.	22	20	20	Destroyed.	Destroyed.	Badly scarred.	Unaffected.	Flattened.	Unaffected.	Excellent.
7 47M.	Bad. Other cases.	29	2	27	Destroyed.	Destroyed.	Badly scarred.	Unaffected.	Partially destroyed.	Unaffected.	General muscular paralysis.
8 33M.	Bad. Father, tbc.; mother, gangosa.	4	2	2	Slightly affected.	Uv. destroyed. S. affected.	Perforated.	Badly ulcerated.	Flattened.	Unaffected.	Excellent. Voice absent.
9 22F.	Only case. Father, tbc.	5	2	3	Destroyed.	Destroyed.	Perforated.	Unaffected.	Flattened.	Unaffected.	Excellent.
10 12M.	One on her case. Sister, tbc.	27	1	28	Destroyed.	Destroyed.	Perforated.	Unaffected.	Flattened.	Unaffected.	Fair. Cervical adenitis.
11 17F.	Bad. Other cases.	17	1	16	Unaffected.	Unaffected.	Slight active ulceration.	Unaffected.	Destroyed.	Unaffected.	Fair. Large ulcer L. F. A.
12 19M.	Only case. Father, tbc.	17 1/2	1/2	1/2	Unaffected.	Very active ulceration.	Unaffected.	Slight ulceration.	Unaffected.	Unaffected.	Excellent.
13 26F.	Bad. Mother, tbc.; father, gangosa.	7	5	2	Slightly affected.	Unaffected.	Unaffected.	Slight ulceration.	Flattened.	Unaffected.	Excellent. Voice hoarse.
14 42M.	Good. Only case.	23	3	20	R. destroyed.	Right side destroyed.	Right side perforated.	Voc. cords scarred.	Unaffected.	Unaffected.	Excellent. Voice absent.
15 40M.	Good. Only case.	12	1	11	Destroyed.	Destroyed.	Perforated.	Slight ulceration.	Destroyed.	Right eye destroyed.	Fair. Healed lupus scars.
16 19F.	Unobtainable.	8	1	1	Destroyed.	Uv. destroyed. S. affected.	Unaffected.	Unaffected.	Unaffected.	Left eye destroyed.	A leper.
17 19F.	Good. Only case.	9	1 1/2	7 1/2	Destroyed.	Destroyed.	Perforated.	Slight ulceration.	Septum destroyed.	Active destruction.	Poor. Lupus neck, face and chest.
18 37F.	Good. Only case.	34	10	24	Unaffected.	Partially destroyed.	Perforated.	Slight ulceration.	Destroyed.	Active destruction.	Excellent.
19 21F.	Good. Only case.	14	7	7	Unaffected.	Destroyed.	Unaffected.	Unaffected.	Unaffected.	Unaffected.	Excellent.
20 46F.	Bad. All affected.	35	35		Destroyed.	Destroyed.	Extensive necrosis.	Badly ulcerated.	Totally destroyed.	L. destroyed. R. ulcerated.	Good, except asthma. Voice hoarse.
21 22M.	Bad. All affected.	5	5		Destroyed.	Destroyed.	Perforated.	Badly ulcerated.	Totally destroyed.	Unaffected.	Excellent. Voice hoarse.
22 21F.	Only case. Father, tbc.	13	13		Destroyed.	S. P. unaffected.	Unaffected.	Badly ulcerated.	Posterior ulceration.	Unaffected.	Excellent.
23 17F.	Uncle, gangosa.	7	4	3	Badly scarred.	Uv. destroyed. S. affected.	Unaffected.	Unaffected.	Destroyed.	Unaffected.	Excellent.
24 21F.	Good. Only case.	11	11		Destroyed.	Destroyed.	Extensive necrosis.	Badly ulcerated.	Entirely destroyed.	Lower lids destroyed.	Poor. Large ulcers L. F. A.
25 30M.	Bad. All affected.	14	2	12	Destroyed.	Partially destroyed.	Extensive necrosis.	Unaffected.	Flattened.	R. lower lid destroyed.	Good.
26 21M.	Bad. All affected.	6	6		Destroyed.	Destroyed.	Extensive necrosis.	Slight ulceration.	Destroyed.	Unaffected.	Good. Voice hoarse.
27 21M.	Only case. Cousin, tbc.	2	2		Destroyed.	Uv. destroyed. S. affected.	Unaffected.	Badly ulcerated.	Unaffected.	Unaffected.	Excellent. Voice almost gone.
28 56M.	Fair. Cousin, gangosa.	40	2	38	Destroyed. L.	Uv. destroyed. S. affected.	Perforated.	Badly ulcerated.	Partially destroyed.	Unaffected.	Excellent. Voice absent. Deaf 26 years.
29 20F.	Only case. Father and brother, lepers.	10	4	6	Destroyed.	Slightly affected.	Perforated.	Slight ulceration.	Flattened.	Unaffected.	Fair. A leper.
30 37F.	Mother, gangosa.	7	2	5	Destroyed.	Partially destroyed.	Perforated.	Slight ulceration.	Partially destroyed.	Unaffected.	Fair. A leper.
31 62M.	Good. Only case.	50	2	48	Destroyed.	Destroyed.	Perforated.	Slight ulceration.	Partially destroyed.	Unaffected.	Excellent.
32 34F.	Mother, tbc.; father, gangosa.	28	1	27	Destroyed.	Destroyed.	Perforated.	Unaffected.	Partially destroyed.	Unaffected.	Excellent.
33 38F.	Good. Only case.	10	1 1/2	9 1/2	L. T. destroyed.	Slightly ulcerated.	Perforated.	Unaffected.	Partially destroyed.	Unaffected.	Excellent.
34 40F.	Good. Only case.	7	7		Destroyed.	Slight ulcerations.	Unaffected.	Slight ulceration.	Very slight ulceration.	Unaffected.	Excellent.
35 46F.	Good. Only case.	21	21		Destroyed.	Destroyed.	Perforated.	Slight ulceration.	Septum destroyed.	Unaffected.	Excellent. Voice hoarse.
36 38F.	Good. Only case.	22	1	21	Destroyed.	Destroyed.	Perforated.	Slight ulceration.	Unaffected.	Unaffected.	Excellent. Voice hoarse.
37 15F.	Good. Only case.	6	6		Unaffected.	Uv. slightly ulc.	Unaffected.	Slight ulceration.	Unaffected.	Unaffected.	Excellent. Voice hoarse.
38 36F.	Good. Only case.	12	1	11	Unaffected.	Unaffected.	Perforated.	Unaffected.	Partially destroyed.	Unaffected.	Excellent.
39 21F.	Poor. Father, tbc.; mother, gangosa.	9	2		Slightly affected.	Slightly affected.	Perforated.	Unaffected.	Flattened and septum destroyed.	Unaffected.	Good. Scars of old ulcer, large joints, chest and neck.
40 37F.	Good. Only case.	24	14	10	Destroyed.	Destroyed.	Perforated.	Unaffected.	Flattened.	Unaffected.	Fair. Many scars of old ulcers.
41 41F.	Fair. Mother, tbc.	12	2	10	Destroyed.	Destroyed.	Perforated.	Unaffected.	Destroyed.	L. destroyed.	Poor. A leper.

TABLE OF CASES—Continued.

General History.		History of Disease. (Years.)		Present Condition.							General Condition.	
Age and Sex.	Family History.	Duration.	Active.	Quiescent.	Tonsils and Pillars.	Uvula and Soft Palate.	Throat.	Larynx.	Nose.	Eyes.		
42 18F.	Good. Only case. Mother a leper.	2	1	1	Destroyed.	Destroyed.	Perforated.	Slight ulceration.	Flattened.	R. lower lid ulcerated.	Good. Voice hoarse.	
43 15F.	Good. Only case.	4	2	2	Destroyed.	Destroyed.	Scarred.	Unaffected.	Destroyed.	R. lower lid ulcerated.	Excellent.	
44 12F.	Good. Only case.	2	1	1	Destroyed.	Destroyed.	Destroyed.	Unaffected.	Destroyed.	L. lower lid ulcerated.	Excellent. Feeds through nasal opening.	
45 40M.	Good. Only case.	24	9	15	Destroyed.	Destroyed.	Destroyed.	Unaffected.	Destroyed.	L. destroyed. R. unaffected.	Excellent.	
46 21M.	Good. Only case.	6	1	5	Unaffected.	Unaffected.	Unaffected.	Unaffected.	Destroyed.	Unaffected.	Excellent.	
47 29M.	Good. Only case.	9	4	5	Destroyed.	Uvula destroyed.	Perforated.	Unaffected.	Destroyed.	Unaffected.	Excellent.	
48 48M.	Good. Only case.	22	1	21	Destroyed.	S. P. unaffected.	Perforated.	Unaffected.	Flattened.	Unaffected.	Excellent.	
49 39F.	Good. Only case.	6	1	5	Destroyed.	Destroyed.	Perforated.	Unaffected.	Destroyed.	Destroyed.	Excellent.	
50 14F.	Good. Only case.	4	1	3	Unaffected.	Uvula unaffected.	Unaffected.	Unaffected.	Unaffected.	Unaffected.	Excellent.	
51 59M.	Good. Only case.	22	13	9	Scarred.	Destroyed.	Perforated.	Unaffected.	Destroyed.	Destroyed.	Poor. A leper.	
52 11F.	Bad. Four other cases.	4	3	1	Destroyed.	Uvula unaffected.	Perforated.	Unaffected.	Destroyed.	Unaffected.	Excellent.	
53 33F.	Bad. Four other cases.	8	1	7	Destroyed.	S. P. perforated.	Perforated.	Slight ulceration.	Destroyed.	Unaffected.	Voice hoarse.	
54 14M.	Bad. Three other cases.	1	1	1	Destroyed.	Uvula unaffected.	Perforated.	Unaffected.	Unaffected.	Unaffected.	Excellent.	
55 38F.	Fair. Only case. Two cases of leprosy.	1	1	1	Destroyed.	Uvula unaffected.	Scarred.	Slight ulceration.	Flattened.	R. lower lid ulcerated.	Good.	
56 30F.	Good. Only case.	1	1	1	Destroyed.	S. P. scarred.	Scarred.	Unaffected.	Filled with adenoid growth.	Unaffected.	Fair. Mass of adenoid tissue protrudes from nose to right cheek.	
57 30F.	Good. Only case.	12	2	10	Destroyed.	Destroyed.	Perforated.	Unaffected.	Flattened.	Unaffected.	Good.	
58 45F.	Bad. Two other cases.	10	2	8	Destroyed.	Destroyed.	L. side destroyed.	Slight ulceration.	Unaffected.	Both l. lids ulcerated.	Voice hoarse.	
59 22M.	Bad. Mother gangosa.	4	2	2	Destroyed.	Scarred.	Perforated.	Unaffected.	Flattened.	Unaffected.	Excellent.	
60 60F.	Good. Only case.	1	1	1	Destroyed.	Destroyed.	Active ulceration.	Slight ulceration.	Flattened.	Unaffected.	Voice hoarse. Very active.	
61 15M.	Bad. Mother dead of gangosa.	2	1	1	Destroyed.	Active ulceration.	Active ulceration.	Unaffected.	Flattened.	Unaffected.	Good.	
62 20M.	Good. Only case.	2	1	1	Destroyed.	Destroyed.	Destroyed.	Unaffected.	Sunken. Three-lobed anterior opening.	Unaffected.	Good. Large ulcer on left leg.	
63 12M.	Bad. Uncle, gangosa.	2	1	1	Destroyed.	Destroyed.	Perforated and scarred.	Slight ulceration.	Sunken. Two-lobed.	Unaffected.	Good. Voice hoarse.	
64 21M.	Good. Only case.	1	1	1	Unaffected.	Unaffected.	Unaffected.	Unaffected.	Septum destroyed.	Unaffected.	Good. All internal nasal structure destroyed. Nose inflamed.	
65 57M.	Good. Only case.	12	1	11	Destroyed.	Destroyed.	Scarred.	Unaffected.	Partially destroyed.	Unaffected.	Excellent.	
66 28M.	Good. Only case.	1	1	1	Scarred.	Destroyed.	Scarred.	Unaffected.	Unaffected.	Unaffected.	Excellent.	
67 26F.	Bad. Brother, gangosa.	1	1	1	Unaffected.	Destroyed.	Destroyed.	Slight ulceration.	Unaffected.	Unaffected.	Voice hoarse.	
68 15M.	Bad. Sister, gangosa.	1	1	1	Destroyed.	Destroyed.	Perforated.	Unaffected.	Flattened and sunken.	Unaffected.	Excellent.	
69 29M.	Good. Only case.	5	1	4	Destroyed.	Destroyed.	Destroyed.	Slight ulceration.	Destroyed.	Unaffected.	Voice hoarse.	
70 47M.	Bad. Two other cases.	27	1	26	Destroyed.	Destroyed.	Almost destroyed.	Unaffected.	Destroyed.	Lower lids destroyed.	Good. Upper lip absent. Face badly scarred.	
71 49F.	Good. Only case.	25	1	24	Destroyed.	Destroyed.	Perforated.	Slight ulceration.	Flattened and sunken.	Unaffected.	Good. Voice hoarse.	
72 21F.	Good. Only case.	1	1	1	Destroyed.	Scarred.	Scarred.	Slight ulceration.	Scarred, flattened and drawn to the right.	Unaffected.	Good. Mouth badly scarred. Voice hoarse.	
73 27M.	Good. Only case.	3	1	2	Destroyed.	Right side of S. P. destroyed.	Perforated.	Unaffected.	Flattened. L. nostril partially destroyed.	Unaffected.	Good.	
74 47F.	Good. Only case.	16	1	15	Destroyed.	Destroyed.	Scarred and perforated.	Unaffected.	Sunken. Tip turned upward.	L. lower lid destroyed.	Good. Face badly scarred.	
75 18M.	Good. Only case.	6	1	5	Destroyed.	Destroyed.	Scarred.	Unaffected.	Destroyed.	L. lower lid destroyed.	A leper. Face badly scarred.	
76 13F.	Bad. Three other cases.	1	1	1	Pillars scarred.	Scarred.	Scarred.	Extensive ulceration.	Unaffected.	Unaffected.	Good. Very hoarse voice.	
77 12F.	Only case. Mother, tbc.	2 1/2	12	12	Unaffected.	Destroyed.	Perforated.	Extensive ulceration.	Unaffected.	Unaffected.	Fair. Leper. Very hoarse voice.	
78 20F.	Good. Only case.	1	1	1	Scarred.	Scarred.	Scarred.	Unaffected.	Destroyed. Active ulceration.	Unaffected.	A leper. Face badly scarred.	
79 44F.	Good. Only case.	6	1/2	5 1/2	Destroyed.	Destroyed.	Destroyed.	Unaffected.	Destroyed.	Both lids destroyed.	A leper. Face badly scarred.	
80 41F.	Good. Only case.	1	1	1	Scarred.	Scarred and contracted.	Slightly scarred.	Unaffected.	Scarred externally.	Unaffected.	Good. Face badly scarred.	
81 19M.	Good. Only case.	2 1/2	12	1/12	Unaffected.	S. P. perforated to post-nares.	Unaffected.	Unaffected.	Unaffected.	Unaffected.	Excellent. (New case.)	

absorbed, the organ being replaced by an opening. The sense of smell is destroyed early in the course of the disease.

Hard Palate and Superior Maxillæ.—The hard palate may be perforated or entirely destroyed. The central portions of the superior alveolar arches are frequently

PROGNOSIS.

Except in the fulminating type, the disease *per se* is never fatal. The mutilations and complications are permanent and cause corresponding disability. Intercurrent diseases follow a typical course and are not severer than in those unaffected with gangosa.

Fig. 1.—Case 52. Extensive involvement with no external mutilation.

Fig. 2.—Case 62. Flattened nose.

Fig. 3.—Case 35. Notice extensive scarring of upper lip.

destroyed. This causes loss and deformity of the upper teeth. This deformity produces the typical "cleft palate voice" from which the disease takes its name.

Larynx.—In 81 cases the larynx was involved in 33. Involvement varies from slight ulceration to total destruction, and as a result the voice is husky or absent.

Face.—Ulceration extends peripherally from the oral and nasal openings or from the common opening when the upper lip is destroyed.

Eyes.—These become involved when facial ulceration destroys the lids. The eyes were involved in 21 cases. The sequence in destruction is conjunctivitis, corneal opacity, staphylomata and finally blindness.

Tongue.—The tongue and muscles of deglutition are unaffected.

Hearing.—Hearing was affected in only 2 cases. No sequelæ have been observed.

DIAGNOSIS.

To those who have ever seen a case of gangosa, diagnosis offers no difficulty. The primary rapid ulceration, consequent mutilation and subsequent chronic ulceration produce a picture seen during the active stage. During the quiescent stage the ulceration is replaced by scar tissue.

Differential Diagnosis.—Gangosa is differentiated from (1) leprosy, by the sudden onset with immediate localization, absence of fatal termination and absence of *B. lepræ*. (2) Lupus of the mucous membranes, by the sudden onset, local appearance of the throat symptoms, absence of tubercle bacilli and absence of tuberculosis elsewhere. (3) Syphilis, by the absence of history and symptoms and the failure of specific treatment. The disease bears no clinical resemblance to blastomycosis, actinomycosis, rhinoscleroma or mycosis fungoides.

Fulminating gangosa is diagnosed from diphtheria by the absence of Klebs Loeffler bacilli and by the typical mutilation in those cases which survive.

PROPHYLAXIS.

The contagiousness of the disease has been recognized by the Spaniards in Guam for almost a century. They established a colony for the segregation of these cases, which was discontinued at the time of the American

Fig. 4.—Case 38. Old scarring with labial destruction.

occupation. There is good evidence showing that this abandonment of segregation has increased the prevalence of the disease in Guam. In view of this fact, the governor of Guam, on April 9, 1906, issued the following order:

EXECUTIVE GENERAL ORDER NO. 111.

In view of the existence in this island of an infectious and contagious disease known as *Gangosa*, it is of paramount neces-

sity, in the interest of humanity and for the protection of society, to adopt proper measures to arrest the spread of that loathsome and dangerous malady. Less rigorous methods being either futile or of doubtful efficacy, it is hereby ordered and decreed that:

1. All persons within the Island of Guam who, on examination before a board consisting of all the medical officers of the

established, the unaffected members of the family being removed and kept under observation.

TREATMENT.

There is little doubt that treatment in the early stages limits the progress of the disease. The treat-

Fig. 5.—Case 45. Extensive skin involvement has destroyed left eye.

Fig. 6.—Case 42. Notice beginning of total destruction of nose.

Fig. 7.—Case 20. Very severe scarring; left eye destroyed.

Naval Station of the island ordered by the governor, are found to be suffering from *Gangosa*, shall, on the unanimous recommendation of said board, be segregated in the same manner and under the same conditions, rules and regulations as are now provided by General Order No. 43, in the case of lepers.

2. The place of segregation shall be a tract of land in the

Fig. 8.—Case 75. Notice co-existence of leprosy and gangosa.

locality of Ypao, district of Agna, to be known as the Ypao Colony, having an area of about six (6) hectares, the boundaries of which shall be constructed of convenient and suitable material and bear the sign Ypao Colony.

3. This order shall take effect on the eighteenth day of April, 1906.

T. M. PORTS, U. S. N., Governor.

In the fulminating type rigid quarantine should be

ment is essentially local and should aim to destroy the affected area. Tincture of iodine applied freely appears to be the best agent for this purpose. Lunar caustic, phenol and chromic acid have been used. It is believed that in some cases the actual cautery would be the most effective agent. An antiseptic mouth wash should also be used. Tonic treatment when indicated should be used. Potassium iodid even in large doses and long continued appears to have no effect on the disease. In chronic cases deodorants should be used. Potassium permanganate, in 1 per cent. solution has proved the most advantageous. When possible, complications should receive their appropriate treatment. It seems reasonable to believe that the x-ray, Finsen light and radiotherapy would be of service in the treatment of this disease, more especially in the chronic external ulcerations. In the fulminating type, treatment should be symptomatic, combined with active local disinfection.

We desire to express our thanks to Surgeon C. P. Bagg, United States Navy, for valuable suggestions and assistance in the preparation of this article, and to K. S. Hart, United States Navy, for careful photographic work.

THE BACTERIA IN SCARLATINAL AND NORMAL THROATS.*

GUSTAV F. RUEDIGER, M.D.
CHICAGO.

INTRODUCTION.

It is a well established fact that streptococci are constantly found in great abundance on the tonsils of scarlet fever patients. This was shown very clearly by the investigations of Weaver, Baginsky and Sommerfeld, Charlton, Charbade, Booker, Taugl, Klein, Kurth and

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Fifty-seventh Annual Session, June, 1906.

* This article is from the Memorial Institute for Infectious Diseases and describes work which was aided by a grant from the American Medical Association.