felt perfectly well. On the most careful examination I could find nothing else.

20th.—The same; but the boy could not understand being kept in bed when he felt perfectly well, and I could not sanction his getting up while his temperature and pulse were above normal.

21st.—After another careful and minute examination, owing to his temperature and pulse remaining the same, I detected a very soft and faint mitral systolic murmur.

This gradually increased in intensity every day, and on the 24th his temperature became normal, and continued so, but his pulse remained high for a time. He was kept quietly in bed for another ten days.

I carefully talked over the illness with his father, and told him he had endocarditis, which was a very serious matter; though by dint of the very greatest care, in not allowing any exertion for the next few months, its effects might all disappear; otherwise he would carry a maimed heart for the rest of his days.

On March 8th he was unfortunately suddenly told he might go home for the rest of the term. Next morning when I saw him I was informed that he had been in a curious state, talking incoherently about things that had no existence at all, and he seemed thoroughly dazed at my visit. He was cheerful and happy, though incoherent, and had a quiet pulse. Whether his joy was too much for his state, or whether minute particles of fibrine were liberated from the mitral valve in the excitement of his sudden joy at the idea of going home, I cannot say; but I am inclined to think that this latter is what happened.

On June 11th I auscultated his heart, and there was not the slightest trace of murmur remaining. He was quite well.

The lessons to be learnt from the case, I think, are—
1. That the most trivial amount of rheumatism may be accompanied by endocarditis. 2. That recent murmurs from endocarditis may and do disappear in young healthy subjects by dint of very great care while they are recent.

Rugby.

A CASE OF SUDDEN DEATH DURING ETHER ADMINISTRATION.

By ROBERT N. HARTLEY, M.R.C.S.

THE rarity of cases of sudden death during the use of ether as an anæsthetic is my excuse for entering somewhat fully into the following details.

The patient was a gentleman, aged sixty-six, who had for some weeks previously been the subject of symptoms which eventually culminated in complete obstruction of the bowels, the site of the obstruction being thought to be either in the lower part of the descending colon or the sigmoid flexure.

As the only procedure holding out any hope of prolonging life, it was decided, in accordance with the opinion of Sir James Paget and Mr. Pridgin Teale, and with the full consent of the patient, to open the bowel in the right lumbar region.

Before the commencement of the operation the patient was lying, or half-lying, in bed, with the head and shoulders well propped up with pillows, breathing regularly and without great effort. There was no complicating disease in heart or lungs, and the pulse, though rather quick and small, was not inordinately feeble. There was thus nothing to contraindicate the use of an anæsthetic, or to make one more than usually anxious as to its effect. Some artificial teeth worn by the patient had been previously taken out, and one of the pillows was now removed for the more easy adjustment of the inhaler.

Ether was administered by Clover's smaller inhaler, two ounces being the quantity contained in the ether-chamber. The patient took the ether without the slightest hesitation or resistance, moderately inflating the india-rubber bag with each expiration, and in two minutes was quite anæsthetic. He was now placed on his left side with the head lower, and in five minutes from the commencement of the administration of ether the line of incision was made by Mr. Teale. The patient was never allowed to become deeply "carbonised," and the inhaler was frequently withdrawn entirely. In ten minutes from the commencement of the operation, and

just as I was engaged in counting the carotid pulse, which was beating evenly and with a rather stronger impulse than before the operation, he showed an inclination to vomit, and ejected a quantity of brownish fluid smelling strongly of brandy. He then took one deep inspiration, and seemed as if inclined to vomit again; but his head sank back on the pillow, and he quietly died, making no further respiratory effort, except one short gasp during the performance of artificial respiration. Prompt exploration of the fauces failed to show any mechanical obstruction to the breathing, and in spite of an immediate resort to artificial respiration and all other available means for restoring animation, our efforts proved useless. An examination of the inhaler showed that a little more than an ounce of ether had been spent.

No post-mortem examination was made.

The interest of this case centres in the difficulty of defining the exact cause of death. The mode of death is one which unfortunately is not rare during the use of *chloroform*, as an anæsthetic, but is almost unknown while a patient is under *ether*.

There was no mechanical obstacle to respiration by the regurgitation of anything into the larynx during the act of vomiting. Nothing solid had been introduced into the stomach for seven or eight days previous to the operation, and the vomited material consisted entirely of thin fluid. Moreover, there was no paroxysmal or convulsive seizure such as would be likely to follow a sudden mechanical interference with the act of breathing.

Two circumstances may have contributed to the suddenly fatal result:—

First, the distension of the bowels possibly interfering to some extent with the descent of the diaphragm, and hence with easy respiration, though neither before nor during the administration of ether had the patient shown any considerable difficulty or distress of breathing. Mr. Teale regrets, however, that he did not at once complete the operation, and rapidly open the bowel, and thereby reduce one factor—the distension by gas.

Secondly, the undoubted tendency to death from asthenia in cases of abdominal obstruction. In this particular case, the lungs having been more or less emptied of air by the act of vomiting, the additional quantity of blood drawn into the heart and large vessels by the next inspiratory effort might easily have brought the feebly acting heart to a standstill. Mr. Clover suggests to me that by the eructation of fluid containing brandy the ether may have been mixed with alcoholic vapour, which would add to its narcotising effect, and thereby tend to increase the cardiac depression.

Brixton-road, S.W.

ABSCESS OF THE CEREBELLUM.

By W. A. BERRIDGE, M.R.C.S., L.S.A.

I TRUST a general practitioner may be allowed to put upon record the following brief notes of a case of abscess of the cerebellum, as a small token of his gratitude for the instruction he received from Dr. J. Hughlings Jackson, F.R.S., and for the care bestowed upon his clinical education by his former teacher and now esteemed friend, Dr. Stephen Mackenzie.

A. D—, aged seventeen, had suffered from time to time since he was two years old from "gatherings in his head," which "broke," and "smelled very bad." He went to school when five years old, but never got on well. The schoolmaster said "he was stupid, and would not learn." Still he learned his letters, and could hold his own in play with other boys. He had no particular illness, never had scarlet fever, and had no signs of congenital syphilis, but often suffered from headaches and the aforesaid "gatherings."

When first seen (August, 1879) he was tall, pale, and rather thin. He spoke as I have noticed patients with organic cerebral mischief generally do, his words beginning with a "jerk" and ending with a drawl; or, in other words, the first part of a word being sharp and rather acute, the latter part grave and always prolonged. His mother said, "I have sent for you because he has a cold in his head, and cannot smell." On further inquiry, "he had headaches, occasional retchings, and felt giddy"; but he could not describe his "giddiness." He had no "reel." He ate ravenously. Urine 1015, acid, no albumen, no sugar. Temperature normal. He had no paralysis. Pupils dilated, but responded to light.

On examination with the ophthalmoscope, "the edges of both discs were rather indistinct, but there was nothing definite." After some treatment he improved, and was lost sight of for several weeks. When seen again (December 1879) his smell had returned, but he had lost his sight; he was quite blind. The edges of the discs were now quite obscured, the vessels partly covered by exudation; that is to say, he had well-marked double optic neuritis. He vomited frequently, and complained of intense pain at the back of his head. He had now the three cardinal symptoms—headache, vomiting, and double optic neuritis; he had also a long history of ear-discharge; so there was no difficulty in saying he had an abscess in his brain, and that he would die, most likely suddenly.

There was no marked change, and he died quite suddenly on March 12th, 1880. Next day, assisted by my friend and partner, Mr. F. B. Hallows, I opened the head. The convolutions were pale, and somewhat flattened. Nothing was noticed until slicing the cerebellum. There was a large abscess involving nearly the whole of the right lobe of the cerebellum, and containing about three ounces of what a surgeon would call "laudable pus." Permission could only be obtained to examine the head.

Redhill.

NOTES OF A CASE OF LYMPHADENOMA.

By W. BERRY, M.R.C.S.E., L.R.C.P. & S. ED.

HONORARY SURGEON, ROYAL ALBERT EDWARD INFIRMARY, WIGAN.

ON December 10th, 1879, I was asked to see Patrick O'B—, twenty-six years of age, a collier, living in a small house with deficient hygienic surroundings, and was given the following

History.—In August, 1879, he met with an accident in the coal-mine, being injured between two tubs and thrown on his back across the tram-rails. He was able, however, to walk to the surgery of the colliery doctor, who prescribed a liniment for him, which he applied. In consequence of this accident he was off work for one month; then he returned, working only one day, when he stated that he sprained his back owing to his work being heavy. He sought advice this time at the Wigan Infirmary, where he was admitted some time in September. He was discharged on the 9th of December without any improvement, feeling, if anything, weaker than when he entered.

Physical examination.—When I saw him in the evening of Dec. 10th he was lying on his back in bed, having a sallow, pasty appearance, with some puffiness underneath the eyes; respiration somewhat short and hurried; pulse 96, feeble; tongue slightly furred, with slightly raised papillæ, and dryish, although he did not complain of much thirst; bowels slightly constipated, and appetite very poor; urine moderate in quantity, of high colour, with some lithates, but no albumen; respiration somewhat puerile, probably owing to its hurried character; no dulness on percussion; heart's sounds very feeble, especially second sound, which could not be heard distinctly, but no murmur. On examining his abdomen, in left iliac region was found a round hard swelling about the size of a large orange; this tumour was immovable, and seemed to increase in size at its base. The inguinal glands and femoral glands of both sides were hard and enlarged, but not painful. He complained of pain behind the abdominal tumour in his back, and attributed all his symptoms to the accident. I did not take his temperature, but the house-surgeon of the infirmary told me that when he was in that institution he had a temperature of 103° F. His wife informed me that he was strong and healthy previous to the accident, but after his death I learned he had not been in good health for some time. The man and his friends were anxious to have all his ailments put down to his accident, so that he might receive benefit from the Permanent Relief Society; hence the difficulty in obtaining a correct history and arriving at a true diagnosis.

I prescribed a mild aperient tonic for him, and ordered plenty of milk and nourishing food.

I saw him two and three times a week, but with no marked improvement in his symptoms. The pasty, puffy appearance of his face increased, his left leg and thigh became cedematous, bed-sores commenced to form over his hip and great trochanter, and the femoral glands enlarged. His breathing got more impeded, and debility much greater.

On Jan. 7th, 1880, the glands just above the clavicle on the left side were much enlarged and almost in one mass, and the posterior triangle of the neck also contained glands enlarged in size.

No history of syphilis could be made out. I came to the conclusion that we had a case of *lymphadenoma* to deal with; through neglect, however, a microscopic examination of his blood was overlooked, but the ænemic appearance of his face was sufficient evidence of impoverished blood. He gradually got worse and died on Jan. 18th.

Through the kindness of the coroner, I got an order for a post-mortem examination, owing to the deceased's relatives seeking to obtain pay from the Permanent Relief Society. On January 20th, at 8 P.M., sixty-two hours after death, with the aid of my assistant, Dr. O'Reilly, and my pupil, Mr. Richards, I made a post-mortem. The body was emaciated, with swellings of enlarged glands in the neck, inguinal regions, and left side of abdomen, œdema of the left leg, a small bed-sore over left trochanter major, hæmstatic congestion over the buttocks and lumbar vertebræ.—Head: On removing the calvaria, the dura mater was found to be slightly adherent, the brain-surface (convolutions) pale, the arachnoid filled with serous fluid; when cut into it showed the same absence of blood; serum was also found filling both lateral ventricles.—Chest: Lungs in a state of collapse, lying at the bottom of the thorax, the pleuræ being filled with fluid. Lungs, on removal, found to be adherent in places. Heart lying flaccid in the pericardium, which contained very little fluid; on opening its cavities very little dark-coloured blood was found in them; the valves were healthy.—Abdomen: The liver was prominently in view, extending over into the left hypochondriac region, being much enlarged, but healthy in appearance when cut into. The spleen weighed about six ounces, but did not appear diseased. The stomach was distended with gas, as were also the intestines. The coats of the stomach were somewhat thinned, but not to any marked extent. The kidneys were normal in size, capsules easily removed, but the cortical portions thinned, otherwise healthy. On pulling over the small intestines and mesentery, there was found lying on the left side of the bodies of the two lower lumbar vertebræ a solid mass, firmly adherent. I removed a good portion of this tumour with the knife for subsequent examination; it was about the size of two kidneys, and was outside, but enveloped by, the peritoneum. A number of enlarged glands were found on the under surface of the mesentery, about the size of buttons, and these passed down chain-like into the pelvis, along the rectum, being larger on the left side. The glands in the femoral canal were also enlarged. The bladder was healthy.

Wigan.

A Mirror OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

WEST LONDON HOSPITAL.

A CASE OF HÆMORRHAGIC DIATHESIS; HÆMORRHAGE
FROM THE MOUTH; DEATH.

(Under the care of Dr. THOROWGOOD.)

FOR the following notes we are indebted to Mr. Sidney H. Lindeman, house-physician.

William A—, aged eighteen, fairly well nourished, but presenting a strumous and anæmic appearance, and who for some time past had lived badly, was admitted on July 9th, 1880. He first attended as an out-patient on July 8th, and stated that on July 4th about 3 A.M. he found that he was bleeding from the gums. This lasted for about four hours and then stopped, but reappeared shortly before coming to the hospital. He had never, he said, noticed it before, but had been subject to profuse hæmoptysis, and at times had found "large blue lumps" on the chest, over the sternum, coming without any cause. These swellings lasted a few days, and then disappeared. He had never had any hæma-