This article was downloaded by: [130.132.123.28]

On: 27 December 2014, At: 04:53

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 7

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tnah13>

XIII.—Notes on the family Hetrodidæ, with a list of the described species

W.F. Kirby F.L.S. F.E.S.

Published online: 28 Sep 2009.

To cite this article: W.F. Kirby F.L.S. F.E.S. (1899) XIII.—Notes on the family Hetrodidæ, with a list of the described species , Annals and Magazine of Natural History: Series 7, 3:13, 97-101, DOI:

[10.1080/00222939908678083](https://doi.org/10.1080/00222939908678083)

To link to this article: <http://dx.doi.org/10.1080/00222939908678083>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content

should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

XIII.—*Notes on the Family Hetrodidæ, with a List of the described Species.* By W. F. KIRBY, F.L.S., F.E.S., &c.

THE family Hetrodidæ includes a number of rather large spiny apterous Orthoptera, a synopsis of which was published by Dr. Karsch in 1887 in the 'Berliner entomologische Zeitschrift,' vol. xxxi. Several additional species have been described since. I have just completed the rearrangement of the specimens in the Natural History Museum, and have made several notes on synonymy &c., which I thought it would be useful to publish at once, and I prefix a list of all the species hitherto described.

With the exception of one or two species which are said to occur in Asia Minor, the family is wholly African, and the species appear to be more numerous in Eastern and, perhaps, Southern Africa than on the West Coast. The Spanish and North-African genus *Pynogaster*, Graells, is now referred by Brunner von Wattenwyl to the family Ephippigeridæ instead of to the Hetrodidæ.

The synonymy of the true Hetrodidæ is given by Karsch in the paper just quoted, and I have therefore only given full references where additions or corrections were required.

Species marked † are wanting in the Museum collection; ‖, as usual, denotes a preoccupied name; * denotes type of genus.

List of described Species of Hetrodidæ.

Family Hetrodidæ.

Subfamily HETRODINÆ.

HETRODES, Fischer de Waldheim (1833).

- *1. *Hetrodes pupus*, Linn.—Africa, Cape Verde Islands, Asia Minor.

[Typ. fig., Roesel, Insectenbelust. ii. Heuschrecken, tab. vi. fig. 3.]

Hetrodes pupa et spinulosus, F. de W. Orth. Ross. pp. 368, 369, pl. xxxiv. figs. 1, 2 (1846).

2. *Hetrodes productus*, Walker, Cat. Derm. Salt. ii. p. 227. n. 4 (1869).—Congo.

3. *Hetrodes macrurus*, Walker, l. c. p. 228. n. 6 (1869).—?

4. *Hetrodes marginatus*, Walker, l. c. p. 226. n. 3 (1869).—Cape.

5. *Hetrodes abbreviatus*, Walker, *l. c.* p. 227. n. 5 (1869).—S. Africa.

†6. *Hetrodes Bachmanni*, Karsch.—Troe Troe, W. Cape Colony.

†7. *Hetrodes variolosus*, Fieber.—?

Hetrodes variolosus, Fieber, *Lotos*, iii. p. 259 (1853).

ACANTHOPLUS, Stål (1873).

§ Femora spined beneath.

*1. *Acanthoplus longipes*, Charpentier.—Benguela, Congo.

§§ Femora not spined; no spines on front margin of pronotum.

2. *Acanthoplus pallidus*, Walker.—E. Africa.

Hetrodes pallidus, Walk. Cat. Derm. Salt. ii. p. 231. n. 20 (1869).

3. *Acanthoplus discoidalis*, Walker.—S. Africa.

Hetrodes discoidalis, Walk. *l. c.* p. 230. n. 18 (1869).

†4. *Acanthoplus Speiseri*, Brancsik, *Trencs. Várm. Term. Egylet*, 1894–5, p. 258, pl. viii. fig. 9 *a–d* (1896).—Zambesi.

5. *Acanthoplus desertorum*, Kirby (*vide infra*).—Kalahari Desert.

6. *Acanthoplus serratus*, Kirby (*vide infra*).—King William's Town, S. Africa.

7. *Acanthoplus germanus*, Kirby (*vide infra*).—King William's Town.

†8. *Acanthoplus stratiotes*, Brancsik, *l. c.* p. 259, pl. viii. fig. 10 *a–d* (1896).—Zambesi.

§§§ Femora not spined; two central spines on front margin of pronotum.

†9. *Acanthoplus Jallæ*, Griffini, *Boll. Mus. Zool. Torino*, xii. no. 290, p. 3 (1897).—Zambesi.

CLOANTHELLA, Bolivar (1890).

Cloanthella, *Bol. Jorn. Sci. Lisboa*, (2) i. p. 226 (1890).

*†1. *Cloanthella clypeata*, *Bol. l. c.*—Braganza.

COSMODERUS, Lucas (1868).

- *1. *Cosmoderus erinaceus*, Fairmaire.—Gaboon, Cameroons, French Congo.
Ephippiger erinaceus, Fairmaire, Thomson, Archives Ent. ii. p. 260, pl. ix. fig. 1 (1858).
- 2. *Cosmoderus Kingsleyæ*, Kirby, Ann. & Mag. Nat. Hist. (6) xviii. p. 260, pl. xii. fig. 7 (1896).—Ogové.

APROPHANTIA, Kirby (1896).

[Kirby, Ann. & Mag. Nat. Hist. (6) xviii. p. 261, note (1896).]

- *1. *Aprophantia maculata*, Kirby, l. c. pl. xii. fig. 8 (1896).—Cameroons.

ENYALIOPSIS, Karsch (1887).

Types *E. ephippiatus* and *E. Petersii*.

- 1. *Enyaliopsis ephippiatus*, Gerstaecker.—Zanzibar.
Eugaster ephippiatus, Gerstaecker, Arch. f. Nat. xxxv. p. 213 (1869);
Von der Decken's Reisen, iii. (2) p. 27, Taf. ii. fig. 2 (1873).
- 2. *Enyaliopsis Petersii*, Schaum.—E. Africa.
- †3. *Enyaliopsis Bloyeti*, Lucas.—Kondoa, E. Africa.
- 4. *Enyaliopsis Durandi*, Lucas.—Zambesi.
- †5. *Enyaliopsis obuncus*, Bolivar.—Angola.
Enyalus obuncus, Bol. Jorn. Sci. Math. Lisboa, viii. p. 119 (1881).

GYMNOPROCTUS, Karsch (1887).

- †1. *Gymnoproctus abortivus*, Serville.—Senegal, Soudan.
Eugaster Maurelii, Lucas (sec. Karsch).

Subfamily EUGASTRINÆ.

ANEPISCEPTUS, Fieber (1853).

Pornotrips, Karsch (1887).

- †1. *Anepisceptus horridus*, Burmeister (type of both the above generic names).—Syria, Arabia, Egypt.
|| *Hetrodes spinulosus*, Charpentier.

2. *Anepisceptus Servillei*, Reiche and Fairmaire.—Abyssinia, Somali.

Hetrodes Servillei, R. & F., Ferret and Galinier, Voy. Abyss. iii. p. 426, pl. xxviii. fig. 1 (1847).

- †3. *Anepisceptus Revoilii*, Lucas.—Somali.

- †4. *Anepisceptus hippolyti*, Kirby (*vide infra*).—Abyssinia.

|| *Eugaster Servillei*, Lucas, Ann. Soc. Ent. France, (4) ix. p. 83, pl. iii. figs. 1-6 (1869).

5. *Anepisceptus Ruspolii*, Schulthess-Rechberg.—Somali.

Pornotrips Ruspolii, S.-R. Zool. Jahrb. Syst. viii. p. 81, pl. iv. fig. 4 (1894).

6. *Anepisceptus suakimensis*, Kirby.—Suakim.

Eugaster suakimensis, Kirby, Ann. & Mag. Nat. Hist. (6) xvii. p. 122 (1896).

- †7. *Anepisceptus Robecchii*, Schulthess-Rechberg.—Somali.

Pornotrips Robecchii, S.-R. Ann. Mus. Genov. (2) xix. p. 209, pl. ii. figs. 15, 15 a (1898).

EUGASTER, Serville (1839).

1. *Eugaster spinulosus*, Linné.—Morocco.

[Typ. fig. Edwards, Gleanings Nat. Hist. ii. p. 161, pl. cclxxxv. figs. 3-5 (1760).]

Eugaster spinulosus, Kirby, Ent. M. Mag. xxvii. pp. 210, 295 (1891).

2. *Eugaster frater*, Kirby, Ann. & Mag. Nat. Hist. (6) xvii. p. 123 (1896).—Mombasa.

3. *Eugaster Woodii*, Kirby, Ent. Monthly Mag. xxvii. p. 211 (1891).—Somali.

4. *Eugaster loricatus*, Gerstaecker, Arch. f. Nat. xxxv. p. 213 (1869); Von der Decken's Reisen, iii. (2) p. 26, pl. ii. fig. 1 (1873); Karsch, Berl. ent. Zeitsch. xxxi. p. 65 (1887), xxxii. p. 463 (1888).—Zanzibar.

5. *Eugaster Powysi*, Kirby, l. c. p. 294 (1891).—Morocco.

- *6. *Eugaster Guyoni*, Serville.—Algeria.

Eugaster Guyoni, Kirby, l. c. p. 295 (1891); Vosseler, Jahreshefte Ver. Württemb. xlix. p. xciii (1893); Finot, Ann. Soc. Ent. France, lxxv. p. 553 (1897).

|| *Eugaster spinulosus*, pt., Karsch (excl. syn.), l. c. xxxi. p. 65 (1887).

7. *Eugaster Lucasi*, Kirby, *l. c.* p. 295 (1891).—Tunis.

Hetrodes Guyoni, var., Lucas, Bull. Soc. Ent. France, (2) ix. p. 4 (1851).

Eugaster Guyoni, Lucas, Ann. Soc. Ent. France, (4) i. p. 217 (1861);
White, Proc. Ent. Soc. Lond. 1888, p. xxv.

Eugaster Guyoni, var. *Lucasi*, Krauss, Jahreshefte Ver. Württemb.
xlix. p. xevi (1892); Zool. Jahrb. Syst. ix. p. 552 (1897).

†8. *Eugaster inornata*, Krauss.—Oran.

Eugaster Guyoni, var. *inornata*, Krauss, *ll. cc.* (1892, 1897).

|| *Eugaster Guyonii*, Karsch, *l. c.* xxxii. p. 462 (1888).

ACANTHOPROCTUS, Karsch (1887).

†1. *Acanthoproctus* (?) *capreolus*, Pictet, Mém. Soc. Genève,
xxx. (6) p. 69, pl. iii. figs. 33, 33 a (1889).—S. Africa.

*2. *Acanthoproctus cervinus*, De Haan.—Cape.

Hetrodes fortis, Walker, Cat. Derm. Salt. ii. p. 229. n. 11 (1869).

3. *Acanthoproctus militaris*, White, Methuen, Life in Wilderness, p. 316, pl. i. fig. 4 (1846).—S. Africa.

4. *Acanthoproctus Howarthæ*, Kirby (*vide infra*).—E. Karoo,
Cape Colony; Brak Kloof, near Grahamstown.

HEMIHETRODES, Pictet (1889).

Hemihetrodes, Pictet, Mém. Soc. Genève, xxx. (6) p. 74 (1889).

1. *Hemihetrodes vittatus*, Walker.—S. Africa.

Hetrodes vittatus, Walker, Cat. Derm. Salt. ii. p. 229. n. 12 (1869).

Hemihetrodes Peringueyi, pt., Pictet, *l. c.* pl. iii. figs. 30, 30 a (1889).

2. *Hemihetrodes Peringueyi*, Pictet, *l. c.*—S. Africa.

APHRACTIA, Kirby (*vide infra*).

|| *Enyalus*, Stål, Öfv. Vet.-Akad. Förh. xxxiii. (3) p. 58 (1876).

Eugaster, div. a, Stål, Rec. Orth. ii. p. 22 (1874).

1. *Aphractia diademata*, Stål (1858).—S. Africa.

Hetrodes crassipes, Walker, Cat. Derm. Salt. ii. p. 231. n. 19 (1869).

Acanthoproctus ibex, Pictet, Mém. Soc. Genève, xxx. (6) p. 72, pl. iii.
figs. 31, 31 b (1889).

MADIGA, Kirby (1896).

Madiga, Kirby, Ann. & Mag. Nat. Hist. (6) xvii. p. 124 (1896).

|| *Prionocnemis*, Karsch.

*1. *Madiga verrucifera*, Karsch.—Mombasa.

Prionocnemis verruciferus, Karsch, Berl. ent. Zeitschr. xxxi. p. 67, Taf. ii. fig. 3 (1887); Ent. Nachr. xiii. p. 261 (1887).

†2. *Madiga aberrans*, Schulthess-Rechberg.—Somali.

Prionocnemis aberrans, S.-R. Ann. Mus. Genov. (2) xix. p. 210, pl. ii. figs. 16, 16 a (1898).

SPALACOMIMUS, Karsch (1887).

1. *Spalacomimus talpa*, Gerstaecker.—Zanzibar.

Eugaster talpa, Gerst. Arch. f. Nat. xxxv. p. 214 (1869); Von der Decken's Reisen, iii. (2) p. 28, Taf. ii. fig. 3 (1873).

BRADYOPISTHIUS, Karsch (1887).

1. *Bradyopisthius paradoxurus*, Karsch.—Somali.

[To be continued.]

XIV.—*New Species of Oriental Lepidoptera.*

By Colonel C. SWINHOE, M.A., F.L.S.

Fam. Nymphalidæ.

Subfam. EUPLEINÆ.

1. *Caduga ethologa*, nov.

♂ ♀. The Perak form of *C. tytia*, Gray, figured by Distant in Rhop. Malay. pl. xli. fig. 15, differs constantly from both Indian and Chinese forms in the long series before me, not in the division or otherwise of the large pale spot in the anterior wings between the second and third medial nervules as stated by Distant (p. 409), this character being inconstant in the group, but in the difference in the form of the subapical hyaline streaks of the fore wings, which is very characteristic in the group—the lower streak being short and the upper ones reduced to mere spots, whereas in *tytia* the lower streak is always long; the fore wing of this form is also much shorter than it is in *tytia*, the apex less produced, and the outer margin more abrupt.

Expanse of wings $3\frac{1}{2}$ inches.

Perak. Many examples.

This form is in the B. M. from Perak, unnamed.