- New York Medical Journal*, 1885, XLII. 145. 1886, XLIII. 384. 1886, XLIV. 373. 1886, XLIV. 322. 1887, XLV. 238. 1887, XLV. 273. 1887, XLV. 624. 1887, XLV. 640. 1887, XLVI., II. 1887, XLVI. No. 5.
- Journal American Medical Association*, 1886, VI. 147. 1886, VI. 186. 1886, VI. 426. 1886, VII. 35. 1887, VIII. 199. 1887, VIII. 342. 1887, VIII. 337. 1887, VIII. 359. 1887, VIII. 291. 1887, VIII. 701. 1887, IX. 135.
- Letter to E. Fletcher Ingalls. Reprint from the *New York Medical Journal*, July 2nd and 9th, 1887.
- Statistical Records of Intubation, *Medical Record*, July 23, 1887; *New York Medical Journal*, August 8, 1885.
- Transactions of the American Laryngological Association*, 1887.
- Chicago Medical Journal and Examiner*, 1885, L. 475. 1885, LI. 511. 1886, LIII. 132. 1886, LII. 214. 1885, LI. 401.
- Medical and Surgical Reporter*, 1886, LV. 586.
- New Yorker Med. Presse*, 1887, III. 259. 1887, June.
- Medical Register*, 1887, I. 174. 1887, I. 146.
- Medical News*, 1887, L. 341. 1887, L. 630.
- Maryland Medical Journal*, 1886, XVI. 67. 1886, XVI. 168.
- Medical Review*, Pittsburg, 1887, I. 114.
- Weekly Medical Review*, St. Louis, 1887, XV. 57.
- Pacific Medical and Surgical Journal*, 1887, XXX. 129.
- Chicago Medical Times*, 1887-88, XIX. 53.
- Cincinnati Lancet and Clinic*, 1887, N.S., XVII. 97. 1887, N.S., XVIII. 321.
- Hahnemannian Monthly*, June, 1886.
- American Practitioner and News*, 1886, N.S., II. 321.
- St. Louis Medical and Surgical Journal*, 1887, LII. 115.
- American Lancet*, Detroit, 1886, N.S., X. 401.
- Canadian Practitioner*, Toronto, 1887, XII. 1.
- Archives of Pediatrics*, 1885, II. 657. 1886, III. 215. 1887, IV. 154.
- Medical Age*, Detroit, 1886, IV. 313.
- Mississippi Valley Medical Monthly*, 1886, VI. 345.
- Albany Medical and Surgical Journal*, 1887, II. 331.
- Boston Medical and Surgical Journal*, 1887, CLVI. 518.
- American Journal of Obstetrics*, 1816, June 17.

REVIEWS.

CHIARI (Vienna).—**Observations on Diseases of the Throat and Nose.** *Topfritz and Deutsche, Leipzig and Vienna*, 84 pages.

THE author relates observations on more than 2,000 patients seen in the clinic of Prof. Nothnagel, many of which are very interesting. One case of laryngitis hæmorrhagica (Strübing) was combined with a fibroma; acute subglottic laryngitis (chorditis hypertrophica inferior) was seen in two cases. Chronic laryngitis was treated with nitrate of silver and iodized glycerine. Once a case of circumscribed abscess of the vocal cord occurred: cured by incision and removal of the pus. Phlegmonous inflammations of the larynx occurred seven times. In two cases of tuberculosis, tracheotomy was necessitated by reason of stenosis. Lactic acid is preferred for the

treatment of tubercular ulcers. The use of cocaine in laryngeal operations is discussed. Intoxications are rare, and are due to idiosyncrasy. The author favours Voltolini's sponge method for papillomata, which, however, recur after every method of operation. Cysts of the vocal cord were seen in two cases, and one was operated upon. Five cases of carcinoma were seen, and two were cured by extirpation of the larynx. Stenoses are treated with Schroetter's tin and hard rubber bougies.

Many cases of laryngeal paralysis were seen, two of aphonia spastica, and one of chorea laryngis. Compression of the trachea was met with in ten cases of struma. Many cases of reflex neuroses of the nose were seen, and local applications, particularly of the galvano-cautery, were most beneficial.

Ozena was treated with $\frac{1}{2}$ per cent. salt irrigations, and eczema with diachylon ointment. A frequent cause of epistaxis was noted in enlarged vessels of the septum and excoriations of this locality, and the galvano-cautery was the best treatment. Polypi and other growths in the nares were operated on by galvano-cautery, cold wire, or forceps.

A rhinolith was found in one case. Adenoid vegetations were operated on with cutting forceps.

Michael.

HERING, T.—The Curability of Laryngeal Phthisis by Surgical Methods. A work based on Microscopic, Anatomico-Pathological, and Clinical Observations. Warsaw, 1887.

THIS is a very careful work, illustrated with many plates, and deals with the etiology and pathological anatomy, and pathological proofs of the curability of phthisical ulcers of the larynx; a description of the surgical methods invented and employed by the author, viz., curetting, frictions, and injections of lactic acid, and injections of iodoform emulsion; a description, with drawings, of the instruments invented by the author; and lastly, the history of thirty-five cases in which these methods were employed. From the author's observations, the conclusions are arrived at, that laryngeal phthisis ought to be treated locally by surgical methods, just as tuberculosis of other organs accessible to surgical manipulations; that by following these methods cicatrization of ulcers can be procured, and a more or less lasting cure, with disappearance of the sufferings of the patient, and gain in strength. Recurrences, however, occur from time to time, and the cure of these ulcers does not influence the progress of the disease in the lungs. Speaking in general terms, this work is careful, giving conclusions which are not too optimistic, even for an author so enthusiastic of his ideas and methods. Hering's treatment consists in the following:—

1. At the commencement of the disease, parenchymatous injections of iodoform emulsion, or of 10 per cent. to 20 per cent. lactic acid, should be made into the larynx.

2. During the stage of softening and inflammation of the infiltrated parts, deep incisions into the affected parts (epiglottis, posterior wall of the larynx, aryepiglottic ligaments), or extirpation of the affected parts by means of forceps, guillotines, and curettes, should be performed. Anæsthesia should first be obtained by 10 per cent. solution of cocaine,