so as to avoid uncertainty and confusion both in clinical therapeutics and in current medical literature. To physicians who desire an extra pharmacopeia containing valuable formulæ not contained in the text, I would recommend the "Formulary" of the American Pharmaceutical Association; which contains efficient substitutes for many of the most prominent proprietary remedies, and thus offers a means of avoiding those of secret composition, the use of which the Code of Ethics condemns as unscientific and unworthy of educated physicians. It seems to me, if persistent effort were made by both physicians and pharmacists to instruct the community as to the dangers to the public health resulting from the constant use of patent medicines, that the business might be made less profitable than it now is, and the consumption of such articles very much curtailed.

In these brief introductory remarks, I have purposely refrained from discussing exhaustively any single topic, and have been content to indicate a few directions in which advancement may be made in the departments of medicine represented in this Section. Improvements might also be introduced in the work of this Section so as to make it more popular and attractive, and I believe that some recommendations will be made to this end at the present session, by the Business Committee.

FIBROID TUMORS OF THE UTERUS.

Address of the President, read before the Mississippi Valley Medical Society, Oct. 4-6, 1893.

BY R. STANSBURY SUTTON, M.D.

PITTSBURG, PA.

Definition.—The familiar terms, fibroid tumor, fibro-cystic tumor, and fibroid polypi of the uterus, every medical man recognizes as belonging to the same class of neoplasms. We all understand by the fibro-cystic tumor that a solid fibroid tumor has undergone a cystic degeneration. Notwithstanding our familiarity with these terms, they are objectionable, because they do not convey to the mind a comprehensive idea of the pathology of the neoplastic growths to which they are applied. The term fibroid (fibrosus and eidos), and the term fibrous mean fiber-like, and are only applicable to a tumor composed entirely of fibrous tissue.

The fibrous or fibroid tumor found in various localities is always situated in connective tissue; it contains only occasionally a few yellow elastic fibers, is often quite vascular, and must be dissected out.

But the so-called uterine fibroid contains a variable quantity of muscular tissue as well as of fibrous tissue, both of which are derived from the parenchyma of the uterus in which the tumor grows. By reason of the presence of muscular tissue, Virchow proposed that these tumors be called myomata. This term designates only their muscular element, and is as objectionable as the term, fibroid. The term, myofibromata or fibro-myomata, fully designates the solid tumor, and the prefix cystic, that variety in which cystic degeneration has occurred.

Etiology.—The etiology of these tumors is still a mystery. They are more frequent in the black than in the white race. In both races they are more frequent in the body than in the neck of the uterus, and more frequently situated in the posterior than in the anterior wall. Prior to puberty they present no positive symptoms of their existence.

From a careful analysis of cases under my care, I suspect the neoplasm does sometimes exist prior to puberty. After the menopause is established these tumors occasionally disappear, unless they have undergone cystic degeneration.

If, prior to the establishment of the menopause, the fibro-myomata has undergone cystic degeneration, it will continue to grow. The removal of the ovaries and Fallopian tubes in such cases has proved insufficient to stop the growth of the cystic or very soft fibro-myoma. Months after removal of both ovaries and tubes I have seen it necessary to make a supra-vaginal hysterectomy. Concerning the causes which produce these tumors nothing is known. By reason of their existence a woman may be sterile, but it is not yet proved that sterility favors their origin or growth.

Pathology.—The tumor always has its origin in the substance of the uterine wall or subperitoneal connective tissue. It begins as a hard nodule, involving in its further development both the muscular and fibrous tissues. Later it becomes invested with a thick layer of fibrous tissue, which, with the superimposed uterine tissue, constitutes its envelope or capsule. Within this capsule the blood vessels ramify and form a network, sending prolongations to feed the inclosed growth, which now lies like a foreign body in or attached to the wall of the uterus. Rarely, large blood vessels penetrate the substance of the tumor. Such tumors have, however, been observed by Virchow, Leopold and Schröder, in which large blood-sinuses existed, the walls of the sinuses being composed of the muscular fiber of the uterus involved in the growth of the tumor.

Klebs has proved the existence of lymphatics, and Lorey of nerves in this class of growths. The presence of lymphatics often determines the variety of future cystic degeneration. As the tumor enlarges it separates the tissues composing the uterine wall, or compresses it as it pushes the peritoneum outward. The consequent irritation invites a greater blood supply, and hyperplasia of the walls begins. As the tumor excites more and more irritation, and the muscular tissues of the uterus grow stronger, the neoplasm, if developing within the walls, may begin to migrate either toward the peritoneal covering of the uterus or toward the endometrium. Should, however, its location favor, it may separate wider and wider the uterine tissues, keeping its relative position from either surface. Thus the tumor is finally fixed in the uterine walls, and is designated as interstitial or intramural; or if it is developed or forced toward the peritoneum, carrying the latter with it as an investment, it is then designated as submucous. The tumor may consist of a single nodule or center of formation, or it may consist of a number of nodules or centers of formation interlaced closely and invested by a common capsule. This fact suggested to Mr. Lawson Tait a new nomenclature for these growths—namely, uninodular and multinodular fibro-myomata. As the tumor grows, either the fibrous tissue or the muscular tissue will preponderate.

When the fibrous tissue is bountiful, the tumor will be hard, even as hard as cartilage; when the muscular tissue is most bountiful the tumor will be softer, even sufficiently so as to merit the name, fleshy tumor or myoma. A tumor largely composed of fibrous tissue will usually be of moderate size and of

very slow growth, but the reverse is apt to occur in those tumors in which the muscular tissue preponderates; and these tumors are especially liable to cystic degeneration. On the other hand, the latter class, if cystic degeneration does not occur, are liable to rapid disappearance after the establishment of the menopause. Not infrequently several separate and distinct fibro-myomata are found in the same uterus. Dr. Thomas exhibited to the New York Pathological Society a uterus containing thirty-five tumors of various sizes, and Schröder of Berlin, cites a case by Schultz, in which at least fifty tumors existed in the uterus. A remarkable feature of Schultz's case was that the woman was 83 years of age. The uterus was obtained postmortem. In cases of multiple myomata, the growths may be so distributed in the uterus as to present the subperitoneal, the intramural, submucous, and polypoid varieties.

The changes occurring in the uterus itself are not constant. The muscular tissues hypertrophy to a greater or less extent in all cases, and while hypertrophy is going on in one direction, atrophy of the muscular tissue on the side toward which the tumor is advancing may be in progress. After the tumor has become subperitoneal, the uterus may atrophy to a very marked degree—a condition not infrequent in old subjects.

The cavity of the uterus is usually increased in depth, while in the submucous and intramural variety it may also be obstructed. The lining membrane of the uterus, by reason of the constant hyperemia, is more vascular, bleeds more readily upon the introduction of the sound, than in cases of normal condition, while in cases of submucous fibro-myomata or polypoid fibro-myomata, menstruation is almost always profuse, and between the regular periods blood is frequently lost or even constantly. In addition to this, a serous discharge, intermingled with the vaginal and uterine secretion and having an unpleasant odor, is not always wanting. The uterus in no instance occupies its normal position, but is either pulled or pushed into an abnormal one, dependent on the size and position of the tumor. The cut section of fibro-myoma will vary in appearance according to the preponderance of its constituents. If the fibrous tissue be very abundant, the section will cut like cartilage, and have a gray color and satin-like glossy appearance. If the muscular tissue exists to a marked degree, the consistency of the section will be softer and the color will vary from pink to red.

The lymphatic spaces of Klebs are sometimes seen between the bundles of muscular tissue. When but little muscular tissue is present the section will be smooth, but if the reverse is the case the surface is uneven as the contracting fibrous bands force the softer muscular tissue above the surface. At the margin of the section and surrounding it is found a layer of condensed, fibrous, muscular tissue forming the capsule. This is sometimes closely, at other times loosely, attached to the tumor. Frequently it is easy with the thumbnail to detach it entirely, a process we shall yet learn to be that of enucleation. On microscopic examination of the harder variety are seen wavy bundles of fibrous tissue with a small amount of unstriped muscular tissue. The softer variety presents, in addition to the fibrous tissue, a greater amount of unstriped muscular fibers in twisted bundles, the muscular fiber being identical

with the muscular fiber of the uterus. If the specimen be stained in carmine solution and washed in a solution of acetic acid, the rod-shaped nuclei of the spindle-shaped cells will be brought into view.

The important practical point in this is to establish a positive differential diagnosis between the fibro-myoma and sarcoma. The latter, when removed from the uterus, as elsewhere, is almost sure to return, and constitutes what has been erroneously called the recurrent fibroid tumor.

The nuclei of the spindle-shaped sarcoma cells are round or oval; of the fibro-myomata, they are rod-shaped. The limit of growth for the myomata is difficult to determine. They have, after removal, been found to weigh fifty or sixty pounds. Until expelled from the walls of the uterus they maintain a globular form. After extrusion into the cavity of the peritoneum they are free from the pressure of the uterine muscular tissue, and generally lose the globular form. After extrusion into the cavity of the uterus they usually become pear-shaped—a fact probably due to the lateral pressure exerted upon them in the expulsive efforts of the uterus. Their density varies with the amount of fibrous tissue in their composition, and is a determining element in the shape of their future growth, when they become subperitoneal. The globular form is not, however, infrequently retained in very solid tumors after they have become polypoid tumors, either upon the exterior or interior surface of the uterine walls. The irregularity of surface occasionally seen is due often to the multinodular composition of the neoplasm, some nodules growing faster than others by reason of obtaining a better blood supply. Another element in determining the shape of the subperitoneal tumor is the pressure against the walls of the pelvis; regularity of contour is more apt to exist when the tumor has become too large to enter the small pelvis. Occasionally, in addition to the mass in the pelvis, a prolongation of the tumor upward, cone-like, is felt through the abdominal wall. I have seen the tumor pear-shaped, the big end up, and the other end filling the pelvis completely.

Frequently the tumor, when subperitoneal, may be felt extending above Poupart's ligament, dragging the uterus with it, side by side, it being difficult or even impossible to decide, when the os is almost or entirely out of reach, which is tumor and which is uterus. The sound may determine. If the cervix can be seized with a volsellum forceps and the uterus be thus moved, while the free hand over the abdomen takes cognizance of the movements, it may also be determined.

The changes which occur in the fibro-myomata are as follows—varieties of softening:

Edema.—At the time of the menstrual period, they occasionally become swollen or edematous, this condition disappearing again a few days later. But this condition may continue to such a degree that a spurious fluctuation may exist. Such tumors have been tapped for ovarian cysts, and a few drops of yellow, slimy mucus escaping told of the error of diagnosis, or the tapping has been dry. Such tumors may give apparently real and distinct fluctuation after being extirpated and laid on the table. The tumors when cut through may be completely infiltrated; in the meshes of fibrous tissue hundreds of very small cysts exist, and the muscular tissues of the growth may have wellnigh disappeared. Such

tumors have been found by many operators. The edema is sometimes intermittent, returning after entire subsidence.

Fatty Degeneration.—After the menopause, and occasionally after pregnancy, complicated with intramural fibro-myoma, fatty degeneration occurs in the muscular element of the growth; this is absorbed and the fibrous tissue alone left. In very old subjects the remains of prior fibro-myomata are simply nodules of the concentrated fibrous tissue.

Myxomatous Degeneration.—Occasionally, in the tissues comprising the growth, there exists mucoid tissue which secretes mucus, and forms often cavities of considerable size; this condition of the tumor is known as myxomatous degeneration. (Virchow.)

Suppuration.—Should the blood supply be entirely cut off from the growth, as in twisting of the pedicle in the subperitoneal variety or of the polypoid submucous variety, or if the capsule be widely detached in an effort at enucleation per vaginam, the tumor may die and soften. This change has been designated erroneously as suppuration. True inflammatory softening and suppuration of a fibro-myoma is rarely seen.

Gangrene.—The submucous variety is especially liable to necrosis and spontaneous expulsion. An inflammation of the capsule resulting in interference with the nutrition of the growth, or inducing an ulceration on the surface of the capsule, is the usual cause. The resistance of the capsule being impaired by the opening, the tumor is expelled by uterine contractions entire or piecemeal. This fact led to the treatment of this variety by means intended to open the capsule through the cavity of the uterus. Atlee and Brown gouged holes in the capsule or divided it with the knife. Greenhalgh attacked the capsule with the cautery-iron. Byford induced rupture of the capsule with ergot. Cures have been obtained by all of these methods, none of which were free from danger.

Cases of inflammation of the capsule, adhesion to and perforation of the abdominal wall, followed by the escape of the gangrenous neoplasm, have been reported by Soir, Dumesnil, Gutierrez, Hofmokl and Schmidt.

Varieties of Induration.—After fatty degeneration of the muscular elements, their absorption follows, but the fibrous tissue is left behind and contracts to form a very hard but small tumor. When the tumor contains very little muscular tissue it is nearly as hard as cartilage.

Calcification.—Chalky or phosphatic degeneration of these tumors has long been known. Hippocrates relates such a case, a Thessalian woman aged 60 years; and Salius the case of a nun who had such a tumor. Schröder refers to such cases by Louis, Velpeau, Jaffe and Saxinger.

Years ago I removed a calcified subperitoneal fibro-myoma, which was larger than a goose egg and as hard as a stone. It was reached through the posterior vaginal wall. The cretaceous matter appears first, says Schröder, in streaks through the interior of the tumor. It may finally become so dense as to produce a stone of sufficient solidity to require to be cut through with a saw. The small tumors, especially the subperitoneal and intramural, are most liable to this change. When the blood vessels are cut off, the nutrition ceases. The tumor is now a foreign body, and is apt to act as such. It

may cause an inflammation of the adjoining tissues, and when these are softened it may escape into the peritoneal cavity, causing fatal peritonitis, or, escaping into the uterine cavity, be expelled, leaving the patient to recover. True suppuration of a fibro-myoma must be rare, yet authors authenticate it after traumatic injury to the growths, and also in association with cretaceous degeneration beyond the changes referred to; these tumors, by various processes of degeneration, become cystic, and occasionally are apparently attacked with that most malignant of diseases, sarcoma.

Cystic Degeneration.—A solid fibro-myoma may become cystic through fatty degeneration, mucous degeneration, suppuration, serous infiltration, the formation of lymph-cavities filling with a clear fluid coagulating on exposure to the air, or from breaking down of clots in large blood cavities already existing in the tumor. In the multinodular tumors one or more of the nodules may break down, while others maintain their original solid condition, and a mixed tumor results. As already observed, these growths take largely to themselves the connective fibrous tissue of the uterus. Connective tissue is that in which alone we find sarcoma developing in the body. The fibro-myoma and sarcoma are first cousins as to origin, and when the sarcoma begins, with its spindle cell and oval or round nucleus, to invade the myoma, it finds the natural tissue of its selection to work upon. Myomata thus invaded grow softer, and may even undergo cystic degeneration, and give rise to a tumor properly designated sarcomatous cystic fibro-myoma.

The termination of the cystic fibro-myoma is widely different from that of the simple fibro-myoma; while the latter occasionally destroys the woman, the former always tends to terminate fatally. Fibro-cystic tumors grow more slowly than ovarian cystomata, but by mechanical interference with respiration, circulation and nutrition, and by producing nervous exhaustion and organic diseases of the heart, and also occasionally by causing a great loss of blood from the uterus, they tend with equal certainty to the destruction, sooner or later, of the life of the patient. The location of simple fibro-myomata has everything to do with their importance; it is therefore excusable to make the arbitrary division of subperitoneal, interstitial, or intramural, and submucous; also to consider each variety separately.

Description. Subperitoneal Fibro-myomata.—Virchow termed this variety the peritoneal polyp, and there is no difference of structure between it and the fibro-myomatous polyp found in the uterine cavity. When the tumor is forced out of the uterine wall into the cavity of the peritoneum, it carries with it the peritoneum investing the uterus, except in those cases where it projects from the side of the uterus and is pressed out between the layers of the broad ligament, which subsequently forms a serous covering for it. The tumor may remain closely adherent to the wall of the uterus, or gradually leaving it, develop a pedicle which may, after a time, remain thick or become so attenuated as to consist of nothing but two layers of peritoneum, the intervening cellular tissue and blood vessels, some lymphatics, and nerves. The tumor continues to grow in the cavity of the pelvis, toward which it gravitates, retroverting the uterus early if attached to the pos-

terior wall, until finally it reaches such a size that it can not enter the superior strait of the pelvis. When the tumor projects from the top or anterior wall of the uterus, the latter is retroverted as soon as the growth has attained any considerable size. The growing of these tumors thus in the pelvis gives rise to distressing vesical and rectal irritation, and often to retention of urine, to the narrowing of the caliber of the rectum, to constipation, and to hemorrhoids. Besides this, there is often pain in the course of the sciatic or crural nerves of either side. The irritation and obstruction to the circulation, both from the mechanical presence of the growth and the malposition of the uterus, produces profuse bleeding at and between the menstrual epochs. That such is the pathology is proved by putting the patient in the knee-chest position, pushing the incarcerated tumor and uterus up, and draining the vaginal tissues with pledgets of cotton saturated with glycerid and by the free use of large quantities of hot water thrown against the vault of the vagina after the replacement is effected. Thus long-continued hemorrhages are abated.

Once the tumor has become subperitoneal it may contract adhesions to the viscera against which it rests, and thus it often happens in abdominal section that we find these tumors nourished by large blood vessels entering them through adhesions which, when the tumor is large and projecting into the cavity of the abdomen, frequently contain veins of enormous size. Twisting of the pedicle, when long, in these tumors may occur as in the ovarian tumor. If a new blood supply has been established, the tumor will not die, even though it may be eventually separated entirely from the uterus. When this variety of tumor drags the uterus upward, or when the base of the pedicle is broad and the tumor falls backward, bending the uterus, the cavity is increased in depth; as the pedicle becomes elongated the uterine cavity shortens whether the tumor decreases in size or not. The uterus may be flattened out completely and adherent to the side of the tumor. After it has contracted adhesions to the intestine, any rotary motion of the tumor is liable to produce intestinal obstruction and to demand immediate operative interference to save the life of the patient. Frequently I have removed fibroid tumors from which it was necessary to detach several loops of the small intestine. This variety of tumor is frequently accompanied by others of the same variety, either also expelled from, or still existing in the walls of the uterus. The greatest limit of growth as a rule for this variety is the size of an adult head, but they have been met with having a weight of fifty to sixty pounds.

Submucous Fibro-myomata.—As already stated, at least two varieties originate in the walls of the uterus as round tumors; when forced toward the lining membrane of the uterus they become submucous. As they project into the uterine cavity they carry the lining membrane before them. The pedicle of this, the polypoid variety of the tumor, may be thick, containing muscular tissue, the lining membrane of the uterus, and very small blood vessels. As long as the tumor lies beneath the unupheaved uterine lining it is round, but after it enters the cavity of the uterus and is macerated in constant discharge, and pressed upon by the contracting uterine walls, it becomes pear-shaped, or if nipped near the center by the

fiber of Bandle's ring it may resemble a dumb-bell or hour-glass. This variety of myomata does not undergo cretaceous degeneration and is usually expelled too soon to undergo cystic degeneration, which occurs rarely in this variety. Nevertheless, it is probable that lives are as often lost from hemorrhages occasioned by this variety as by either of the other varieties of the myomata. This variety drags down the uterus when of moderate size, and after expulsion from the uterus, may by its weight as it descends drag the fundus uteri down and invert the uterus. Women often hide these growths, even after they appear at the vulva.

The size of the growth is determined mainly by the length of time it remains in the uterus or vagina. When in the vagina the tumor may be mistaken for an inverted uterus.

The Intramural or Interstitial Fibro-myoma.—This variety simply remains and grows within the uterine walls. Surrounded on all sides by uterine tissue, it receives a greater blood supply than either of the other varieties. It consequently grows with greater rapidity, and often to an enormous size, enlarging a patient like a full term pregnancy, and weighing ten pounds and upward. The great weight stretches the anterior wall of the belly, and the tumor overhangs, while it rests upon the pelvic brim. The abdominal walls grow very thin from pressure, the recti muscles atrophy and separate, and the tumor lies immediately under the superficial tissues of the belly wall. These tumors, bulging out the uterus laterally, spread apart the layers of the broad ligament, and the hypertrophied tubes are spread out high up on the tumor; the uterus and the tumor develop in such a way as to entirely alter the relative position of the uterine appendages. In these cases the cavity of the uterus is deepened or shortened and often rendered crooked, while at the same time it is very difficult to find the os uteri in the vagina. These externally large tumors are usually single, and the walls of the uterus may be found either greatly hypertrophied, or much atrophied. These tumors occasionally undergo cystic degeneration, and grow to an enormous size.

Fibro-myoma of the Cervix Uteri.—These tumors follow the same law as to location and development that they do in the body of the uterus. They may be subperitoneal when developed in the supra-vaginal portion; when developed too low down, they very rarely appear upon the outer surface of the cervix, namely, the outer surface of the vaginal portion. The growth is much more rare in the cervix than in the body. In the twenty-eight years I have been in practice I have met with but five or six cases where the growth was in the cervix. In one the growth was submucous, and in another, interstitial and as large as a lemon; in another, the tumor was as large as a fetal head. Where they become submucous, they are gradually forced into the vagina, retaining their uterine connection through a pedicle of varying density. When interstitial and large they pack the pelvis to a great extent. The lip of the cervix invaded is spread out over the growth, while the opposite lip is thinned and stretched as a band along the circumference of its enlarged neighbor. The uterus is dragged down until the growth has become large, when its fundus is deviated, at first in a direction corresponding to the lip of the cervix invaded. When the tumor is expelled from the cervix it will be oval

or round-ended, and when pouting between the labia may readily be mistaken for the fundus of the inverted uterus. On the other hand, the uterus has been mistaken for this variety of fibro-myoma, and been cut off with the ecraseur. Large growths in the cervix are more liable to produce vesical and rectal symptoms than those growing either from the posterior wall or fundus of the uterus.

Symptoms.—The symptoms which accompany the presence of these tumors in their various localities and stages of growth are widely different, as we shall learn in considering the diagnosis. However, a class of indications are patent as found in a vast majority of the cases, and may be tabulated as follows:

- A. Vague pain in the pelvic region;
- B. Pain referred to the front or back of the leg;
- C. Irritability of the bladder or rectum;
- D. Uterine tenesmus;
- E. Menorrhagia or metorrhagia;
- F. Dysmenorrhea;
- G. Profuse leucorrhœa;
- H. Serous discharge from the uterus;

The disorders of menstruation, the tenesmus, the leucorrhœa and the serous discharge are typical, within the menstruating age, of the existence of the submucous fibro-myoma, and in a less marked degree in the interstitial variety. They may be entirely absent in the subperitoneal variety. It may be inferred as a rule in diagnosis that tumors producing these symptoms to a marked degree are situated immediately beneath or close to the lining membrane of the uterus. Exceptions to the rule do occur, and hemorrhages may accompany the existence of a subperitoneal fibro-myoma incarcerated in the pelvis. Progressive anemia from loss of blood, feeling of weight, depression of spirits, the development of unusual nervous conditions, irritability, loss of cheerfulness and gloomy apprehensions are frequent. When the tumor rises out of the pelvis it may be easily felt through the abdominal wall. If cystic degeneration has occurred, spurious or true fluctuation may be present; but absence of the symptoms will not be proof that cystic degeneration has not occurred.

Diagnosis.—Scarcely is it possible to name anything more difficult than the diagnosis of the very small fibro-myomata, when many of the leading symptoms are absent. This is sure to be the case with subperitoneal growths, and the surgeon is left entirely to exploratory skill. If the tumor has retroverted or anteverted the uterus, and become at all prominent upon either surface, bimanual palpation with a finger in the vagina or rectum will usually determine its existence. If the patient be fat, it will be necessary with the volsellum forceps to pull down the uterus, so that the finger in the rectum may be swept over the posterior surface of the uterus, or the sound may be well curved and made to drag the uterus well forward against the finger in the vagina. The tumor, if at all defined, will present an outline within which the tissue is harder than that of the normal uterus; if well defined, it will be easily recognized. Even at so early a date the uterine cavity is often found deeper than normal, and the uterus congested and abnormally heavy. When the tumor springs from the supra-vaginal portion of the cervix, it is readily felt through the vagina. But if it be from the posterior wall of the fundus, and of considerable size and firmly adherent in the cul-de-sac, a sound in the uterus may or may not enable the

examiner to determine that the growth is or is not attached to the uterus, or, if it is so attached, that it is a myoma. The direction taken by the sound and the consistency of the growth are the questions to be considered. The round form and solid feel of the myoma will aid in differentiating it from a hematocèle or ovarian cyst, but it is not possible to differentiate it from a solid tumor of the ovary which is adherent to the uterus. Twice I have seen the abdomen opened, once by Billroth and once by Mr. Lawson Tait, for supposed myoma of the uterus, to find the tumor a solid one of the ovary. The differentiation from a cyst is less difficult; its elasticity, its rare close connection with the uterus, and its softer feel, are suggestive points. But if doubt exist, the aspirator-needle may be used.

When the tumor is interstitial and very small it will be very difficult, if not impossible, to determine its presence. The existence of some of the symptoms, alphabetically tabulated, may give good presumptive evidence; in addition, the uterus may be deeper than normal, or it may be possible for one of unusual skill to determine, by the aid of the sound, that one wall of the uterus is thicker than the opposite wall; or that a portion of the wall is more solid or thicker. If the tumor is small and in the anterior wall of the uterus, the canal will be so displaced backward that the direction taken by the sound will suggest that the body felt forward is not alone the fundus of the anteflexed uterus.

When the tumor has attained a considerable size and other symptoms are present, and the possibility of enlargement from pregnancy, chronic metritis and subinvolution are differentiated, the case is not so difficult to determine. But given a small interstitial fibro-myoma associated with chronic metritis or pregnancy, and the diagnosis may be impossible. Time alone will solve it. Should, however, in early pregnancy the fetus be dead and the woman present an enlarged uterus, with irregular periods of bleeding, a constant leucorrhœa, or flow of disagreeable odor, the case may be mistaken for a fibro-myoma, interstitial or submucous, and nothing except dilatation of the cervix and exploration with the finger will make a diagnosis possible. In chronic metritis the uterus is more or less tender; its walls are flat and soft, the os open, and frequently nausea exists. In cases of fibro-myoma, the uterus is rarely sensitive, and especially when the growth is interstitial or subperitoneal, the body as well as the cervix is hard, and the os normal. If chronic cellulitis with extensive deposits be encountered, the uterus is fixed, while the reverse is usually the fact when the symptoms depend on a neoplasm.

In early pregnancy the cervix and lower segment of the uterus are softer and the os occasionally more patent than normal; while in cases of fibro-myoma the cervix and os usually remain unchanged and the tumor is hard. In pregnancy the color of the vagina is bluish. Further, in bimanual palpation the pregnant uterus will be found soft or elastic, symmetrical, rhythmical, and nearly in the central line; in cases of even small fibro-myoma it will be hard, inelastic and usually misplaced. In small submucous neoplasms the bleeding is usually so prominent a symptom that the introduction of a sound or dilatation of the cervix and introduction of a finger will suggest themselves as the speediest way to decide the diagnosis. In a submucous growth, with the cervix

dilated sufficiently to admit the finger, the diagnosis is not difficult. When the tumor grows from the intravaginal portion of the cervix, if interstitial its early symptom is simply an enlarged lip; but later its solidity, freedom from tenderness, its circumscribed hardness, and the absence of the evidence of malignant disease will warrant an incision into the lip and an enucleation of the growth at the same time that the diagnosis is made. When the small interstitial fibroid is low down, its development toward the os externum enlarges the lip beneath it and protrudes it forward into the vagina. The opposite lip is spread out and the os loses its form, becomes a slit, and may be very difficult to find. When the neoplasm is entirely in the cervix the same difficulty will occur. The liability of mistaking such cases for inversion of the uterus is to be guarded against. A retroflexed or anteverted uterus has frequently been taken for a fibro-myoma. The groove presenting to the examining finger between the neck and the fundus misleads; but the sound and bimanual examination will determine whether the round body felt is the fundus or a fibro-myoma. If the neoplasm arise from the posterior wall the uterus is retroverted, and the bimanual method of examination will enable us to trace its close connection with the uterus unless the pedicle be unusually long. If, however, the uterus and tumor fill the pelvis, the patient should be placed in the knee-chest position, or, better, in Sims' position, with the side of the table elevated. The uterus and tumor may now be pushed upward and forward out of the pelvis. A sound may then be introduced into the uterus and held with the thumb and forefinger of the right hand; the left hand passing over the patient's hip can grasp both uterus and tumor, and their connection may be determined by the movements of the sound. If the tumor can be grasped separately and moved without affecting the position of the uterus, it can not be uterine; but if its movements, as determined by the sound, do affect the position of the uterus, it is attached to the uterus and probably of uterine origin.

In thin subjects, especially, the sound may not be required; but bimanual examination with one or two fingers behind the cervix will enable the examiner to determine a close connection between the tumor and the uterus, or decide whether or not the tumor and uterus move together. But as it is the early development of fibro-myomata that will always give the case most difficult of diagnosis, a more lengthy examination of this subject will be required. A case or cases presenting, in which no change in the os or cervix has occurred, with no displacement, either retro- or anteversion or prolapsus, with or without much increase of the depth of the uterine cavity, with no marked elevation of either uterine wall to be felt bimanually, but with disordered and painful menstruation, or irregular discharges of mucus or blood, or the watery discharge of serum, will try the skill of the best diagnostician, and frequently compel him to summon time to his assistance. These symptoms, however, should always put us on the alert.

All growths of this nature are at first supposed to be interstitial, but if they are not, or become submucous, the leading clinical features of the case will be disordered menstruation, bloody discharges at irregular periods, and the presence of leucorrhea consisting of mucus more or less watery. If, however, the tumor be developing toward the peritoneal coat of the

uterus, these symptoms may be entirely wanting, and pain usually present between the periods, and especially severe at the periods, may be the only symptom present. Meadows has observed that the location of the pain is much determined by the location of the growth: "If the pain is felt in the lower dorsal or upper lumbar region, then it is probable that the tumor is growing on the fundus uteri. If, on the other hand, the tumor is more confined to the body of the uterus, then the pain will be felt in the lumbar region above; and lastly, if the cervix be the seat of the disease, the pain will be felt mostly over the sacral region."

Pain due to ovarian disease, like the neoplasm of the ovary, is to one or the other side of the central line; due to uterine neoplasm, it is usually in the central line. Small ovarian neoplasms are not always accompanied by menstrual disorders and irregular discharges. Where the little tumor is subperitoneal, and pain and displacement alone point out a suspicion of its presence, we must wait until further development occurs. Later, the tumor may be made out, and the differential diagnosis from chronic metritis and pelvic cellulitis may be at once determined by the absence of sensitiveness to the touch of the finger; hematocoele may be ruled out by the absence from the history of its sudden invasion with symptoms of shock. The slow growth of the tumor, its irregularity and solidity, are characteristic of fibro-myomata. Should, however, amenorrhea be present, the density of the growing tumor, and possibly its irregularity, the asymmetrical condition of the tumor and the uterus together, the deviation of the uterus from the central line, the absence of changes in the mammary glands or vaginal mucous coat, the presence of a serous leucorrhea of a peculiar odor, will usually lead to a correct diagnosis. Supposing the tumor and the uterus to be still in the pelvis, the uterus may be crowded to one side; or, if pulled down by the descending tumor, it will be retroverted, while the cervix is crowded forward; or, if the tumor proceed from the anterior wall, the uterus will lie retroverted beneath or to one side of the tumor. If the tumor has become so large as to lie above the pelvis in the cavity of the abdomen, the uterus will be dragged up and its cavity lengthened. While the tumor occupies the pelvis the pressure is extended to the bladder and rectum, and both retention of urine and difficult defecation are frequent. The ureters may also be so interfered with by the pressure of the tumor and uterus as to convey a diminished amount of urine to the bladder: with this condition almost complete suppression of urine may occur, but after the tumor is lifted out of the pelvis the flow of urine becomes greater than normal for a day or two. The presence of free fluid in the cavity of the abdomen is not very frequent, but when it does occur by reason of the presence of the tumor, it may come from irritation of the peritoneum, but usually from the covering of the tumor itself, the soft or edematous variety. Adhesions are the result of patchy or general peritonitis, usually the former.

The mechanical irritation of the tumor produces peritonitis, which may involve that portion of peritoneum covering the tubes, and the adhesions which follow may shut off the canal of the tube; or the pressure of the tumor may shut off the uterine canal and sterility result; or the irritation of the tumor may produce so much congestion of the uterus as to

set up the condition known as chronic metritis, viz.: congestion with hyperplasia. When this is the case the symptom of pain is much increased, and bleeding will occur.

It is not uncommon at the menstrual period to find the tumors and uterus more enlarged than ordinarily, and in some cases at these times there is frequently retention of urine requiring the use of the catheter. We may recapitulate the symptoms of the solid, subserous fibro-myomata thus:

- A. Uterine displacement;
- B. Pressure upon or irritability of the bladder and rectum;
- C. Bearing-down pain and backache;
- D. Uterine tenesmus at the menstrual period;
- E. Serous leucorrhœa with or without peculiar odor;
- F. Pressure on the nerves and blood vessels;
- G. Increased depth of the uterine canal;
- H. Occasional retention of urine;
- I. Sterility;
- J. A solid, non-sensitive tumor,
- K. Mobility of the tumor and uterus together;
- L. Ascitic fluid in the abdominal cavity.

Of these symptoms, nearly all will be found in those cases where the growth nearly fills the pelvis, and nearly all of them will be wanting if the growth is too large to enter the pelvis, and therefore lies above it. In one case the patient may be suffering from nervous irritability and a disorder of all her functions to a considerable extent; she may be confined to her room, locomotion being painful. In another case, the pelvis being but little encroached upon or entirely empty, the patient may have no symptom of suffering, and may enjoy good health.

When the tumor and uterus are above the true pelvis in the cavity of the abdomen, their connection is usually determined by bimanual examination without the aid of the sound. If cystic degeneration has occurred, palpation may detect it, and aspiration will produce a fluid having in it the fiber-cell or a fluid coagulating on exposure to the air, or blood alone may be drawn. The cavity of the uterus will likely be found lengthened, and the connection between the tumor and the uterus may be determined by seizing the latter with a volsellum forceps, while an assistant seizes the tumor; alternate pulls by each other will determine a connection with the uterus, but it will not determine positively that the tumor has not had another origin and formed a uterine attachment. In such cases the proper method of diagnosis is by an exploratory incision, the operation for removal of the growth following immediately.

Interstitial Fibro-Myoma, Solid Variety.—As in the case of the subserous variety, the earliest symptoms will be uterine displacement with vague symptoms of but little certainty. As the tumor increases, if in the anterior wall, ante flexion at first, and later retro flexion, will occur. If in the posterior wall or at the fundus, retroversion will soon follow. The uterus in all varieties, by increased weight, partly due to the growth and partly to increased blood supply, will descend to some extent. As the tumor grows, dysmenorrhea, menorrhagia and leucorrhœa develop. The irritation of the growth hastens uterine tenesmus, and that organ, becoming congested and heavy, sags down in the pelvis. As the tumor is increased the uterine canal is pressed upon; the symptoms of pelvic engorgement spoken of in the

last section occur, and continue until the tumor and uterus, by reason of their size, are lifted up out of the pelvis. The introduction of a sound, or, better, bimanual examination, proves the growth to be the uterus enlarged by the tumor. Disordered blood flows from the uterus and leucorrhœa more or less serous are the other symptoms most generally encountered.

Fibro-Myoma of the Cervix.—Here the symptoms differ from the others in this, that menorrhagia is not so frequent, endocervicitis is more common, and the enlarging lip of the uterus is in sight if a speculum be used, and within easy reach of the finger. The differential diagnosis involves the malignant growths of the cervix, hyperplasia of the cervix following laceration, and inversion of the uterus. The tumor, if pedunculated, may be traced to its connection with the uterus. The consistency of the growth, its want of sensibility, the inability to separate it from the uterus by bimanual examination, the inability or difficulty of finding the os, the fact that it has lost its normal or usual contour, the slow growth of the tumor, the presence of the uterine fundus beyond, the absence of all cachexia, the fact that the growth is of sufficient size to impact the pelvis, will clear up the case. Should it be deemed necessary, however, an incision into the growth can be safely made and its nature fully diagnosed. Even should the tumor prove to be the inverted uterus, in the hands of a good surgeon no injury would result.

Submucous Fibro-Myoma.—The inception of this variety may be early followed by dysmenorrhea, menorrhagia, metrorrhagia, leucorrhœa and serous discharge, uterine tenesmus and displacement, the symptoms steadily increasing with the growth of the tumor. Occasionally, menorrhagia and metrorrhagia are wanting. The uterus enlarges, its cavity increases in depth, its walls become heavier, and because of increasing weight it sinks in the pelvis. Anemia, neurasthenia, dyspepsia, great depression of spirits and prostration often rapidly follow, and unless the patient is relieved of the tumor, either by nature or her physician, she will die. Should the patient not get rid of the tumor early, it may so enlarge the uterus by its growth that the former is lifted out of the true pelvis.

General Remarks.—Solid fibro-myomata are differentiated from ovarian cystomata and fluid accumulations in the cellular tissue of the pelvis by the entire absence of fluctuation, their slow growth, their connection with the uterus, and the marked derangements of the functions of that organ. From solid tumors of the ovary it is not always possible to differentiate them, and as already noted, I have seen both Billroth and Lawson Tait open the abdomen for a supposed uterine tumor, and find instead a solid ovarian tumor. The moving about of the solid ovarian growth may move the uterus, and give the idea of a uterine tumor with a pedicle.

The most difficult diagnosis is encountered in very small subserous and interstitial tumors. But time or the bimanual method of examination carefully employed, or pulling down the uterus with a volsellum and retroverting it toward the examining finger in the vagina or rectum, will sooner or later discover the small growth. If it be subperitoneal and located on the anterior wall, and producing symptoms of stone in the bladder, I see no objection to dilating the urethra, dragging down the uterus, passing the

finger into the bladder and examining both bladder and the anterior wall at the same time. The interior of the uterus may be reached with the finger only after thorough dilatation, which may be accomplished with Molesworth's dilator or the metallic dilator of Marion-Sims, or Gooddell, either after or before division of the vaginal portion of the cervix.

Previous to entering upon either of the above-described processes, the bowels should be completely cleaned out, and all pelvic congestion further relieved with antiseptic hot water douches, and the last may be continued as a safeguard after the procedure. Both ante flexion and retroflexion of the uterus have been mistaken for fibro-myoma. The uterine probe will locate the position of the fundus in either case, and careful manual palpation will further solve the problem. Fecal accumulations in the caput coli or sigmoid flexure would not follow the movements of the uterus, and would be affected by enemata and cathartics. Pelvic hematocele is of sudden appearance, accompanied by evidence of loss of blood; the tumor is fixed and painful to the touch.

Pelvic cellulitis produces a painful swelling which soon fixes the uterus, rendering efforts to move it painful; the temperature rises and examinations are painful; the vagina is hot and the pulse beats perceptibly in the vaginal arteries. These constitute the main clinical features, differential and otherwise, of solid fibro-myomata of the uterus.

Caution.—The uterine sound as an aid to diagnosis is beyond doubt valuable. Its use is entirely precluded in case pregnancy exists. It should be used with the utmost care and without the exercise of any force. The direction of the uterine canal should be first determined by means of a flexible probe, which may also be used as a substitute for the sound. An extensive experience in bimanual examination will enable any one to dispense with the constant use of this instrument with advantage.

(To be Continued.)

SICKNESS AND PAUPERISM.

Read before the Cambridge Society for Medical Improvement,
Nov. 22 1893.

BY L. L. BRYANT, M.D.

ASSISTANT CITY PHYSICIAN OF CAMBRIDGE, MASS.

There are over eight thousand histories of families who have applied for public aid on file in the office of the overseers of the poor of this city. This represents twenty-four thousand names, in round numbers which appear on the pauper books of Cambridge. To this list, names are continually being added at the rate of fifteen hundred a year. What are we, as a Society, doing to prevent this? The question has often been asked me: "What are you doing?" and I have many times propounded the same question to myself. This paper is written in the effort to reply. In the month of May, 1883, ten years ago, I was elected to the office of Assistant City Physician. The said office was created Dec. 1, 1880, to enable the local Board of Health to conform to statutory law and yet retain the incumbent City Physician, Dr. Henry P. Walcott, a man occupying an honored position in the State Board of Health, and standing deservedly high in the respect and esteem of the people, and of this Society, of which he is a member.

My predecessor, Dr. O. C. Turner, a most estimable

man, died in October, 1892, of typhus fever, contracted while in harness; and during his illness, and subsequently until my election the affairs pertaining to the office were satisfactorily conducted by my friend, Dr. Wetherbee.

With a nervous organization particularly susceptible to external influences, and with absolutely no knowledge of the refinements of modern economics as applied to the poor, I sought and found the office which I still occupy; and my only qualifications were a fair knowledge of general medicine, and that experience which usually follows the first few years of practice subsequent to graduation.

The gaunt, hollow-eyed, cadaverous poor, having nothing and wanting everything, I knew nothing about, and as little what to do for, or with. Entering the service of the city under the circumstances, and in the condition mentioned, I first made myself acquainted with the duties prescribed by the city ordinances.

I found the office of the overseers of the poor to be the abiding place of one of the most perfect systems with which I had ever been made acquainted, with its thousands of histories of individuals, each one necessary to the proper registration of an applicant for aid and the establishment of a legal settlement, numbered, indexed, and filed away in bound volumes. I found in this office all the usual machinery and labor-saving devices necessary to the conduction of a large business, and presided over by a Secretary, D. P. Muzzey, devoted to his work; and apparently constantly employed. I found that all appeals for assistance must be made to this Secretary, who was empowered by the Board to dispense the city funds at its command as it deemed for the best interests of the applicants, and then to so manipulate the official machinery in the interest of the city as to secure from the various cities and towns wherein they had resided long enough to acquire a settlement, a monetary equivalent for everything paid out and for all services rendered. I found that the fixing of a settlement on some other city or town was one of the most important duties of the office, and required the services of a so-called "Visitor," who, likewise, was apparently constantly employed. This labor necessitated calling on the applicant, asking for, and recording various personal questions, searching city and town records to prove the truth or falsity of his or her statements, and, in general, the making of such a return to the office as would enable it to form as accurate an idea as possible of what he had been, what he was, what he wanted, and "how we are to get anything back?" I found that the Secretary was the confidential clerk and advisor of the Board, the disburser of supplies, and came into more intimate relationship with the Board, the poor, and the general public than any one else, and conscientiously performed his duties.

The Visitor, Vespasian Danforth, enjoyed the distinction of being considered one of the best versed men in pauper law in the State. Both of these public servants were ever on the watch for fraud, and worked the machine, seldom allowing sentiment to change the order of its working. Not but they were susceptible, but experience continually taught them that yielding to it was an unprofitable indulgence save as a reflection from the Board. I found as a working formula that the overseers of the poor expected me to visit such of the poor of the city as the