ANATOMICAL TREATMENT OF FRACTURES OF THE FEMORAL NECK.*

C. E. RUTH, M.D.

KEOKUK, IOWA.

In the Association JOURNAL of August 28, 1899, I published a report of 17 cases of intracapsular fracture of the neck of the femur treated according to the method advocated and practiced first by Prof. T. J. Maxwell thirty years ago. It is what I have seen fit to denominate the anatomical method, and is the only method, to my knowledge, that entirely neutralizes the displacing tendencies of muscular action and weight.

Of the 17 cases reported 5 recovered with no discoverable shortening whatever on careful and repeated measurements, 7 gave one-half to three-fourths inch shortening, 3 with one inch shortening, and 2 cases with useless flail-like limbs in which the treatment was abandoned after a few days of trial for lack of co-operation on the part of patient and friends. One death occurred from complications after union had been secured.

Fig. 1.—Muscle pushing itself or other soft parts between the fragments.

Fourteen cases, ranging in age from 25 to 86 years, have obtained union with useful limbs, in no case exceeding one inch of shortening. Union was therefore secured without perceptible shortening in 35 per cent. of cases in which the treatment was continued with anything like regularity.

Slight eversion of the foot was noticed in one case, owing to the lateral pull being too slight and directed too nearly straight outward, instead of the heavy incline upward sufficient to control the tendency to eversion. In two cases of negative results in which treatment was abandoned after a few days, we have the usual outcome of non-treatment, while the amount of shortening obtained in the other cases is much less than

the average of those treated by simple extension or other device.

Case 17 was 60 years of age, and so feeble that but little hope was entertained of saving her life; yet she recovered without shortening, and now walks without crutch or cane, without limp or halt. I am convinced that in my earlier cases I could have secured union with less shortening had I known better how much weight was required to overcome muscular action and weight displacement.

The notoriously bad results following fractures of the femoral neck makes apology for this paper unnecessary.

The following cases I wish to report and offer as additional evidence:

CASE 18.—Dr. D., aged 28, was thrown 15 feet from a cart in August, 1894, and sustained a fracture of the right femoral neck. Shortening existed of two inches with eversion and external rotation, treated with extension, counter and lateral extension four weeks. Result, union with one-half inch shortening. Walks without limp in any manner perceptible. Physician, C. A. Runyon, Elvaston, Ill.

CASE 19.—Mrs. Nancy M., aged 68 years, in 1892 sustained a fracture of the left femoral neck. No union was obtained. Two inches of shortening exists with a useless limb, treated by the old sand-bag method, and eight pounds of extension. In

Fig. 2.—Course and relations of psoas and iliacus to hip-joint, with thigh flexed at right angles to the trunk.

November, 1899, she fell and fractured the right femoral neck. Two and a half inch shortening was found on examination. She was treated by extension, counter and lateral extension, and recovered in six weeks with seven-eighths inch of shortening. Can bear weight on the limb and walk with crutches as well as before fracture of the right femur. The left leg still dangles, as it has for the past nine years. This patient would have been confined to the bed or chair the rest of her life had she been treated for the second fracture as she was for the first. Physician, C. A. Runyon, Elvaston, Ill.

CASE 20.—Mrs. F., aged 88, sustained intracapsular fracture of the left femoral neck in 1894, and was treated by Dr. H. A. Kinnaman and reported by me in my former paper as Case No. 11. She recovered with a useful limb with one-half inch shortening. She fell Jan. 11, 1900, and sustained a fracture of the right femoral neck. Treatment was not begun for ten days, but was by extension, counter and lateral extension. The treatment was intermittent, owing to the patient being

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee of the Section: Drs. W. J. Mayo, H. O. Walker and A. J. Ochsner.

intolerant of restraint. The result was union in four weeks with about one inch shortening. She died two months later of intercurrent malady, nature not determined. Physicians, Drs. Gray and C. E. Ruth.

CASE 21.—Tim H., aged 80 years, habits intemperate. Fractured the left femoral neck by a fall on the sidewalk, Nov. 27, 1899. Treated as above. Union was obtained in five weeks; position and motion perfect; shortening one-half inch. Examined and measured leg, April 25, 1900. January, 1901, still well with good use of leg. Physicians, R. H. Fegers and C. E. Ruth.

CASE 22.—Mrs. Obediah H., aged 70. Sustained a fracture of the right femoral neck from a fall in January, 1901. Eight days later I was called, and treatment was begun as in the above-mentioned cases. Union was obtained in four weeks without discoverable shortening.

CASE 23.—Boy 10 years old fell twenty feet and sustained a fracture of the femoral neck. Treatment as in the former cases, with scarcely demonstrable shortening. Union was secured in four weeks. Physician, A. Ives.

The above physician has had two other cases, 73 and 86 years of age, in which union was secured, but death resulted from exhaustion or complications, and as I have not the histories, can furnish no farther report of them.

CASE 24.—David B., aged 60 years, April 10, 1900, was thrown from an ore wagon in a runaway, and thrown down the mountain thirty or forty feet. Examination under chloroform revealed fracture of the femoral neck. Treatment as in the former cases for six weeks. Union was obtained with one-half inch of shortening. He has for several months been driving the ore wagon, and walks with very little limp. Occasionally complains of some pain in the hip in bad weather. Physician, C. M. Tinsman, Adin, Cal.

CASE 25.—Mrs. Elizabeth F., aged 86, sustained fracture of the right femoral neck in January, 1900, with shortening, eversion and backward displacement of the trochanter major. Treated by Buck's counter and lateral extension. Union was obtained with no perceptible shortening. The fractured leg gives as good service as before the injury. Physician, Dr. F. A. S. Rebo, Alexandria, Mo., assisted by Dr. W. A. Rebo, of Canton, Mo.

The first indication in the treatment of all fractures is correct adjustment; second, immobilization; third, overcoming muscular action which would tend to disturb the normal relations. We should be careful to eliminate all our own errors before blaming Nature with our unsatisfactory results. In these cases there is no muscular effort or influence acting on the upper fragment. The position of the lower fragment will depend upon the position of the patient, the kind of retaining force, and his own muscularity; for muscular action is of more importance in determining results of fracture here than anywhere else in the body. This is due to the great number of powerful muscles passing over the fracture line to be inserted somewhere beyond, and which must exert a powerful displacing force.

The position, at least for a time, that must be taken by these cases is upon the back, and the displacement is known to be upward, backward and outward rotation, with all the internal displacement of the lower fragment which the muscular tissues will permit.

The pronounced tendency to eversion and external rotation is increased greatly by the external rotary power of the psoas and iliacus, which was neutral or else an internal rotator now must be a pronounced external rotator, because the insertion into the back and internal portion of the upper end of the femoral shaft and lesser trochanter throws the line of action internal to the point of resistance, whereas it was formerly external.

The result is that we have the following muscles acting to produce the following results:

Those pulling nearly directly upward and tending to shorten the limb by over-riding the fragments, are the sartorius, rectus, gracilis, semi-tendinosus, semi-membranosus, long head of the biceps, vertical portion of the abductor magnus.

Those pulling very strongly inward as well as upward are the adductor magnus, except vertical portion; adductor longus, adductor brevis, pectineus, gluteus maximus, gluteus medius and gluteus minimus.

Those which pull almost directly inward and rotate outward are the pyriformis, obturator internus, gemellus, superior and inferior, obturator externus, quadratus femoris, lower half of gluteus maximus.

Those which pull inward, upward and rotate inward, are the tensor vaginae femoris and the anterior portions of the gluteus medius and minimus.

Fig. 3.—Fracture of neck of femur in Case 2. At 72 years of age, used 20 years afterwards. No perceptible shortening. Physician, Prof. T. J. Maxwell, Keokuk, Iowa.

Those formerly internal rotators exerting a strong upward pull and now external rotators, often drawn directly between the fragments, and absolutely preventing approximation of the bony parts, psoas and iliacus (Fig. 1).

It will thus be seen that the external rotary force greatly predominates over the internal because of the conversion of internal rotary force of the psoas and the iliacus to external, besides the power of the limb's weight to accomplish the same ends. Lastly, we have the important displacing force weight tending to carry the upper end of the lower fragment behind the line it should occupy because its old attachment support is gone. Posterior displacement and external rotation are only limited by the Y ligament.

Adjustment of the fragments is best accomplished by flexing the thigh upon the abdomen to relax the psoas

and the iliacus, bringing them above the fracture line, thereby preventing them from being permanently caught between the fragments (Fig. 2).

This position also relaxes nearly all the external rotators or changes them into abductors; then make vertical traction on the shaft of the femur, which now stands at right angles to the trunk, while moderate eversion is being maintained; next abduct to the normal line and make extension in the long axis of the trunk, while an assistant makes traction one-half to two-thirds as strong outward, slightly upward and forward from the upper end of the femoral shaft. These manipulations should be made by firm, steady traction, not by jerks, which is to become continuous by Buck's extension with a weight of from ten to twenty pounds, according to the muscularity of the patient, with elevation of the foot of the bed enough to counteract the tendency of the patient to slide down to the foot. This means an elevation of from six to fifteen inches. A binder's board, or other splint material, should now be molded to the upper inner aspect of the thigh, over which a band of adhesive plaster four to six inches wide should pass outward, upward and sufficiently forward that the weight over the pulley

Figs. 4 and 5.—Vertical section of femoral neck fractured 20 years before. Case 2 of former report.

shall overcome the internal pull of all the rotators and adductors, and at the same time raise the lower fragment to its normal level and maintain it there. It must entirely overcome the tendency to external rotation.

The weight on this lateral pulley will be from five to fifteen pounds, according to requirements. The side of the bed corresponding to the injured side of the patient must be raised enough to prevent the individual from being drawn out of position toward the lateral pulley. Patients so treated will have absolutely no pain whatever after spasmodic muscular action is entirely overcome.

The claim has been persistently made that bony union can not be expected. In Case No. 2, treated by Dr. T. J. Maxwell in 1881, bony union was secured, as proven by postmortem specimen which I now present you (Figs. 3, 4, 5). This patient was 72 years of age when he sustained the fracture. He made a good recovery without perceptible shortening, and had good use of his leg for twenty years.

The specimen shows the line of union very distinctly. There was slight eversion, but not sufficient to cause the patient any marked annoyance. This case is sufficient

to forever settle the question of Nature's ability to repair these injuries if given a chance.

The figures I herewith present will show the actions of the psoas and iliacus muscles in fractures of the femoral neck; and as some might question the correctness of the photographs, I will illustrate by models to which I have attached artificial muscles (Figs. 1, 2 and 6).

The only essential illustration difference between the action in life and on the model, however, is the anterior portion of the capsular ligament will probably be intact to its lower border and prevent the muscular tissue from falling between the fragments; but muscular action tends to force the capsule in such cases between

Fig. 6.—Shows normal position, relation and line of action of years before. Case 2 of former report.

the fragments, and thus interfere with union quite as much as though the muscular tissue itself occupied the space.

We do not come to you with a single case in the hands of a man of slight experience, but with 25 cases of unquestioned fracture of the femoral neck in the hands mostly of men of large experience. In 22 of these cases union was obtained; useful limbs were secured in persons ranging in age from 10 years to almost the extreme limits of old age, and yet the shortening was no more than one inch in any; and after the most careful measurements no shortening whatever was discoverable in six of the cases.

In one of the cases reported with one-half inch shortening, the physician in charge informed me that he could find no shortening, and that the patient walked without a particle of halt; but one-half inch shortening was claimed in the prosecution of a suit brought against his employers for damages. This man fell thirty feet when the fracture was produced. In some of my own cases I am certain that I did not use heavy enough weights, and in others the treatment was interrupted occasionally by having the weight removed to satisfy some meddlesome attendant, nurse or the patient.

The only failures to secure union were in two cases that absolutely refused to have the treatment carried out. I might cite you many cases of flail-like limbs from this cause, as can be done by most physicians of any considerable practice. Where little or no shortening took place, and the limb was functionally perfect, there can be no doubt that the union was bony.

Whatever may be the opinions in regard to repair of this bone, results that are almost perfect, and make the patient comfortable while repair is taking place, are certainly worthy of our attention, endorsement and most earnest efforts to establish, even though most of the patients are advanced in years. The usefulness of this plan of treatment is not alone limited to fractures of the femoral neck, but is equally applicable to the treatment of hip-joint disease with or without operation in cases confined indoors.

In some of these cases treatment was not begun until several days after the injury, and in some the treatment was interrupted occasionally for a short time. In no case was pain complained of after the first day, if weights were enough to prevent muscular spasmodic disturbance of the fragments.

the time required for procuring union. It also proves that the influence of the synovia is nil as a preventive of union. I do not wish to be considered as claiming that perfect results can be confidently expected in all, or in fact any of these and thereby invite prosecution for malpractice in every case of failure. What I do want, however, is to see the day when we will be able to lay aside prejudice and custom, if need be, and use the method that will give the best results. By this plan of treatment considerable freedom of movement can be allowed the upper part of the body as well as the opposite limb; so that the confinement is usually not irksome after the first few days.

DISCUSSION.

DR. C. E. THOMSON, Scranton, Pa.—My patient, an old man, 62 years of age, sustained a fracture in August, 1899. He was a miner by occupation, and the injury was caused by a fall of rock. No diagnosis was made at the time of accident. He was bedridden for three months; at that time he came under my care. A diagnosis of fracture of neck of femur was made and afterwards confirmed by *x*-ray photographs. The operation recommended by Gillette, of St. Paul, was proposed, accepted, and performed on Dec. 12, 1899. It consisted of a

Fig. 7.—Showing Buck's extension and lateral extension. Foot of the bed raised; the side of the bed corresponding to the injured limb is also raised to overcome tendency to displacement towards the lateral pulley.

By this plan of treatment the patient can be raised, if needs be, to the sitting or semi-sitting posture for the purpose of cleansing. Some of these cases had bedsores, and were lying in their own excrement and suffering severely when the treatment was begun. These were raised daily to the sitting posture, cleansed and the sores healed during the use of the extension treatment.

The patient can be easily handled in the use of the bedpan. The method is easy of application, and only fulfills the indications we require for all other solutions in continuity of bone, viz., securing coaptation of fragments without the possible intervention of the soft parts, overcomes all displacing tendencies of powerful muscular action instead of ignoring the action of more than one-half the muscles and displacing power of the limb's weight (Fig. 7).

The results prove that it is not necessary to allow most of the fractures of the femoral neck in the aged to produce cripples for the rest of their lives, or that they should suffer, except from the confinement, and that could be scarcely less by the plan of treatment proposed than to give the patient every possible liberty during

horseshoe incision around the trochanter major down to the periosteum. The flap being elevated, a second incision including periosteum and a section of bone (about one-third thickness of shaft) $2\frac{1}{2}$ inches below trochanter was made, liberated with an osteotome, and turned up. This enabled us to get nearer the seat of injury. In this particular case the fracture was found to be comminuted. Several pieces of bone were removed. The fractured ends were then adjusted and a solid silver nail three-sixteenths of an inch in diameter and two and one-half inches long was driven through the neck and into the head of the femur, as the skiagraph shows. We were gratified to find that the nail held the fragments quite firmly together. The wound was closed with catgut suture and the entire leg and body encased in a plaster-of-paris spica, extending from the axilla to tip of toes. Patient had no elevation of temperature. The initial dressing was removed at the end of two weeks. At the end of five weeks the plaster-of-paris spica was removed and patient allowed to move about. He left the hospital on crutches and improvement continued. Seven months after operation he complained of pain, and a few weeks later a sinus appeared at seat of incision, and this was followed later by an accumulation of pus in the gluteal region, which was evacuated under cocain, but a slight sinus persisted. Improvement continued until last month (eighteen months after operation) which was the last time he was seen alive by us, when

he walked into the hospital without cane or crutch. He had 90 degrees of free and painless motion. Last Saturday night he retired as usual, and was found dead in his bed at 5 a. m. The coroner made a diagnosis of "heart trouble." I secured this specimen which is now being passed around. You will see, both by the specimen and the *x*-ray photographs, which

were taken directly on the plate, that the bony union is firm and complete.

DR. D. N. EISENDRATH, Chicago—I think we should discriminate between the different cases of fracture of the neck. The percentage of mortality is still very large and the reason for this is we do not recognize the importance of the fact of getting elderly patients up as soon after the fracture as possible. They die of bed-sores and hypostatic pneumonia; they are people with frail constitutions who can not stand confinement to bed, just as the last speaker showed us. The ideal treatment for these cases is in the use of the Thomas Ridlon hip-splint. It enables those people above the age of 60 and 65 to get up within twenty-four hours after the fracture. We confine these patients for 4 or 6 weeks in bed and then obtain no better results than obtained in ambulatory treatment.

DR. T. J. MAXWELL, Keokuk, Iowa—I had my first case of intracapsular fracture of femur in 1870. The case was a spare woman aged 52. I found the foot everted and leg shortened. I put her in a Day splint and made extension; she suffered pain continuously. At the end of a week there was flattening of the hip and I devised this plan or method that has been presented to you by Dr. Ruth. I put her under lateral extension and she recovered without shortening. Ten years afterwards I had another case—the bone that has been presented to you of an intracapsular fracture. However, this bone may not have been an intracapsular fracture. It might have been an impacted fracture, except for this: There were two inches of shortening, eversion of the foot and crepitation on extension and rotation. Now, the 25 cases presented to you ought to be evidence sufficient of the benefit; where there is no wearing, no pain after extension, the bed-pan can be used. This bone belonged to a man I treated 20 years ago. He was nearly 72 years old at the time of fracture. He had walked on it for twelve years.

DR. W. W. GRANT, Denver—This specimen is a most important one, if the man did have an intracapsular, impacted or

non-impacted fracture. In examining it I am forced to express a grave doubt of its having existed. The cancellous tissue of the neck and the compact substance of the under or lower surface do not reveal any evidence of disease or past lesion whatever. The upper surface of the neck, near the trochanter shows a few osteophytes—evidence of old inflammation.

I call attention to the fact that the neck of this bone, from a man 90 years old, is of normal length. If there was an impacted fracture, how could you explain it? If, as stated, there was immediate shortening of two inches or more, with loud crepitus, it would be equally difficult, if not impossible of satisfactory explanation. Fracture of the rim of the acetabulum,

Army Medical Museum.
Rarifying osteitis with interstitial absorption.

with dislocation, would, with the extension treatment used, better explain the condition past and present.

There is a specimen in the Army Medical Museum at Washington which involved the very question at issue. I have examined it repeatedly and carefully. It looks something like this specimen, showing more evidence of the influence of pressure in walking. The neck is shorter and the angle more acute. The man, an old soldier, was treated for a fracture of the neck many years before his death. Dr. B. E. Frayer, a retired army surgeon, now of Kansas City, made the postmortem and sent the specimen to the surgeon-general as an

Army Medical Museum.
Rarifying osteitis. Front view.

illustration of bony union. A Philadelphia surgeon, visiting Washington, examined it. The specimen was sawed lengthwise and disclosed satisfactory evidence that no fracture had existed. It was regarded as evidence of interstitial absorption from traumatism, notwithstanding the doubt entertained by some of such a condition.

Yet, there seems no doubt of a dry form of inflammation, with fatty degeneration, and change of angle of neck, the latter produced or promoted by walking and absorption. Such a condition, with or without injury, seems more rational than the conclusion that the specimen here presented—perfect in coaptation of fragments, bony union and outline—was a complete fracture of the neck.

The condition involves deeply not alone the question of treatment and the interest of the patient, but in a medicolegal sense, the welfare and reputation of the surgeon.

CHAIRMAN—How old and how many years between the fracture, in the Washington specimen, and death of the patient?

DR. GRANT—He was about 44 at time of injury and he died 10 to 20 years afterwards, I believe.

DR. MAXWELL—How can he account for two inches of shortening with crepitation on extension and rotation.

DR. GRANT—You might have it in the acetabulum without fracture of the neck of the bone and that extension would reduce it and you would have no shortening. I have often seen that crepitation mistaken for fracture of the neck of the bone.

DR. A. C. BERNAYS, St. Louis—When I saw that specimen at the Tri-State meeting four months ago I was skeptical; I thought it was not the bone of a man 90 years old, or a woman. I was inclined to doubt it as has been done by others this morning. Dr. Maxwell showed me the specimen at Keokuk. The thing impressed me so much that I got the superintendent of the poorhouse to let me have the bones of two of the oldest men that died during the month. They were over 80 years of age; I would not state now the exact age. I saw bones there just as solid in texture and as firm as that bone is. I became convinced that the bone is the bone of a man over 90 years of age. I simply want to bring that point out, because it is no more than just to the gentleman who presented it and to Dr. Maxwell.

DR. C. E. RUTH, Keokuk—I thought that this specimen might be questioned, and referred to that fact in my paper

Normal arrangement of bone columns. Gray.

because the result is so nearly perfect. If this fracture had occurred in the hands of a man of slight experience and ability, I might question the diagnosis of a fracture of the neck of the femur with a shortening of two inches and complete eversion of the foot; but Dr. T. J. Maxwell, a man of unquestioned ability and enormous experience, had this case, and now in regard to this specimen twenty years' time has elapsed. The man was vigorous and muscular; that shows by the prominence of the lines, tuberosities and tubercles. Again, I said the line of solution of continuity shows. It does show, but not as pronouncedly as in a case of marked displacement of the fragments; consequently, there is no large callus to mark the line of solution of continuity. I said there was eversion of the foot following the injury until the patient's death. Can you question that fact when I can barely pass this lead pencil between the great trochanter and the head of the femur? Did you ever see a femoral head in a well-developed individual as close to the great trochanter as that in a bone not previously broken? I never did, and I have examined a considerable number. This man walked with eversion of the foot from the time of his recovery from the fracture until his death, when it was removed postmortem. The only point where you can see evidence of solution of continuity externally is on this roughened line on the anterior surface of the neck nearer to the head than to the intertrochanteric line. Did any of you ever see so prominent a roughened line on the front side of the femoral neck between the intertrochanteric line and margin of the

cartilage in a femoral neck not previously broken? Not by any means. Why? Because there is no capsule or other tissue requiring a roughened surface for attachment to the anterior part of the femoral neck. Why do we have evidence of solution of continuity on the anterior surface indicated in this roughened line if it does not show on the posterior surface? The reason is, we had an eversion of the foot and there was not so close an approximation of the fragments on the anterior surface as there was on the backside, and then the fracture line ran so close to the base of the neck at the great trochanter that external union is not discoverable. Many of you have seen a union of bone not discoverable by examination of the surface when adjustment was perfect. We have made a longitudinal section to show the condition of the bone inside. You can see the position from this point to behind the center of the bone, the dense compact tissue extending along the line of fracture in a curved direction [illustrating diagrammatically]. Let this represent the head, the great trochanter and the lesser trochanter. Now, here we have the solution of continuity running across about this line [indicating] as we have indicated by the roughened line on the front side. When we section the bone we find that the line does not run through all the thickness in the same position or course. We find here dense, compact tissue extending from the inner side just above the lesser trochanter beyond the center of the neck, where we have the more pronounced projection of compact tissue (callus). This is not the only specimen we may obtain. We

Heavy shading indicates position of compact tissue from callus, broken lines between indicate cancellous tissue devoid of columns; other lines practically normal, save neck has been drawn much too long.

have had twenty-five cases in which the diagnosis was beyond question. There is one case reported by a physician where the fractured leg has a better function than the other, and this patient was 86 years old at the time of the accident and there is no perceptible shortening.

In regard to pinning of the bones, as was mentioned by Dr. Thomson, this beautiful specimen shows bony union. A skiagraph has been taken and it shows three-fourths inch of shortening. This plan was all right if it was necessary. I do not say it is not necessary in every case; it has not been in my experience. It is certainly a good method if we have no better. I do not say we should never use it, but when we can avoid an operation in these cases we should do it, and not add an additional burden to an already serious condition. Unquestionably, fracture of the femoral neck in an aged person is serious. If we can avoid operation complications it is an important matter, and should always be done except in cases of delayed or non-union; then the pinning is certainly good surgery. I have not much experience with the Thomas splint. It would be all right if it held the femur out and forward as well as down. It makes no provision whatever for the tendency of the upper end of the lower fragment to drop behind the normal line when the individual lies on the back, as he will part of the time. It makes no attempt whatever to bring the femur out in its normal position. The trouble has been, in the treatment of these fractures, that we make our extension directly downward, instead of in the line of the neck of

the femur. You would not think of applying this principle of opposing muscular action to any other part of the body as here. We seem to forget that some of the muscles pull inward, while others pull directly upward; so it is absolutely necessary that we apply two forces, and that we overcome the muscular action so that the resultant of the two applied forces shall be in the direction of the neck of the femur and give the result of one acting in the long axis of the body, and the other in an outward direction.

HERPES ZOSTER OPHTHALMICUS WITH BRIEF REPORT OF FIVE CASES.*

WM. C. BANE, M.D.

Clinical Professor of Ophthalmology and Otology, Denver College of Medicine; Ophthalmologist and Otolgologist to St. Joseph's Hospital; House of the Good Shepherd; National Jewish Hospital for Consumptives; C., R. I. & P. Ry.; Consultant for Diseases of the Nose and Throat to the State Home for Dependent Children.

DENVER, COLO.

Herpes zoster ophthalmicus has been well known for the past third of a century. It is an acute inflammatory affection appearing in the course of the first and second branches of the fifth nerve, and is almost invariably unilateral. It occurs in elevated groups of firm oval or circular vesicles, on an edematous base, at the terminal points of the nerve filaments. The groups of vesicles do not all appear at once and their area is limited to the median line, and to filaments of the branches involved in the inflammation.

Herpes zoster ophthalmicus is a neuritis in which the ganglion of Gasser is involved, as has been proven by De Wecker. Charcot and others have demonstrated that in herpes zoster the ganglion of the posterior root of the spinal nerve is inflamed.

The disease is characterized by severe pain, which usually precedes the eruption by a period varying from hours to several days. The pain is generally sharp and shooting, and interferes with sleep and labor, but sometimes the pain is dull and causes less disturbance.

In the ophthalmic type, the first branch is the one most frequently involved, which includes the upper lid, forehead as far as the scalp, and also the nose. The second branch includes the lower lid, superior maxillary region, upper lip, and over the malar bone.

When the nasal branches are involved, the cornea is very likely to suffer, though exceptions to this rule have been observed.

The error of diagnosing herpes zoster ophthalmicus as facial erysipelas has frequently been made. Some of the typical features of the two diseases are as follows:

HERPES ZOSTER OPHTHALMICUS.

1. It develops as a bright red inflammation limited to one side of the face, and appears in clusters of firm vesicles on an edematous base, that does not spread when once developed.

2. The eruption is preceded for hours or days by severe neuralgic pains, and as a rule no fever or rigors mark the disease.

3. The inflammation being of a neuritic origin, it is limited to tissues where the nerve filaments terminate.

ACUTE FACIAL ERYSIPELAS.

1. It develops as a purplish-red inflammation that spreads over all parts, without regard to the median line; the vesicles, when they appear, are in large blebs.

2. It is ushered in by a chill, and marked elevation of temperature, also a burning sensation in the inflamed tissues. A dull headache may be present.

3. The inflammation being due to infection by the streptococcus erysipelatis, spreads through the lymph channels, and when deep invades the lymphatic glands.

4. The inflamed points are exceedingly painful to the touch.

5. Groups of punched-out scars are invariably left in the wake of the vesicles and remain through life.

4. The inflamed tissue is moderately tender to the touch.

5. Seldom is there a trace left after the exfoliation of the skin, though sometimes a thickening of the cellular tissue remains.

The following cases of herpes zoster ophthalmicus have come under my observation:

CASE 1.—On Jan. 8, 1891, I saw Mrs. H. C. K., in consultation with Dr. C. Jane Vincent of Alleghany City. Some seventeen days previous to my visit, the patient had developed herpes zoster ophthalmicus of the left frontal and temporal regions. The patient had suffered severe pain in the head and the left eye. The vision was reduced to counting fingers at two feet. She had considerable congestion of the eyeball, and a small central ulcer of the cornea. The iris was free and responded to light and accommodation. One week after my first visit the ocular congestion had subsided, the corneal ulcer had healed and vision so improved that she could see the features of a face a few feet from the eye. No record was made of any vesicles on the nose.

CASE 2.—Rev. J. S. T. M., aged 65, consulted me May 5, 1891. He stated that one month previously his left eye had been injured by a bit of steel. His family physician removed the foreign body three days after the accident. He then developed what was diagnosed facial erysipelas, the disease being confined to the left side of the forehead and nose with involvement of the left eye. He gave a history of having had three corneal ulcers and iritis of the left eye. Upon examination I found a moderately congested eyeball and a slight opacity of the cornea marking the location of the ulcers. There were also several pigment deposits on the anterior capsule of the lens, marking the points where the iris had been adherent. On the forehead, the supra-orbital ridge and the nose were the scars where vesicles had existed. Vision of the right eye was 6/12, and that of the left eye 6/18. The patient remained under my care for about a month, during which time his vision improved and the congestion of the eyeball in great part disappeared.

CASE 3.—In February, 1896, I was called by Dr. J. W. Higgins to see Mrs. J. W. P. The patient was a delicate woman of 56 years of age. She was suffering with intense neuralgic pains in the left side of the head and face. Her face presented the typical appearance of an aggravated form of herpes zoster ophthalmicus, involving the first and second branches of the trigeminus. The area involved was of a bright-red color with large groups of vesicles on the forehead, nose and cheek. The left eye was moderately congested and painful, but no involvement of the cornea existed. The neuralgic pains were excruciating. Two weeks after I saw the patient she became insane and was taken to an asylum, where she remained for four years, finally dying without regaining her mental balance. Upon inquiry of her husband I learned that she had manifested some evidence of mental disturbance for a month previous to the development of the herpes zoster.

CASE 4.—E. A. P., aged 36, a nurse, was referred to me Nov. 30, 1899, by Dr. H. T. Pershing, on account of having some irritation of the left eye. Patient came to Colorado eight weeks previously, owing to some pulmonary disease. Two weeks after reaching Denver he was taken with chills, having three or four daily for a week, with a temperature of 100 to 104. On Oct. 24,

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Ophthalmology, and approved for publication by the Executive Committee of the Section: Drs. J. A. Lippincott, Casey A. Wood and H. V. Würdemann.