the percussor adjusted to it, and five moderate taps were given with the hammer, which crushed the stone, and the operation was finished in a few minutes. Our patient then expressed his intention of going to the harvest field, but he was of course put to bed, and kept there.

"The bladder was washed out, when some particles came away, and in a few days many fragments passed with the urine. He has now resumed his employment, being quite recovered, and free from any inconvenience.

"It may be observed, that this operation was unattended with any pain, except that arising from the injection of the bladder. The patient was not sensible of any increase of pain on the crushing.

"ED. FRED. DEHANE, M.R.C.S."

Remarks.—Having within the last six years performed the operation of lithotomy thirteen times, and one case only of them being unsuccessful (an encysted stone being removed), owing to an abscess having burst among the bowels, from the kidney, on the third day, when the patient suddenly sunk, my experience entitles me to a hearing on the comparative merits of the two operations. The great evil of lithotomy is the imperfect restoration of the apparatus of the bladder; for let the case be as successful as it will, there most frequently is a defect in that organ, an occasional dribbling. This does not depend on the healing of the external incision in ten days, or five weeks,—on a small opening being made into the bladder, or an extended one; neither on much or little violence being exerted in the extraction, or whether the gorget or the knife, or the straight, the grooved, or the curved staff is used.*

If the opening be made very low down in the membranous part under the pubes, the upper part is left whole, as it ought to be; still in all these cases we do not make the valvular apparatus quite perfect, except in the horizontal position in bed, where the bladder is perfectly retentive. It is the hanging down of the penis that favours the jet, and this would suggest its being kept with the glands confined

upwards. With children, however, it would be impossible to enforce this.

Do away with this evil, and lithotomy will not yield to lithotritry, except in the early formation of stone. In such a case as that of "William Corns," it is a beautiful adaptation of mechanism to surgery, its success triumphant; and if stone were never suffered to get large, the application of lithotritry would render lithotomy an unnecessary operation.

Walsall, Sept. 14, 1834.

EXTERNAL EMPLOYMENT

OF

PITCH IN LEPROA VULGARIS.

To the Editor of THE LANCET.

SIR,—The review of a work in your last Number (page 909), which recommends the internal and external employment of pitch in the lepra vulgaris, has induced me to relate briefly the two following cases, which are strikingly illustrative of its efficacy when employed externally in that frequently intractable affection. I am, Sir, your obedient servant,

R. HUTCHINSON, M.D.

Nottingham, Sept. 18th, 1834.

Case 1.—Mary Barton, æt. 48, unmarried, of Newark, in this county, was admitted into the General Hospital, March 19th, 1833. She has been affected for the last four years with an eruption upon the extremities and body, for which numerous remedies have been ineffectually tried; she has been three times under the complete influence of mercury without benefit. The arms, legs, and trunk, are now nearly covered by a scaly eruption, of the distinct character of lepra vulgaris; the scales in many parts are very thick and hard, the fissures between them deep, and discharging an irritating ichor; she complains of no pain, but a disagreeable tingling sensation; the eruption occupies every part of the legs and arms, with the exception of the palms of the hands, soles of the feet, and flexures of the joints, and no portion of the whole trunk retains its natural colour. The young woman's health is good, her face florid, and complexion light.

She was first ordered a vapour-bath to soften the scales; to continue in bed afterwards, in order that the *ung. picis* might be applied over the whole eruption, to remain there, and to repeat the application every night and morning. Aperients and slightly alterative pills to be taken every

* I came to the conclusion how desirable it was to open the urethra very low down, from the external wound readily healing. The inquiry arose from observing some cases healing in ten days, while others were protracted to five weeks. In some cases the bladder was retentive in three days, and the natural passage restored. Then again, how much violence the perineum will suffer, and yet get well, whilst the upper part of the urethra, from slight causes, occasions great constitutional disturbance! In a conversation with Mr. B. Cooper on this subject, he agreed in opinion with me, and referred me to his opinions thereon as stated in his work.

night, and some lac sulphur and supertartrate of potash in the morning, occasionally, to keep the bowels regular; to have a vegetable and milk diet.

March 23. The scales are entirely removed; the usual redness of the skin remains, but the leprous character is quite gone. Contin. remed.

My reports from this day to the 23rd of April state, that the natural colour of the skin gradually returned, when she was discharged quite free from any complaint.

Case 2.—G. Starkey, æt. 19, shoemaker, admitted August 4th of the present year. For the last five weeks an eruption has been gradually spreading from the extremities to the trunk; at the present time it occupies nearly the whole body; the arms are swollen, and the scales so thick as to interfere materially with their functions; the fissures on the backs of the hands particularly deep, with an inflammatory appearance, and somewhat tender to the touch; has much tingling pain in the parts; has occasional slight rigors, with uneasiness at the stomach; the pulse is rather full, and the bowels are regular. The eruption has the true character of *lepra vulgaris*. Ordered V.S. ad 3viii, vapour-bath, aperient pills at night, *Decoct. Sarsæ*, co. to take during the day; to remain in bed, and apply extensively and freely the *ung. picis* upon the whole eruption.

Aug. 8. The scales fallen off, the usual discoloration of the skin is seen; bowels confined. Contin. remed. Capt. *Pulv. Purg.*, mane 2nde quaque.

10. Much improved; feels in health well.

28. Discharged quite well.

Remarks.—These two cases may be taken as fair specimens of two stages of lepra, which may be called the acute and the chronic. In the second case the acute or inflammatory stage was clearly evident, and the disease would not, I believe, have been so speedily removed, unless depletion had been had recourse to. In the first, which I would call the chronic, on account of the absence of pain or tenderness, and from its long duration, mercurialization had invariably aggravated the eruption, and no application had entirely removed it, until the tar was tried. Although lepra has great tendency to recur, I have not heard that it has done so in this case. In the second case, sufficient time has not elapsed to ascertain if the cure be permanent. In both, the beneficial results of the application of the *ung. picis* were extremely gratifying and strikingly evident. Perhaps, something may be attributed to the necessary remedies employed,—to the bleeding and the purging

in the second case, to the alteratives in the first; but in a disease so constitutional as I believe lepra to be, no one could be justified in neglecting the employment of remedies calculated to improve that state of the digestive organs which most probably is the proximate cause of its existence, acting upon a constitutional and probably hereditary predisposition.

VAGINAL CÆSAREAN OPERATION,

WITH OTHER CASES, OCCURRING
IN THE PRACTICE OF

MARTIN A. EVANS, M.D., *Galway*.

ABOUT three months ago I was called to see Mrs. O'Maley, æt. 27, residing about five miles from this place. She had been then nearly forty-eight hours in labour, and on examination I found that scarcely any portion was perceptible which might be termed the os uteri. There was a general round mass—the uterus, enveloping the child's head, presenting itself in the upper part of the vagina; and it was only after a protracted search that I could discover a slight depression, the size of which scarcely equalled the diameter of a very small split pea.

The midwife was confounded, having never met with such a case before, and she informed the woman's friends that no hope remained, save that of cutting into the uterus through the abdomen, and thus extracting the child. A vague belief of recoveries effected in this way prevails pretty generally in this district. To execute this purpose I was sent for. The patient herself was a woman of strong nerve and frame, and unceasingly insisted on my commencing the operation, her pains being extremely severe. However, notwithstanding her entreaties, and those of her friends, I postponed any operation for twelve hours, and until I had given a fair trial to nature, having, during that interval, bled her, and found everything unavailing. The pains, which at the early part of my visit were so powerful, now began to flag; her pulse became hurried and weaker, and there was no alteration whatever in the state of the os uteri.

Under these circumstances I commenced the operation, by introducing the index-finger of the left hand, and fixing it at the small aperture. I then passed, with my right hand, Sir A. Cooper's hernia knife, and, using the index-finger as a director, divided the os uteri in various directions, each cut being about an inch in length. There was not much hemorrhage; the