paratively tender foliage. Where similar conditions prevail in Washington and its vicinity, a third generation of larvæ may develop, though small in numbers, but the writer is convinced that even in Washington late developing beetles of the first generation may hibernate.

Remedies.—The only thoroughly satisfactory safeguard against this insect consists in spraying the trees with an arsenical solution. The only other remedy which is worthy of mention is the destruction of the larvæ at the surface of the ground before or after they transform to pupæ. The latter remedy, however, is not complete, and even where it is carefully carried out for every tree in a city, it will do no more than reduce the numbers of the insects by perhaps two-thirds.

Ten years ago a proposal to spray the enormous elms which are to be found in many No thern towns would have been received with ridicule, but of recent years the practicability of the plan has so frequently been demonstrated that there is no hesitancy in commending it to more general city use. Probably the largest elm tree in America, the Dexter elm, at Medford, Mass., has been successfully and economically sprayed by the Gypsy Moth Commission. It is necessary to have especial apparatus constructed, and it is equally necessary to have the work done by men who are accustomed to it or at least are good climbers. The first successful work of this kind was probably that done by Prof. John B. Smith on the campus of Rutgers College. He had a

which they eat, not only will a great deal of leaf perforation by the beetles themselves be prevented, but the number of eggs laid will be very greatly lessened. A second spraying should be conducted two weeks later. This is directed against the larvæ, the majority of which will perhaps have hatched by that time or soon after. A third spraying, and even a fourth, or under exceptional circumstances a fifth, may be required if it is considered necessary to keep the trees fresh and green, and particularly if the earlier sprayings have been followed by rains, as is apt to be the case in the earlier part of the season. In Bridgeport, Conn., where only a part of the trees are sprayed, and these by private enterprise, an even greater number of operations have been found desirable. Three thorough sprayings of all the trees in a given precinct will probably be as much as will be required, especially if this be done year after year and some pains be taken to destroy such of the larvæ as may successfully develop and descend for transformation. Even two sprayings, covering all the elms of a city or town, will be well worth the expense.

The substance to be used in these spraying operations may be Paris green, London purple, or arsenate of lead. The directions for the use of these substances have been so often repeated that it is not worth while to mention them here.

The other remedy—the destruction of the descending larvæ and the quiescent pupæ—is, as above stated, and

FREAKS, AS PERTAINING TO DISEASES OF THE SKIN.*

By JAMES C. MCGUIRE, M.D., Washington, D. C.

IT is not alone in post graduate schools and hospitals we should look for clinical material; the dime museums of our large cities, the side shows of the circus, furnish excellent advantages for instruction. Especially is this true as regards skin diseases. Many anomalous examples of cutaneous lesions, that would be difficult to find elsewhere, are here seen to their best advantage; for, as a rule, the disfiguring lesions have not been treated in any way; more likely still are they to have been cultivated, till they are in full bloom. It may be of interest to cite a few of these cases and—instructive from a scientific standpoint—to properly classify them as diseases.

The popular names given to many of these deformities are given them on account of their resemblance to the skin of lower animals, as the "alligator boy," the "elephant man," "elephant leg," the "fish-skin girl," the "dog-face boy," the "leopard boy," etc. Others are designated as the "elastic skin man," the "bearded lady," the "blood sweater," the "albinos," the "man with horns," the "blue man," etc.

The "alligator boy," the "fish-skin girl," the "porcupine man," the "hedgehog man," all suffer from different manifestations of the same disease. The appearance of the skin so closely resembles that of fishes it is designated as ichthyosis; it is a congenital chronic disease, characterized by dryness, harshness, and scaling; usually showing itself within twelve or sixteen months after birth. In the milder form, known as xeroderma, there is only slight scaling, but the natural furrows of the cuticle are much exaggerated. In some pronounced cases, the skin is divided into diamond shaped figures, and scales appear in abundance—white or dark green in color—and are especially adherent in the center. These are the cases to which the terms "fish skin," "snake skin," are applied.

In the most severe manifestations of the disease, designated as ichthyosis hystrix, the hypertrophied skin may present dark, corrugated, wrinkled masses, resembling the bark of a tree rather than fish scales. There may be spinous or quilled excrescences, as in those cases designated as "porcupine men," "hedgehog men."

A very remarkable case of ichthyosis hystrix was shown at the Royal Society, March 16, 1731, reported by Dr. McColl Anderson.† The patient was fourteen years of age. "His skin (if it might be so called) seemed rather like a dusky colored thick case, exactly fitting every part of his body, made of rugged bark or hide, with bristles in some places. It did not bleed when cut or scarified, being callous and insensible. It was said he shed it once a year. It is not easy to think of any sort of skin that exactly resembled it; some compared it to the bark of a tree; others thought it looked like seal skin; others like the hide of an elephant. The bristly parts, which were chiefly about the belly and flanks, looked and felt like the bristles or quills of a hedgehog, shorn off within an inch of the skin. The patient was again shown twenty-four years later by Mr. Henry Baker, still presenting the same deformity. He had had six children, all of them presenting exactly the same deformity. 'It appears, then,' said Mr. Baker, 'past all doubt, that a race of people may be propagated by this man having such rugged coats or coverings as himself.'"

Dr. G. H. Fox reports‡ an interesting case of ichthyosis—the "alligator boy." The article is accompanied with two excellent colored plates.

There is still another rare form of the disease, known as ichthyosis congenita. Some authorities—Hebra, Kaposi, and others—regard it as a distinct affection, consisting in a perverted secretion of the vernix caseosa. It is said to begin at an early period of intra-uterine life. At birth, the skin appears too tight, for the body is deeply fissured and hypertrophied, resembling plates of armor; to this form of the disease the term harlequin fetus has been applied.

One of the most common of the freak family is the albino, a name given to those who suffer from a partial or complete loss of pigment, of congenital origin: it is said to be seldom met with in the white race; not only is there loss of pigment from the cutaneous covering, but the hairs and choroid coat of the eyes are affected as well. The white mice and birds of this climate are examples of albinism.

The "leopard girl" is affected with leucoderma, a word derived from the Greek, meaning white skin; the term vitiligo—a blemish—is also applied to the deformity. It differs from albinism from the fact that it is acquired and not congenital. The loss of pigment occurs in patches more or less numerous, that may run together till the whole skin is affected; the hairs on the affected patches also turn white. As to causation, it is said to be due to "disturbance of innervation." Many cases have been on exhibition; and it is not unusual to see it on the hands and face of negroes in this city. In the Louisville Medical News, 1880, Dr. Hall refers to a dark mulatto who became perfectly white, except a patch on the chin.

Leucoderma also affects the lower animals, as in the case of the "sacred white elephant." They are not really white, but of a pale flesh color. We will not look upon these sacred animals with such reverence when we learn the condition may be produced by artificial means. Dr. Piffard has witnessed the experiment of bleaching the skin of the ordinary elephant, by means of peroxide of hydrogen; it was applied several times a day; at the end of four days, the hide had changed to a light ash color, but he did not believe it would be permanent.

There have been several elastic skin men on exhibition; the best known of them is Herr Hagg, who could pull the skin from his chest up to his eyes. Meekren, as long ago as 1657, records a patient who could bring the skin of his chest up to his eyes and down to his knees. These cases are a form of dermatolysis, a rare deformity, that is owing to some defect in the attachment of the cuticle. It is a congenital affection, though it is said the ability to move the skin is increased by cultivation. Other forms of the disease occur in the

FIG. 3.—The Imported Elm Leaf Beetle (*Galerucella luteola*). a, foliage of European elm, showing method of work of beetle and larva—natural size; b, adult beetle; c, egg mass; d, young larva; e, full grown larva; f, pupa—all greatly enlarged; g, mouth parts of full grown larva—still more enlarged (original).

strong barrel pump, and carried the nozzle at the end of a long rubber tube, with a bamboo extension pole, up into the center of the trees by climbing a ladder to the main crotch. From this point the spray was thrown in all directions, and the tree was thoroughly coated with the mixture in a minimum of time.

The Gypsy Moth Commission, in their earlier spraying work, sent their large tank carts through the streets, stopping at each tree and sending one or more men with hose and extension poles into it, thus covering hundreds of large trees in a single day. If steam sprayers are used (and the town or city fire engines can be and have been used to excellent advantage in this way), the necessity for climbing the trees may be largely avoided. By means of multiple discharge hose both sides of a tree, or even of two trees, may be sprayed at once, and the extent of territory that may be covered in a day is surprising. The elm trees in a small park may be treated economically and without much difficulty by two or three men with a handcart tank. This method has been adopted on the large grounds of the Department of Agriculture, with absolute success.

The writer's experience at Washington leads to the conclusion that it is important to spray trees once just after the buds have burst. This spraying is directed against the overwintered beetles. If a large proportion of these beetles can be destroyed by poisoning the leaves

must always be, incomplete. The standard kerosene emulsion, diluted one part to five parts of water, will destroy the insects in either of these stages. This has been successfully used in several New England towns the past season, particularly in New Haven. It must be applied to the base of the trunk and under the entire limb spread of the tree. The rough bark must be removed to a slight extent (the writer does not advocate severe scraping), leaving as few crevices as possible which may harbor the pupating insects. If a tree is very large, it will pay occasionally to climb into the main crotch and destroy such individuals as may have collected at that point. Experience leads us to estimate that on large trees not more than one-half to two-thirds of the larvæ reach the base of the trunk and transform at that point. The extent to which larvæ drop from overhanging branches has been questioned, and it is sometimes a difficult matter to decide. The city forester of Springfield, Mass., however, called our attention to a peculiar and eminently satisfactory case where the drooping branches of a large elm extended completely over a house, on the other side of which there were no elm trees. On the far side of the house beneath the tips of the overhanging branches, the larvæ and pupæ were collected in large numbers in the summer of 1895.

(To be continued.)

* Read before the Medical Society of the District of Columbia, February 19, 1896.

† Diseases of the Skin, 1887.

‡ Journal of Cutan. and Vener. Dis., April, 1884.

abnormal growth of skin, attended with thickening; the integument hangs in folds and overlapping layers. An excellent example of this is seen in plate No. 7, Van Harlingen's Handbook of Skin Diseases. All of the above cases have several peculiarities in common; the deformities are not amenable to treatment; the general health is, as a rule, not affected, and practically nothing is known of the causation of the deformities beyond the fact of inherited tendency.

The "blue man" may be suffering from a condition known as cyanosis, defined in Foster's dictionary as a "bluish discoloration of the skin from defective aeration of the blood, either temporary, as in asphyxia, or permanently, as in the subjects of some malformation of the heart, especially persistent patency of the foramen ovale; or the blue man's condition may have been brought about by the ingestion of nitrate of silver."

An extremely interesting case, occurring without assignable cause, is reported by Mitchell Bruce (Inter. Atlas of Rare Skin Diseases). The patient was a harness maker, 49 years of age; the condition affects the body generally. He suffers from intermittent pains of burning, shooting character, chiefly in the epigastrium and left lumbar region. Pains and discoloration commenced two years previous to his admittance to the hospital, and though under observation for three months, there was no change. There was nothing in the patient's occupation to account for the discoloration.

A few years ago a man exhibited himself in a New York museum, with a horn growing from his forehead; it resembled an animal's horn in respect to shape and size, but was of a dark grayish color, and rough and harsh to the feel. This condition, known as cornu cutaneum, may occur on any part of the body and be of any shape and number; they differ from similar appendages in cattle, from the fact that, in these animals, the horn grows only from osseous tissue. There is a specimen in the British Museum that is said to measure 11 in. in length by $2\frac{1}{2}$ in circumference. Lebert, in 1864, reports 109 cases he had compiled from different sources; in 12 per cent. of these cases it gave rise to consecutive epithelioma.

"Elephant leg," "Barbadoes leg," is the well known disease elephantiasis arabum. Kaposi defines it as an "hypertrophy of the cutis and subcutaneous cellular tissue, which is confined to some region of the body, due to local circulatory disturbances." The most frequent seat of the disturbance is the lower extremity; rarely are both limbs affected. At first there is a deep seated inflammation that leaves the parts in a swollen edematous condition. Every few months this is repeated, leaving the leg still more swollen, till it reaches such enormous proportions as even to interfere with locomotion; the surface may be smooth or warty; sometimes the bones are affected.

An interesting case allied to this disease is described in the British Medical Journal, December 11, 1886, under the name of the "Elephant Man," a most remarkable case of congenital deformity of the skin and bone. The bony exostosis on his frontal bone, combined with a deformity on the superior maxilla, gives a trunk-like appearance to the nose and upper lip, the profile of the face suggesting the profile of an elephant's head. "The tissue is very loose, so that it can be raised from the deeper parts in great folds. The cranial bones are deformed and overgrown, so that the circumference of the patient's head equals that of his waist. Bony exostoses spring from the frontal bone, the posterior part of the parietals and the occipital." The patient can give no family history of similar deformity, but declares his mother was knocked down by an elephant when bearing him. The papillary excrescences are increasing rapidly, and hypertrophy of the integuments of the right hand is causing it to become slowly crippled; general health good.

"Hide-bound skin" is recognized as scleroderma, a chronic inflammatory disease, characterized by a hardening and rigidity of the cuticle over areas of greater or less extent; the surface may be white, yellowish, or even bronze color; the skin may become shrunken and depressed; when limited in extent it is known as morphea, though some authorities consider the latter an independent disease. The disease is supposed to be due to "disorders of the nervous system."

There are some anomalous conditions of the integument which may not be out of place to refer to in this article; though, as a rule, the subjects do not exhibit themselves, they are none the less interesting. In my personal experience, I knew of a young woman, in excellent health, without blemish upon her skin, who exhaled an odor so much like violets it had been remarked upon; in consequence, she is in the habit of wearing violets, or to have an artificial perfume of that flower about her person. The odor is always accompanied with excessive sweating; the perspiration is so excessive she will soil several pairs of gloves during an evening's dance. The condition is usually designated bromidrosis (stinking sweat), but in this case a better name is osmidrosis.

Colored sweat, known as chromidrosis, is usually feigned, but there have been a few well authenticated cases reported. The condition may be due to the growth of micro-organisms; it may also be caused by the medicine the patient is taking, as in Dr. Taylor's case, when blue sweat occurred in a man taking iodide of potash. The condition is supposed to have been brought about by the reaction of the iodine in the perspiration and the starch in the patient's shirt. Occasionally we read in newspapers of hysterical females and religious fanatics "sweating blood;" this is known as hæmatidrosis. Kaposi declares this is not the true sweating of blood, "but the occasional non-traumatic exit of arterial blood from the pores of the skin." In his work on "Diseases of the Skin" he refers to a case observed by Hebra, who once saw a spiral stream of blood spurt from the opening of a sweat gland on the dorsum of a young man's hand. A few cases of phosphorescent sweat have been recorded, though it is said to be caused by fish diet. S. Pollitzer considers it not unlikely that the phosphorescence is due to action of phosphoric bacilli, many varieties of which have been described.

That this article may be fully up to date we will mention those cases in which objects, such as trees, are indelibly impressed upon the skin of those who have been struck by lightning. Prof. Wright, of Yale University, declares that these images are photographed by means of the cathode rays, of which we have recently heard

so much. Some years ago, a lady, living in New York, stated to me that the dark ring about her wrist was caused in this way. While she and her family were sitting at the dinner table, during a severe thunder storm, the window being open, "a ball of fire entered and knocked us all senseless; it was some time before I recovered from the shock, when I found the image of a bracelet which I wore was photographed upon my wrist."

It is, of course, a recognized fact that the hair may be whitened long before the time it would naturally occur from senile changes. Lord Byron, in the Prisoner of Chillon, refers to this condition:

"My hair is gray, though not with years,
Nor grew it white
In a single night,
As men's have grown from sudden fears."

History records, but on doubtful authority, that the hair of Sir Thomas Moore and Marie Antoinette turned suddenly white the night before their execution; that Henry of Navarre was so shocked on hearing that the edict of Nemours was conceded, the hair of his moustache whitened in the course of a few hours. Even if we do not accept these cases as authentic, there have been many others recorded in recent times by competent observers; so that sudden blanching of the hair under great mental emotion is considered a possible but extremely rare occurrence, by all authorities whom I have consulted except Prof. Kaposi, of Verina. He declares that though grayness of the hair may occur in a short time, he does not believe it can occur suddenly and in fully developed hair; the reported cases he considers are based on "errors of observation." Dr. Jackson and others declare the condition is due to the formation of air bubbles between the hair cells of the cortical substance; the air renders it opaque, so that the color of the pigment is obscured. Kaposi's views do not coincide with these; he believes grayness is due to loss of pigment, and does not consider the influence of fright or mental emotion of any kind cause gases to develop in the hair, and that these gases or air conceal the pigment; inasmuch as many normally colored hairs contain air.

Notwithstanding the negative views expressed by such an excellent authority as Prof. Kaposi, that sudden blanching may occur has been proven beyond a doubt, not only in man, but the lower animals as well. Sir Erasmus Wilson mentions the experiments of Sir John Ross. He exposed a lemming—a kind of rat—to the open air at the temperature of thirty degrees below zero; the next morning the fur of the cheeks and spots upon each shoulder had become white; within a week the animal was perfectly white except a small patch on the back.

There has recently been reported a case the reverse of canities, where gray hair suddenly became black.

There is also a rare condition of the hair in which there are alternate rings, one white, the other the natural color. In the London Lancet, 1881, Squire gives an account of a boy sixteen years old, one side of whose head was piebald like a tortoiseshell cat, the other side dark brown.

As has already been said, loss of pigments may occur elsewhere than on the skin and hair. The nails of the hand have been known to become as white as snow. Dr. S. Giovannini, in the International Atlas of Rare Skin Diseases, describes a case of this kind, which he calls canities unguium. An abbreviated report of the case is as follows: Coachman, 29 years of age; all the nails on the fingers are milk white in appearance, but otherwise normal; the condition first occurred during convalescence from typhoid fever, when twelve years old; the second and third toes of each foot are webbed (syndactylia), otherwise the patient is well formed and in good health. The condition is supposed to be due to penetration of air into the interior of the nails, practically the same condition that occurs in canities. Dr. G. P. Unna has reported a second case in the same Atlas. To it he has applied the name of leuconychia, preferring it to the term canities, which he thinks should be confined to blanching of the hair. The patient, twenty-six years of age, at first noticed the condition when a child. In this case the hair is also affected. The pathological changes consist in dryness and lack of luster of the hair, and of white rings of different sizes.

The "bearded women" that grace so many of the museums by their presence must not be left out of this collection of freaks. These females are afflicted with a deformity rather than a disease, known as hypertrichosis, or hirsuties, which simply means a growth of hair abnormal in amount, or grows in places where lanugo hairs are normally found. Chowns speaks of a boy eight years of age who had the whiskers of a man. Dr. Cummins mentions the case of a lady who was noted for the beauty of her face, whose body, from breast to knee, was covered with a profusion of black, thick, bristly hair. These cases and others are referred to in Dr. Jackson's Diseases of the Skin. He says partial congenital hypertrichosis is apt to be of the nature of nevus.

The distinction between the localized hypertrichosis and nevus is mostly made upon the color of the underlying skin. In the former case the skin is perfectly normal, while in the latter it is pigmented, and may be otherwise altered.

The Russian "dog-man," from Ziemssen's Handbook of Diseases of the Skin, is the best known specimen of homines pilosi. Exactly the opposite condition is the universal loss of hair, known as alopecia universalis. Dr. Ohmann-Dumesnil, in the quarterly Atlas of Dermatology, January, 1896, reports a case of this kind. All the hairs of the body were lost, even the fine downy ones known as lanugo hairs. The process was slow and insidious. He says: "As thoroughly a naked individual as could be imagined is a result of the effect of this universal process."

The last freak that will be referred to in this article is probably the most unique of all. It is known as phagmesis, in which feathers adorn the body instead of hair. Dr. G. T. Jackson, in his most excellent treatise on Diseases of the Skin and Hair, is responsible for the following statement: "Dr. T. Robinson cites a case of this nature which occurred in a boy who was exhibited at Bremen, and was reported upon in Bauerle's Magazine for 1831. The boy's head is said to have been covered with feathers in place of hair."—Va. Med. Semi-Monthly.

SELECTED FORMULÆ.

Restoring German Gilt.—The following varnish is useful for restoring tarnished German gilt picture frame moulding:

Gamboge 30 grains.
Dragon's blood..... 240 "
Powder the above and then add:

Turmeric..... 30 grains.
Shellac..... $2\frac{1}{2}$ ounces.
Sandarac..... $2\frac{1}{2}$ "

Place the whole of the above in a bottle, and add two ounces dry oil of turpentine; shake often, keeping in a warm place for fourteen days; filter, and add four ounces clear mastic varnish as above.—Canadian Pharmacist.

Remedy for Codlin Moth on Apple Trees.—One pound Paris green, one pound lime, 250 gallons of water; or, better still, equal parts of Bordeaux mixture and Paris green. Used as a spray.—Gardener's Chronicle.

A Matt Black for Iron.—According to the Revue Suisse de Photographie, a matt black surface on iron can be obtained by the use of the following solution:

Mercuric chloride..... 2 parts.
Cupric chloride..... 1 part.
Hydrochloric acid..... 6 parts.
Alcohol 5 "
Water..... 50 "

The article is carefully cleaned and immersed in the above, or a brush may be used for its application, after which it must be well soaked in hot water. A second application can be given if the color is not dark enough.

Nail Cleaning Liquid.

Tartaric acid 1 drachm.
Tincture myrrh 1 "
Eau de Cologne 2 drachms.
Distilled water 3 ounces.

Dissolve the acid in the water; mix the tincture of myrrh and eau de Cologne, and add to the acid solution. Dip the nails in this solution, wipe, and polish with a chamois pad.

NAIL POLISHING POWDER.

Fine putty powder..... 4 drachms.
Carmine..... 2 grains.
Otto of rose..... 1 drop.

Very carefully reduce the carmine with the putty powder, and rub together until a homogeneous mixture is formed; then mix in the otto. The nails should be dampened, and a little of the powder put on; then polish with a chamois pad.—Bulletin of Pharmacy.

Oxidizing and Blacking Bright Steelwork.—The following direction for oxidizing and blacking the bright work of steel in lieu of paint, to stand heat and to wear well, is taken from a recent issue of the English Mechanic: "Take three ounces of glacial acetic acid, mix it with its weight of water; to this add half an ounce of powdered nut galls, and let stand for a day or two, shaking it up occasionally; then let settle, then pour off the clear, then put a pint of water to the residue. When cold and settled, pour off the clear and mix with the first. Now to this add a grain of nitrate of silver, or sulphate of copper or nitrate of copper. Dissolve whichever you use in a little hot water before mixing with the other liquid. Silver is the best process. Clean all oil off and rust or scabs, paint, etc. Clean all up with bright pumice stone powder. Don't use emery in any form, but the above with a piece of wood. Then clean all off; dry with air slaked lime. Now go over it with the liquid with some cotton wool. If you have saved your powdered galls, take a little of that upon your wool, and you will find that a great acquisition in the first application. Let stand until dry, then give it another coat. When dry, scratch, brush it, and give it another coat, etc. When you have got it to your liking give it some linseed oil and camphor. All bright iron parts can be made like ebony polished, and with the gun-metal mounting you will have a picture in black and gold. Cylinder covers, etc., can be done the same; but you must wash with hot water before oiling it. It will stand any amount of heat, the hammer and friction in wiping; you have no blistering, and you will have some difficulty in eradicating it. Bicycles, repairs, handle bars, etc., can be treated the same way to advantage, well washed with hot water; when dry give them a coat of good copal carriage varnish."

Cement for Porcelain.—1. To cement porcelain to metal, make a mixture of equal parts of water and alcohol (95 per cent. strength), and use this fluid to make a paste with ten ounces finely powdered chalk and eight ounces starch. Then mix in three ounces of Venice turpentine. 2. An excellent cement for china and porcelain may be obtained by melting together seventy-five grains of fish-glue and five drachms of glacial acetic acid, and afterward heating the solution until it becomes of a sirupy consistence, so as to form a jelly upon cooling. To use it, the jelly is placed upon a stove, in order to bring it to a liquid state, after which the edges of the broken crockery are coated with it, and the pieces strongly compressed. 3. A cement for fixing enamel plaques to nickel is prepared by dissolving together by a gentle heat:

Gum damar..... 10 parts.
Copal resin..... 10 "
Venice turpentine..... 11 "
Oxide of zinc..... 3 "
Ultramarine.... quantities to tint the mass.

The coloring matter (zinc white and ultramarine) is stirred into the compound when the solids have been rendered fluid. This cement should be used hot, and when cold can be polished. It is also suitable as a putty for filling up cracks in enameled surfaces.—Pottery Gazette.

Glycerin Suppositories.

Lanolin..... 30 grains.
Glycerin..... 30 "
Cacao butter..... 30 "
White wax..... 15 "

Melt the ingredients together, stirring well, and pour into chilled moulds. The suppositories should weigh 90 rains each.—Re istered Pharmacist.