dissect most readily in a given place. All the blades were designed gradually as successive operations showed the need of peculiar shapes. In the average case not more than perhaps eight different blades will be used, but the odd ones will fill a need at some time. The shank and handle of the knives should be in a straight line. If they form a so-called nasal angle the sense of direction of the hand is interfered with and its motions are made complicated. The little cartilage knives are very important as they enable the operator to remove the entire cartilaginous deflection in one piece in most cases, thus saving a great deal of time and reducing the time of operation in cartilaginous deflections often to but thirty minutes. For forceps I still prefer the Grünwald punch. It reaches to the back of the nares and is not so large that it interferes with exact vision. A forceps that does not reach to the posterior border of the vomer in a large man is too short for use. The use of the chisel I confine to the base of the septum since Dr. Christian R. Holmes, at the Saratoga session in 1902, spoke warningly of its employment on the higher parts of the septum on account of the danger of shattering the delicate bone and creating far-reaching fissures. I operate now on all cases with the patient recumbent on a high operating chair. This avoids the faintness that often overcomes him when sitting, puts him at ease and the operator has more freedom of motion when he stands beside his patient than when he is sitting. To trim off remains of the deviation along the base of the septum I have the patient sit in a chair for a few minutes at the end of the operation. As a substitute for the expensive Kirstein light a 50-candle power, pigtail filament, stereopticon, incandescent, Edison light, attached to a stand, may be used for reflection with the head mirror.

DR. E. E. FOSTER, New Bedford, Mass.—The resection of the nasal septum mentioned by Dr. Gibb has not received in our country the proper amount of consideration and trial that it deserves. The Germans have for a long time resected the septum with much better results than we obtain from the Asch, the Gleason or any of the modifications of these or similar methods. Freer and White have been resecting the septum for the past two or three years and report good results. Their mucous incisions, however, are made in such a way that retraction, curling, perforations and granulations are more apt to occur than in the more simple operation of Killian, Jansen, Menzel, et al. Their operations are also complicated with many specially devised instruments. The submucous resection of the septum as done by Professor Killian consists of one straight incision through the mucous membrane over the cartilaginous part of the septum, about three-quarters of an inch long and nearly at right angles with the floor of the nose. The perichondrium and the periosteum of this side of the septum are elevated from that part of the septum to be removed; then an incision is made through the cartilage, in line with the first mucous incision, to the perichondrium of the opposite side. Through this last incision the opposite mucous surface can easily be elevated from the septum. The two mucous surfaces thus elevated are separated with a long Killian speculum, leaving the cartilage between the blades. The septum can now be removed with scissors, Hartmann's concho'one or Jansen-Middleton's forceps. In a few cases a chisel may be used to remove the lower part of the vomer, but should not be used until the overlying bone has been removed. The mucous incision can easily be closed with one or more sutures, but as a usual thing they are not required. The inferior meati are loosely packed with cotton, to be removed on the following day. I saw a large number of patients in Killian's clinic on whom this operation had been performed. All had a straight, smooth septum. In many of the older cases the septum was firm, there being a re-formation between the two mucous layers. I did not see a deformity of the nose as a result of the operation. The Killian long speculum and the elevator are the only instruments required that are not contained in a laryngologist's armamentarium.

DR. H. H. BRIGGS, Asheville, N. C.—I have had experience with 11 cases of resection operation. The first two were done with the instruments used in ordinary nasal work, and

were successful in restoring the function of the nostril except that perforations resulted, one 3 mm. by 6 mm. and the other nearly twice as large. The remaining cases were done with the Freer instruments devised especially for the resection operation, and the results have been very gratifying. I have had some difficulty in maintaining the flaps in place, but have overcome this in the last few cases by inserting a suture through the corners of the flaps and through the mucous membrane immediately below, after which the packing can be done without fear that the flaps may be misplaced. I consider the resection operation the preferable one in the great majority of deflections. It is tedious and requires much time and patience, but you are amply repaid for this in the end.

DR. KASPAR FISCHER, San Francisco—In all intranasal operations in which splints are not necessary we have to decide whether we should pack the nose or not. If we do not pack the nose we are liable to be called out at any time of the night on account of a secondary hemorrhage; if we do pack, the patient may pass a sleepless night from the discomfort of the packing. I would like to draw your attention to my method of using collodium as a dressing after nose operations, published in the *Archives of Otolaryngology*, 1902.

DR. JOSEPH S. GIBB—I am glad that Dr. Myles and Dr. Mayer are, in the main, in accord with my views as to the indications for both the Asch operation and the Watson-Gleason. I fully coincide with Dr. Myles in that the Watson-Gleason does not meet every indication in bony deviations, and that it is necessary, in some cases, to do supplementary operative work. Dr. Mayer has given me too high a number of good results in the Asch when he states there were 47 cures in 48 cases. The figures were 47 cases in which the result was good, 11 in which the result was fair, and 3 failures. I have not followed the practice of Dr. Mayer to use general anesthesia in the Asch operation. Nearly all the cases reported were done under local anesthesia. This, I presume, is a matter of custom. I have been accustomed, for years past, of employing only local anesthetics in all the various methods for the correction of deviated septa, so that when I began to do the Asch, a much less formidable operation than many of its predecessors, I could see no reason to change my method. In the main the results have been good, and I never use general anesthesia in these operations unless requested by the patient. In reply to Drs. White, Anderson, Foster and Freer, who have spoken so favorably of the "window resection" method, I would say that the time limit did not permit me to reach this part of my subject, and that I have little to add to what is said in the body of the paper. While I have had little experience with this method, I still believe, as stated in the conclusions, that it will not meet every case. I am sure there are cases of marked cartilaginous and osseous deviations in narrow nares, complicated, as is so often the case, by largely hypertrophied turbinates, in which the mucous membrane is so "thinned out" over the deviated portion as to make anything like a clean dissection absolutely impossible.

TUBERCULAR LARYNGITIS; PROGNOSIS AND TREATMENT.*

THOMAS J. GALLAHER, A.M., M.D.
DENVER, COLO.

In considering the prognosis of tubercular laryngitis it is my desire to do so under three heads:

1. In regard to local lesions.
2. In regard to restoration of function.
3. In regard to life.

Irrespective of the character of the local lesions it is impossible to give an abstract prognosis concerning them.

* Read at the Fifty-fifth Annual Session of the American Medical Association, in the Section on Laryngology and Otolaryngology, and approved for publication by the Executive Committee: Drs. G. Hudson Makuen, George L. Richards and John F. Barnhill.

Most careful consideration must be given the pulmonary and general condition of the patient. With extensive and rapidly breaking down pulmonary areas the reparative process in the larynx may be *nil*, as it is in other parts of the body. With slight pulmonary involvement, general condition good, the prognosis is more favorable. Laryngeal manifestations seldom improve under any treatment unless the pulmonary and general conditions improve.

With increased activity of the pulmonary disease, further infection of the larynx is likely to occur through the lymphatics and vessels or by direct infection of the bacilli in the sputum; or, again, the laryngeal disease progresses owing to the general resistance being diminished. We must consider the location of the lesion, area and depth of the infection, and also if many distinct places are involved. If the entire larynx is affected, including the perichondrium and cartilages, the case is hopeless. Ulcerations in the commissure and on the epiglottis are, in my experience, most intractable. Those involving the true and false vocal cords seem to yield more quickly. Most prompt and energetic treatment is necessary to limit destruction of the vocal cords in order to save the voice. There is a direct relation between extent and depth of ulceration and cure. Deep infiltrations are followed by the most extensive ulcerations. The more superficial ulcerations are more amenable to treatment.

LESIONS IN LARYNX.

The lesions found in the larynx may be placed under the following heads:

1. Infiltration: Superficial and deep.
2. Ulceration: Superficial and deep.

Both conditions are associated with more or less edema, which is usually found in the parts with underlying loose connective tissue, as in the epiglottis and aryepiglottic folds.

Infiltrations, both superficial and deep, may remain quiescent indefinitely, depending on the blood supply and the change in the tubercles. The tubercles undergo the same change as in other parts of the body, viz.: *a*, Softening; *b*, fibroid limitation (encapsulation); *c*, calcification. Ulceration follows softening; this in turn may be followed by cicatrization as a result of treatment. I have never seen a case of spontaneous cure of tubercular ulceration in the larynx. If fibroid change surround the tubercle, it may remain quiescent, but it is not destroyed. The term cure can only be applied to given laryngeal lesions which are replaced by scar tissue or to softened areas which have become impregnated with lime salts.

Restoration of voice will depend on the amount of muscular involvement and on the integrity of the vocal cords and freedom of the articulations. Laryngeal tuberculosis seldom if ever produces death *per se*.

INFLUENCE OF CLIMATE.

It is of the utmost importance that the pulmonary and general conditions be improved as rapidly as possible. The removal of the patient to a proper climate is imperative. The arrest of the pulmonary disease will lessen the probability of further laryngeal infection.

While good results have sometimes been obtained by laryngologists despite unfavorable climatic conditions, yet the prognosis in regard to life, as well as a cure of the laryngeal lesions, is much more favorable in a suitable climate.

PROGNOSIS.

To say that all cases of laryngeal tuberculosis are fa-

tal is erroneous; I have seen many cases in one stage or another recover; some who were apparently doomed, in which extensive involvement of the larynx had occurred, including considerable destruction of the epiglottis. I have had two cases in the last year where from one-fourth to one-third of the epiglottis was destroyed, and yet the patients recovered.

Many have recovered where the true and false cords and commissure were affected, either alone or in combination.

In every case it is our duty to give these patients the benefit of our best efforts as early and as thoroughly as possible.

Too often patients present themselves with a hopeless condition of the lungs as well as of the larynx, yet we must not be discouraged in our efforts to save those who have a possible chance for life.

TREATMENT.

In case active surgical procedures are deemed unwise, the treatment consists of thoroughly cleansing the larynx, followed by soothing applications and sprays. I have seen cases of infiltration remain quiescent from one to five years, and still the parts are intact; therefore, we can not deem it best to endeavor to remove this infected area in all cases.

If surgical procedures are decided on, however, the infiltrated area should be very thoroughly removed, and the parts stimulated to promote cicatrization. When once ulceration has occurred, however, I have seen nothing but good follow a thorough curettement and the energetic application of strong germicidal remedies. If vegetations are engrafted on the ulcerated areas they should be removed before any other treatment is instituted, so that the remedies may be more effective. The treatment by electrolysis has not been widely used, nor has the influence of the x-rays or radium been sufficiently demonstrated to warrant their general use.

In the treatment of the ulcerations many remedies have been used, and I desire to again call attention to the value of formaldehyd. Since my report in regard to this remedy, read before this Section in Denver in 1898, I am still further convinced that it is a remedy of very great value. It is satisfactory as a germicide, tends to shrink the exuberant granulations and stimulates the healing process. It is probably more penetrating than any other remedy which we possess. The best results are to be obtained by the persistent use of formaldehyd. In addition to the office treatment, I direct my patients to spray the larynx several times daily with a formaldehyd solution (gtt. 1 to 3 to ounce). The local applications of formaldehyd are made with a stronger solution, 3 to 5 per cent. This I follow with insufflation of aristol and orthoform, and lastly, the intratracheal and laryngeal injection of oil containing menthol, oil of cinnamon, etc. Formaldehyd should be thoroughly rubbed in.

Too much stress can not be laid on the proper use of this remedy, as many have failed to get its beneficial results owing to its too infrequent use. In addition to its influence on the ulcerated areas, it is a splendid prophylactic. My experience with formaldehyd extends over a period of ten years, and I have seen the most gratifying results from its use.

DISCUSSION.

DR. W. FREUDENTHAL, New York City.—It does not accord with my experience of tubercular laryngitis that the local manifestations seldom improve unless the general condition gets better. In laryngeal tuberculosis we may expect anything

and everything, and I have seen cases improve very much while the process in the lungs was getting worse rapidly. When we succeed in putting these ulcerations at rest by means of a narcotic, Nature will do the rest, and they often heal up while the process in the lungs is getting worse. Another method of treatment I have pursued has been the use of radium intralaryngeally. I have introduced it into the larynx and left it there for ten minutes at first, and then for half an hour, daily. The result was very interesting. Some of you remember the time when Koch's tuberculin was first injected and the changes that were produced in the larynx. The changes observed after the use of radium were almost identically the same. This patient had a large larynx, with ulceration of both vocal cords, but after a few sittings infiltration set up, edema appeared, and after the tenth day I had to stop treatment for a day. These infiltrations grew so that she could speak with some voice, and she was delighted. While I did not accomplish a great deal, it was an interesting experiment.

DR. W. S. ANDERSON, Detroit—The treatment of laryngeal tuberculosis with soothing applications gives better results than with those that are irritating. I also wish to indorse the value of intratracheal injections. I find them very efficacious in relieving the symptoms, allaying cough and improving the patient's general condition. I always use olive oil as a base, and such drugs as guaiaccol, camphor, menthol, oil of thyme, oil of wintergreen and oil of eucalyptus, and have found them very helpful.

DR. W. E. CASSELBERRY, Chicago—I welcome any new remedy for tuberculosis of the larynx in the hope of improved results, but confess in advance to skepticism. I have treated many cases in different ways, and the more I see of it the more I feel that local remedies have but little influence on the ultimate outcome of the disease, although often of incalculable value to relieve pain and distress. A few cases recover and some run a slow course, with periods of distinct betterment, but the definite improvements which I have observed have followed increasing the general resistance by correct mode of life rather than local medication. Concerning the surgical treatment, I am in accord with Dr. Gallaher, that one should hesitate before breaking up a mere infiltration. I have seen the use of the curette followed by a rapid aggravation of the condition. If, however, there is already an active ulceration, which is circumscribed and accessible, so that one can make a thorough curettement, I have seen the best results follow, especially if supplemented by the application of lactic acid. Therefore, I do not wish to disparage surgical treatment, but to plead for a wise selection of cases on which the curette is to be used. I differ somewhat as regards tuberculosis of the larynx going *pari passu* with the condition in the lungs. I have seen cases where death was practically dependent on the larynx, and I have also seen the reverse condition, where the larynx continued to improve while the lungs grew progressively worse. However, these cases may be exceptional. I would like to know how Dr. Gallaher manages to have the patient spray his own larynx. I have no apparatus by which a patient can spray his own larynx effectively with other than oily medicaments. In this connection, I would like to call attention to the spraying nozzle of Dr. Freer, which goes around the epiglottis and into the larynx, but it requires a physician to use it.

DR. KATE W. BALDWIN, Philadelphia—I want to mention the use of carbolic acid in these cases of tuberculosis of the larynx. After thorough cocaineization, the ulcer is touched with carbolic acid. I have seen a number of cases where the ulcer was absolutely cured in this way, especially where the treatment was instituted in connection with the x-rays.

DR. ROSS H. SKILLERN, Philadelphia—I agree with Dr. Freudenthal regarding the laryngeal and pulmonary lesions not being dependent on each other, so far as the general condition of the patient is concerned. I have often noted marked improvement in the local lesion, while the general systemic condition of the patient was growing steadily worse. The prognosis depends more on the general systemic condition than on the throat lesion and may be said to be favorable if the case is

seen at an early stage of its development and the tuberculous deposit in the larynx is localized. Extensive curettage should only be employed in those cases in which there is a fair chance of curing the disease. Regarding the preparation anesthesin, I have used it for some months, but fail to note any advantages over orthoform, although when the latter loses its effect anesthesin may be used as a very satisfactory substitute.

DR. OTTO T. FREER, Chicago—I think the galvanocautery should not pass unnoticed as a therapeutic measure in tubercular ulceration of the larynx. It produced complete recovery in one of my cases presenting epiglottic and interarytenoid superficial tubercular ulcers. The edematous infiltrations of tuberculosis are, of course, not suited for it. Recovery in any case of tubercular laryngitis, in my experience, is a rare event, and I was surprised at the rapidity with which thorough cauterization in this case arrested the disease. The good vitality of the patient and her moderate susceptibility to tuberculosis were aids in producing the recovery. If these conditions had not been present the treatment would doubtless merely have aggravated the disease.

DR. J. F. MCCONNELL, Las Cruces, N. M.—I have seen a great many of these cases which have come from high altitudes down to a low altitude in very bad condition. I have used a 2 per cent. solution of methylene blue in these cases with most gratifying results after all analgesics had failed. I apply it in the ordinary manner with the McKenzie forceps, in a 2 per cent. aqueous solution, taking care to get a good quality of methylene blue. The action is so prompt that in three or four hours all the annoying symptoms are relieved and the patient gets great benefit from this palliation. I know of no real effect on the course of the disease other than that due to climatic conditions, but it has a very decided palliative action, and is well worth trying.

DR. THOMAS J. GALLAHER—The method of spraying the larynx is not learned in a day, but these patients make repeated efforts and eventually yield to instruction. It is done with difficulty in a number of cases, but it does a great deal of good. It is simply a question of trial, and I am satisfied I can teach four out of ten to use the spray. After an experience of fifteen years with a large number of cases of laryngeal tuberculosis, I make the statement with a good deal of thought that the condition in the larynx does not often improve unless that of the lung improves also; but the converse of that I have not found to be true. While the larynx may be retrograding, the lung is often improved, this probably being due to the climatic effects on the pulmonary areas. In many cases nothing but breaking down of the larynx can result, though the lung may improve very rapidly. The curette must be used carefully, and we should always hesitate about breaking up a fibrous capsule. If it is once undertaken, it should be done with the idea of thorough removal, and it must not be attempted unless that can be done. After ulceration has begun, I do not know of any other treatment which compares with it. It is capable of producing much good, for the larynx is breaking down more and more; many of these cases will not improve without it. Methylene blue and intratracheal injections of various kinds all have their place. The influence of these on the cough is often quite marked and lasts for some time.

Black Penis from Dose of Antipyrin.—Malherbe of Nantes was consulted by a young man, recently infected with syphilis, on account of a large bluish-black patch which had suddenly appeared on the dorsum of the penis. He supposed that gangrene was imminent, but Malherbe reassured him on learning that he had taken 1.5 gm. of antipyrin six hours before to banish an attack of migraine. Aside from the discoloration the tissues were entirely normal, and there was no other discoloration elsewhere on the body. Fournier called attention more than five years ago to this possible action of antipyrin on the penis. Malherbe thinks that the phenomenon is due to an intense congestion entailing actual ecchymosis. The young man was in the habit of taking 1 to 2 gm. of antipyrin occasionally to cure migraine and had always found it answer the purpose, with no by-effects. The details are given in the *Gazette Méd. de Nantes* of June 18.