fection in an environment previously rendered insatiable.

The sanatorium in the crusade against tuberculosis has passed the experimental stage; the results achieved through persistent effort and honest endeavor on the part of the profession justify the co-operation of the laity through the medium of the state legislature.

Establish state institutions for the care especially of the tuberculosis poor that they may be cured of their disease and returned to the commonwealth able-bodied citizens.

"The prolongation of life by the suppression of preventable disease is of much greater value to the state than the cost of the means employed."

In the light of comparatively recent brilliant achievements in serum therapy, it is reasonable for us to assume that a more specific treatment for tuberculosis may yet be derived. As clinicians, however, we should accept only those measures applicable to the treatment of tuberculosis that are rational, and should not expose the unfortunate victim of this disease to widespread experimentation in exploiting so-called cures.

The open-air treatment of this disease through the medium of the sanatorium has effectually demonstrated that tuberculosis is a curable disease in any climate, and it becomes our duty as physicians to give to humanity the knowledge and facts we have gleaned in this field of medical science, endeavor, if possible, to eliminate the ever prevailing superstition in the efficacy of falsely advertised nostrums as a cure for tuberculosis, refuting in some measure the claims of charlatans.

Could we finally attain this end in our crusade against tuberculosis, much of the sorrow would forever be banished from the millions of afflicted homes and the night of tears changed to the morn of hope.

39 Pine Grove Avenue.

[THIS SYMPOSIUM ON TUBERCULOSIS WILL BE CONTINUED NEXT WEEK.]

AEROTHERAPY AND SOLAR THERAPY IN THE HOME TREATMENT OF TUBERCULOSIS.

WITH DESCRIPTION OF A WINDOW TENT.*

S. A. KNOPF, M.D.

Associate Director of the Clinic for Pulmonary Diseases of the Health Department; Visiting Physician to the Riverside Sanatorium for Consumptives of the City of New York.

NEW YORK CITY.

As the most important therapeutic factor in the modern management of tuberculosis, we must consider fresh and pure air, and we can make the home treatment most successful when we imitate, so far as practicable, all the salient features of the sanatorium treatment. In these institutions the patients live outdoors virtually for twenty-four hours of the day. Let us now see how we can imitate this aerotherapy of the sanatorium treatment in the home of the patient.

In summer, of course, all the windows are open, and during the day we place our patients in the yard, on the veranda or on the roof, whenever and wherever conditions permit us to have the patient take what is known as the rest cure in the open air. Here he rests on a reclining chair with a proper knee bend and comfort-

able back. If the patient can afford it we get him also a half tent. This half tent is composed of a frame of steel tubing, which can be folded together when not in use (Fig. 1). Over this frame strong sail duck is stretched and secured by snap buttons on the inside, so as completely to protect the patient against wind and sun. To prevent the tent from being overturned by the wind, the frame has ground spikes holding it securely. A beach wicker-work chair can also be made to do the service of the half tent. After the seat has been removed the inner walls of the wicker chair are lined with padding. A reclining chair is placed with its back in the interior, and the whole so arranged that the patient is protected from the wind and sun. Whenever the patient is on the chair he should be so comfortable as to allow complete muscular relaxation. Mind and body must be at rest.

For poor patients the simple steamer chair and a few boards joined together to replace the costly half tent will have to answer the purpose. A large and stout umbrella, such as is often used at seaside resorts, can be fastened to the back of the steamer chair, and will answer the purpose when the tent can not be pro-

Fig. 1.—Half tent with patient resting on metal reclining chair taking the rest cure.

vided. The poor patient in cities will probably be obliged to resort to the roof for his rest cure, as the small yard of the tenement house, with many children playing in it, will scarcely be suitable. I do not favor the use of the fire-escape for this purpose. A recent conflagration in New York City, where many lives were lost owing to the obstruction of the fire escape, showed the dangerous results of evading the law in this way. On his reclining chair in the open air the patient should remain during the day whenever he is not taking any walking exercises.

In the modern American, and also in some European sanatoria, the majority of patients have their beds moved out on the veranda during the night, and there they sleep, often in the coldest weather. The brilliant results obtained through this method of sleeping outdoors in cold weather are too well known to all American physicians to need further mention here. What has worried me for years has been the fact that we could provide this open-air treatment at night only for

* This is the chapter on Aerothreapy and Solar Therapy of an address on the Home Treatment of Tuberculosis delivered by invitation before the sixth Annual Conference of Sanitary Officers of the State of New York at Syracuse, Oct. 29, 1906.

so relatively small a number of patients when there are so many who need it, particularly among the consumptive poor. I believe that I have been fortunate enough to solve this problem in a measure by an invention which I call a "window tent." The first model of this device, which I presented to the profession, and in the construction of which I had the kind coöperation of the former resident physician at Riverside, Dr. McLaughlin, had many defects. It was nevertheless put in practical use by a number of physicians and by myself. A second model was somewhat better, but far

Fig. 2.—Window tent in position, with patient in bed and looking through the celluloid window into the room but breathing outdoor air only.

Fig. 3.—Window tent, raised when not in use.

from being perfect. In the third model I have tried to overcome all the defects learned through my own experience and that of others in the use of the tent. The Kny-Scheerer Co. have incorporated all my suggestions in their latest device, and on my request have refrained from seeking a patent. I am particularly grateful for this generosity on their part, for the tent will be of greatest service to the consumptive poor unable to avail themselves of sanatorium treatment. Many of them will

not even be able to pay the modest price at which the window tent is now put on the market, but with the illustrations before them they may have sufficient mechanical skill to imitate the device at very little cost.

The advantages of the last model of the window tent over the former models are greater simplicity and cheapness in construction, greater facility for cleaning and disinfecting and a method of attachment which permits the removal of the tent from the window when not in use.

As the illustration shows, this window tent is an awning, which, instead of being placed outside of the window, is attached on the inside of the room. It is so constructed that the air from the room can not enter or mix with the air in the tent. The patient lying on the bed, which is placed parallel with the window, has his head and shoulders resting in the tent (Figs. 2, 3, 4, 5). By following the description closely it will be seen that the ventilation is as nearly perfect as can

Fig. 4.—View of the window tent with patient, taken from the outside.

be produced with so cheap a device. The tent is placed in the lower half of the window, but it does not quite fill the lower half of the frame, a space of about three inches being left for the escape of the warm air in the room. By lowering the window, this space can be reduced to one inch or less, according to need, and on extremely cold and windy nights there need be left no open space at all above the window frame. The patient's breath will rise to the top of the tent, the form of which aids in the ventilation.

The tent is constructed of a series of four frames, made of Bessemer rod suitably formed and furnished with hinged terminals, the hinges operating on a stout pin at each end with suitable circular washers interposed to insure independent and easy action in folding, the Bessemer rod being hardened to make a stiff, rigid frame to insure maintaining the original form. The frame is covered with 7-ounce extra thick yacht sail

twill, properly fitted, and having elongated ends to admit of their being tucked in under and around the bedding to prevent the cold air from entering the room.

The patient enters the bed and then the tent is lowered over him; or with the aid of a cord and a little pulley attached to the upper portion of the window he can manipulate the lowering and raising of the tent himself. Shutters or Venetian blinds, whether they are attached on the inside or on the outside of the window, can be utilized in conjunction with the window tent as a screen to intercept the gazes of the neighbors and in stormy weather as a protection. The bed can be placed by the window to suit the patient's preference for sleeping on his right or left side, so that he has the air most of the time in his face. Another advantage of the window tent is that it will not attract attention from the outside. The bed being placed alongside the window will be convenient for a majority of the poor who have small rooms; if, however, the bed must be placed at a right angle to the window, it can be so arranged.

A piece of transparent celluloid is placed in the middle portion of the tent to serve as an observation window that the nurse or members of the family may watch the patient if necessary. It also serves to make the patient feel less outdoors and more in contact with his family. He can, if he desires, see what is going on in the room. If the window tent must be placed at a right angle to the window, the observation glass can be put in on the side. It goes without saying, that, as a rule, patients should not smoke; when, in exceptional cases, this may be allowed, the danger of the celluloid window becoming ignited must be impressed on them, and the greatest caution urged. I prefer celluloid to ordinary glass for this purpose, because it can easily assume the vaulted form of the rest of the tent, and thus avoid even the slightest possibility of an air-pocket formation.

If it is necessary to raise the bed to the height of the window sill it can be done with little expense. If the bed is of iron, a few additional inches of iron piping can be attached to the legs by a plumber or by any one handy with tools; a wooden bed can be raised with equal facility. If the window tent is to serve the patient only during the night it can be pulled up and the bed moved away from the window during the day and the window closed; or the tent can be taken from the hooks and put out of the way.

The window tent, of course, will be of greatest service to the consumptive sufferer in winter. If he is feverish, or his stay in bed is advisable, he can spend his entire time in the window tent. If the people are poor, and the room in which the consumptive sufferer lies serves as a living room for the rest of the family, the fact that the well members need not shiver and yet the patient can take his open-air treatment, is of vital importance in many respects. While the room will not be quite so warm as if the window were entirely closed, it will be much warmer than if there were no tent in front of the open window. Laying aside the economic advantages to a poor family in not being obliged to heat more than one room, the patient feels that he does not deprive his loved ones of comfort and warmth and that he is less a burden and hindrance to their happiness. The other members of the family, on the other hand, feel that they can give the patient all the air he needs and that he himself need not suffer for their comfort.

In winter the patient's bed must be covered with a sufficient number of blankets to assure his absolute comfort and warmth throughout the night. Still, this covering should not be so heavy as to press on the body and make the patient feel uncomfortable or tired. The tightly woven blanket is a better protection than the loosely woven one. To the poor, whose supply of blankets is, alas, often very limited, it may be valuable advice to tell them to put several layers of newspapers between the coverings. *Outdoor Life* for December, 1905, recommends having a dozen layers sewed between two layers of flannel. This will certainly make a cheap, light and warm covering. In extremely cold weather the patient, while sleeping in the window tent, should wear a sweater and protect his head and ears with a woolen cap, shawl or helmet, such as shown in Figure 6.

Some patients will complain that the bright light awakens them too early in the morning, and that they have difficulty in going to sleep again. In such instances I counsel the patient to have some light weight, but dark-colored material (such as black lisle thread hose) to put over his eyes. This usually obviates the inconvenience caused by the light.

The pulmonary invalid when in bed should be provided with a bell to communicate with his nurse, relatives or friends who take care of him. He should, of course, have a small sputum cup or pocket flask handy to receive his expectoration. I prefer the flask for use in the window tent, for it seems to me that any kind of a cuspidor which had to stand on the window sill would not be safe, as there is always a danger of its falling. A urinal should also be placed at the bedside, so that the patient will not have to leave the bed in the night.

There is one more benefit to be derived from the use of the window tent that is not to be underestimated. Patients who can only be persuaded with difficulty to sleep with the window wide open will not hesitate when they have this tent as an inducement. Draft, which the consumptive patient usually dreads, particularly when he perspires or in cold weather, need not be feared when sleeping in the window tent. The construction is such that even should the shoulders be accidentally exposed the three tent-walls protect the patient from violent currents of air which may be produced by leaving opposite doors in the room open. In this respect the window tent even has an advantage over sleeping on porches when they are not properly enclosed.

There is another advantage in this window-tent arrangement. The prolonged rest cure in bed will be more endurable when the patient is permitted to look out on the street and to watch life there, than when obliged to gaze at the four walls of his room.

When arranging for the rest cure on the reclining chair during the day, whether it is in the half tent in the garden, on the veranda, in the sleeping shack, on the roof, or on the balcony, one should always bear in mind that it is much more agreeable and conducive to the well-being of the patient, when taking the cure, to have a pleasant view to look on. In building a sanatorium, the greatest attention is paid to the proper selection of the place for the rest-cure gallery or veranda. The more pleasing and entertaining the outlook from these places, the more certain is one to keep the patients quiet and restful.

Lastly, an important point gained by the use of the window tent for consumptives is in regard to drop in-

fection, that is to say, the prevention of the dissemination of the bacilli through particles of saliva expelled during the so-called dry cough, sneezing, etc. While as a matter of course, the patient should be taught to always hold his hand or handkerchief before his mouth, when he coughs or sneezes, this is not always done, and to limit the possible infection to the interior of the window tent is obviously a great advantage. First, the constant exposure of the bacilli, which may have been expelled, to air and light will soon make them innocuous; and second, in this last model of the window tent the canvas of the tent is attached to the frame by simple bands so that its thorough cleansing, washing or disinfection is made easy.

Northrup and others have recently demonstrated that the ideal treatment of pneumonia is likewise pure, fresh and cold air by day and by night. When sleeping on the roof or in the open air is not feasible, the window tent is admirably adapted to the treatment of this disease in adults and in children. Thus, even if an intercurrent pneumonia should develop in the tuberculous patient, the window tent would be still of great service.

When there is no garden, veranda nor roof, the window tent can also be brought into service for the rest

Fig. 5.—Diagram showing ventilation of window tent.

Fig. 6.—Woolen helmet to be worn when sleeping outdoors in cold weather.

cure during the day. The bed is moved away and the reclining chair is put in its place. The latter can be raised to the necessary height by wooden blocks or a platform and with the aid of blankets and comforters the air from the room can be excluded, and the patient being in front of the open window breathes only outdoor air. When beginning this aerotherapy, it is, of course, essential that it be done gradually according to the susceptibility to cold of the patient. Impress on him the fact that night air is as pure as day air, and begin by placing him in the tent for a few hours at night and a few hours during the day in the chair, gradually accustoming him to living in the pure air day and night. A hot-water bottle for the feet, either in bed or in the chair, may often be necessary in extreme cold weather, for the patient's feet must be kept warm if he is to be benefited by the open-air treatment.

As an adjuvant in aerotherapy I must not forget to

call attention to the importance of respiratory exercises. I have repeatedly described and illustrated them in my books¹ and various articles,² so that I do not think it necessary to dwell on the details of this phase of aerotherapy at this time. I will simply say that these exercises judiciously taught and carefully carried out are suitable in nearly all afebrile, early and moderately advanced cases. Even in the third stage I have seen beneficial results from judiciously directed breathing exercises. Most careful clinical observation made under Dr. Otis,³ at the Massachusetts State Sanatorium, prove conclusively the value of careful breathing exercises in the treatment of pulmonary tuberculosis. The pneumatic cabinet must receive mention as a second adjuvant in aerotherapy of pulmonary tuberculosis. Those interested in this subject can refer to the works of Bowditch,⁴ Quimby⁵ and others. I have also described its indication and contraindication in my text-book on Pulmonary Tuberculosis.

After aerotherapy comes solar therapy or heliotherapy, that is to say, the utilization of the rays of the sun in the treatment of pulmonary tuberculosis. To be as brief as possible, let me give the concise directions which I think should be followed in order to get the best possible results from the sun bath:

The sunniest room should be selected for the purpose. Fixed carpets, of course, should not be placed in such a room, and the floor must be kept scrupulously clean. In a private home, where neighboring windows are often near, the arrangement will be more difficult, and low screens will have to be used. In winter the room should be heated to from 70 to 75 F.; as the patient's skin becomes less sensitive, the temperature of the room can be decreased. The room must always be well ventilated, and in summer the upper part of the windows can be left open.

The patient undresses entirely, but if he complains of cold feet he can keep on his stockings and even his shoes until he has become warm enough and desires to take them off. He first places a warmed sheet around his body and then a large blanket; he then lies down on the floor in the sun, his head in the shade and slightly elevated by a cushion. As he begins to feel the warmth of the rays of the sun, he gradually uncovers himself until the whole of his body is exposed to the sun; he exposes his back by turning on his chest. He remains in the sunroom for from half an hour to two hours, according to the instructions given him by his physician. He may change the recumbent to the sitting position or walk about.

Like all curative agents in the treatment of phthisis, the carrying out of this solar therapy should never be left to the caprice of the patient. Too much exposure to the hot rays may cause an erythema, an urticaria or other cutaneous trouble. To prevent these the patient should cover himself with one or even two layers of his sheet whenever the sun's rays produce a slightly burning sensation. Should these cutaneous complications occur nevertheless, the sun baths must be omitted for a

1. Les Sanatoria, thesis, Paris, 2 Ed., 1900. Pulmonary Tuberculosis, Prophylaxis and Treatment, Blakiston, Phila. Tuberculosis as a Disease of the Masses and How to Combat It. Am. Ed., F. Flori, New York City.
2. Johns Hopkins Med. Bul., September, 1901. Ann. of Gyn. and Ped., 1903, xvi. Jour. of Outdoor Life, July, 1905.
3. Otis: Trans. Nat. Assn. for Study and Prevention of Tuberculosis, 1906.
4. Bowditch: Boston Med. and Surg. Jour., July 16, 1885; and THE JOURNAL A. M. A., Aug. 1, 1885.
5. "The Pneumatic Cabinet in the Treatment of Pulmonary Phthisis," Internat. Med. Magazine, January, 1893.

time and the skin bathed in warm water, and friction with lemon juice applied. Headache, or a feeling of discomfort, is the signal to stop, no matter how short a time the bath has lasted. A high temperature is, of course, a contraindication to sun baths; such sensitive patients must remain in bed. Slightly feverish patients may be allowed to try solar therapy, but when experience shows that the baths are followed by an elevation of temperature, they must be discontinued. While taking his sun bath, the patient should perform frequent abdominal and thoracic breathing exercises.

If it is not possible to take sun baths, or enough sun baths, undressed at home, patients should be advised to take them outdoors, and should be told to dress as far as possible in light-colored clothes—never in black, red or brown—so as to permit the better penetration of the actinic rays. Patients should always take an umbrella or parasol with them to be able to shade their heads.

16 West Ninety-fifth Street.

RESUME OF THE COMPLETE GASTRIC ANALYSES IN SIXTY-ONE CONSECUTIVE CASES OF INDIGESTION.*

ALBERT WOLDERT, M.D.
TYLER, TEXAS.

I have previously reported 30 consecutive cases of indigestion which I treated in private practice, and in which complete gastric analyses were made.¹ I now supplement that report by adding to the list 31 new cases, thus making a total of 61 cases reported to this date.

In my treatment of stomach disorders I have to a large degree followed the methods outlined by Ewald, as follows: 1, To secure a complete accurate history of each patient; 2, to determine the percentage of free hydrochloric acid; 3, to determine the percentage of total acids; 4, to determine the motor power of stomach; 5, to determine the absorptive power of stomach; 6, to determine the digestive power of gastric juice on albumin (meats); 7, to determine the digestive power of gastric juice on starch (breads); 8, to determine the presence or absence of the milk-curdling ferment (rennin); 9, to determine the presence or absence of bile in the gastric contents; 10, to determine the size and position of the stomach.

In order to determine the percentage of free hydrochloric acid, I have used the decinormal solution of sodium hydrate, which I always prepare myself by standardizing this decinormal solution of soda with a decinormal solution of oxalic acid, the crystals of which acid have been purified by boiling, recrystallizing and washing with ether. Several titrations are often necessary in standardizing the decinormal solution of sodium hydrate, for on its accuracy depends the most important data to be obtained by the stomach examination, namely, the determination of the percentage of the free hydrochloric acid in the gastric contents.

The indicator used for the free hydrochloric acid was the Töpfer reagent in nearly every instance, being confirmed by the tropeolin test; for the total acids I used the phenolphthalein test; for the digestive power of albumin the white of egg; for the digestive power of

starch the Lugol solution; for the motor power in perhaps ten instances the salol test was used, but in the others lavage three and one-half to four hours after a small meal; for the absorptive power the potassium iodid test; for the milk-curdling ferment (rennin) about 2 drams of milk to three or four drops of gastric juice, and placed in an incubator; and for the presence of bile the nitrous-acid test. The size and position of the stomach was determined by inflating the stomach with air, supplemented by gentle percussion.

To complete the examination, I have in nearly every instance made a careful examination of the heart, lungs, liver, spleen, urine, and when the occasion demanded it tested the eyesight for defective vision, examined the blood for the malarial parasite and made a microscopic examination of the gastric contents.

I have made it an almost invariable rule not to prescribe for a patient suffering with gastric trouble on his first call to see me, but require that he shall wait until the complete gastric analysis has been made before giving medicine. Since I have been engaged in this work I have been much impressed with the importance of making these examinations, and often have brought before me cases showing how futile all efforts would have been if I had tried to guess at the real condition present as suggested by the different symptoms presented.

SUMMARY OF THE SIXTY-ONE CASES OF INDIGESTION.

Entire number of cases in the series and treated in private practice was 61.

Variation in Percentage of Free Hydrochloric Acid.

Free HCl was normal (.040 to .073 per cent.) in 10 cases or 16.66 per cent.

Free HCl was diminished (below .040 per cent.) in 7 cases or 11 per cent.

Free HCl was increased (above .073 per cent.) in 35 cases or 57 per cent.

Free HCl was totally absent in 9 cases or 14 per cent.

Percentage of Free Hydrochloric Acid.

Free HCl above .073 per cent. in 35 out of 61 cases or 57 per cent.

Free HCl above .10 per cent. in 22 out of 61 cases or 36 per cent.

Free HCl above .15 per cent. in 10 out of 61 cases or 16.66 per cent.

Free HCl above .20 per cent. in 2 out of 61 cases or .3 per cent.

Percentage of Total Acids.

Total acids were normal (35 to 60 per cent.) in 37 cases or 61 per cent.

Total acids below 35 per cent. in 14 cases or 23 per cent.

Total acids above 60 per cent in 8 cases or 13 per cent.

Total acids above 88 per cent. were none (highest percentage was 88 per cent.).

Total acids below 4 per cent., none (lowest percentage was 4 per cent.).

Motor Power of Stomach.

Motor power of stomach was deficient in 26 out of 29 cases examined.²

Absorptive Power of Stomach.

The absorptive power of stomach was found deficient (over 15 minutes) in 48 cases or 80 per cent.; normal in 9 cases and not determined in 3 cases.

Digestive Power of Gastric Juice for Albumin (Meats).

Of the series of 61 cases the digestive power of gastric juice for meats was good in every case in which the HCl was in excess or even deficient (except one case), and always totally absent when the HCl was totally absent. The digestive power for albumin was not determined in 2 cases.

* Read by invitation before the Central Texas Medical Association, Corsicana, Texas, July 11, 1906.

1. American Medicine, 1904, June 4.

2. In my opinion our rules employed to determine the motor power of stomach are defective.