SUMMARY OF RESULTS IN 1,000 CASES.

In other words, the condition at present of these 1,000 cases is as follows: Well, 28.7 per cent.; arrested, 6 per cent.; dead, 33.6 per cent.; untraced, 22 per cent. From this it is seen that 444 patients are alive, 336 dead and 220 untraced. Bearing in mind the fact that the patients of the first group were all discharged before the earliest cases of the second, it nevertheless appears that the patients of the second group have done better since discharge than those of the first group. Many of the untraced cases in both groups are undoubtedly dead. The reason why the percentages of the cases that have died do not vary more is probably due to the fact that in the second group a much larger percentage of the cases that have died have been traced. For this reason it is well to compare only the cases still alive in each group. Of the first group 10 per cent. of the cases are alive and 8 per cent. are well from eight to eighteen years after discharge. Of the second group, 44 per cent. of the cases are alive and 28.7 per cent. are well from two to eight years after discharge.

III.—RESULTS OBTAINED IN FIFTEEN HUNDRED CASES DISCHARGED FROM TWO TO EIGHTEEN YEARS AGO.

The number of cases discharged from the Adirondack Cottage Sanitarium until two years ago was 1,500. Of these 366 (24.4 per cent.) were incipient, 778 (51.8 per cent.) were advanced, 343 (22.8 per cent.) were far advanced, 2 cases were doubtful and 11 cases had incomplete records.

A.—Incipient Cases.—The 366 incipient cases were discharged as follows: Apparently cured, 245 (67 per cent.); arrested, 83; improved, 29; unimproved, 8. At the present time 170 (46 per cent.) are well; 13 are in a state of arrest; 14 have relapsed; 7 are chronic; 54 are dead; 108 are untraced. Of the 245 cases apparently cured 131 are well, while 30 of the 83 arrested cases, 9 of the 29 improved cases and none of the unimproved cases are well. Twenty-two of the apparently cured cases are dead; 16 of the arrested, 10 of the improved, 6 of the 8 unimproved cases are dead. The untraced cases are as follows: Apparently cured, 75; arrested, 23; improved, 8; unimproved, 2.

B.—Advanced Cases.—The 778 advanced cases were discharged as follows: Apparently cured, 90 (11.5 per cent.); arrested, 338; improved, 261; unimproved, 78; dead, 10. At the present time 151 (19 per cent.) are well; 51 arrested; 27 relapsed; 39 chronic; 295 (38 per cent.) dead; 215 untraced. Of the 90 apparently cured cases 54 are well, while 80 of the 338 arrested, 16 of the 261 improved and 1 of the 78 unimproved are well. Nine of the apparently cured cases are dead; 106 of the arrested; 118 of the improved, and 52 of the unimproved. The untraced cases are as follows: Apparently cured, 21; arrested, 71; improved, 100; unimproved, 22.

C.—Far Advanced.—The 343 far advanced cases were discharged as follows: Arrested, 51; improved, 116; unimproved, 152; dead, 24. At the present time 6 (1.7 per cent.) of these are well; 5 arrested; 3 relapsed; 9 chronic; 220 dead, and 100 untraced. The cases that are well at present are as follows: Arrested, 2; improved, 1; unimproved, 3. Twenty-eight of the arrested cases are dead; 63 of the improved, and 45 of the unimproved.

SUMMARY OF RESULTS IN FIFTEEN HUNDRED CASES.

In all, 497 (33 per cent.) cases are alive; 569 (38 per cent.) are dead, and 434 (29 per cent.) are untraced. Of the 497 patients that are alive 329 (22 per cent.)

are well; 69 (4.6 per cent.) are arrested; 44 (3 per cent.) relapsed; 55 (3.7 per cent.) are chronic.

If we deduct 434 untraced cases we have left 1,066 cases which have been traced. Of these 31 per cent. are well; 6.5 per cent. are arrested; 4 per cent. have relapsed; 5.2 are chronic; 53.3 per cent. are dead. Deducting from the number of incipient cases those that have been untraced we find that there are left 258 cases. Of these 66 per cent. are well and 21 per cent. are dead. Of the 563 advanced cases which have been traced 28.6 per cent. are well and 52 per cent. are dead; while of the 235 far advanced cases which have been traced 2.5 per cent. are well and 90 per cent. are dead. It must be remembered, however, that these figures are only approximately correct, as no doubt more of the untraced cases are dead than these figures would indicate.†

Clinical Reports.

A CASE OF NEURITIS OF THE RIGHT INFERIOR DENTAL NERVE

INCIDENT ON FORWARD DISLOCATION OF THE MANDIBLE OF THREE YEARS' STANDING.*

GEORGE A. MOLEEN, PH.G., M.D.

DENVER.

Aside from its rarity, the following case will illustrate the importance of a clear history and complete examination even in such a simple malady as facial neuralgia. I acknowledge my indebtedness to Drs. Packard and Silverstein for their prompt care and treatment of what seemed to be the cause.

Mrs. L. K., aged 49, born in Lebanon, Pa., has been in Colorado three years and did not come for her health; occupation housewife. First seen Feb. 7, 1903.

Family History.—Parents born in Pennsylvania; mother died of la grippe, father from dropsy. There were fourteen in the family, of whom five sisters and two brothers are living, the remainder are dead from various irrelevant causes. Tuberculosis, other chronic diseases, insanity and nervous diseases are denied.

Past History.—Of the exanthemata she had diphtheria at 9; measles at 22; denies scarlet fever; has had no sequelæ nor has she at any time complained of aural discharge, dizziness or vomiting. Menstrual history good; climacteric six months ago. When young she complained considerably of toothache, and neuralgia was suggested. Shortly after her arrival in Colorado she began to have paroxysmal pain confined to the right side of face, which seemed to originate, according to her localization, about the right lower bicuspid. This was drawn without relief, and after several medical and dental examinations the lowers were drawn to no purpose, until finally, on the advice of a physician, all her teeth, upper and lower, were removed in April, 1902, without modifying her symptoms.

Present State.—She has slept little for three months past, on account of pain. She says sedatives have given but little relief. The paroxysms come on at varying intervals of from 3 to 10 or 15 minutes, and seem to be excited by movement of the jaw, as in eating or talking, and can be excited by forcibly depressing the jaw. They sometimes begin in the ear and at others in the alveolar process. During the paroxysm the face becomes pinched, the right side flushed, and in severe attacks perspires slightly on the right side only. The pupil of the right eye is seen to dilate when the eye is held open and her hands restrained. Between paroxysms she becomes quiet,

† Since the above was written 5 of the untraced cases have been found to be alive, making 502 cases alive. In 10 cases death was due to other causes than tuberculosis.

* Read before the Rocky Mountain Interstate Medical Association, Salt Lake City, 1903.

though she complains of prickling of the right side of the face and of numbness and tingling on the right side of tongue. Sometimes there is a roaring in the right ear and a burning sensation in the right nostril. She has been partially deaf for some time and can be made to hear with difficulty, though more readily in the right ear.

Examination.—Feb. 7 and 10, 1903. Well-developed German woman, with drawn, careworn, though anxious, expression. All voluntary and co-ordinate motion normal and equal. Reflexes normal and equal and there is no response on tapping the masseter. Tongue is protruded in the median line, without tremor.

Sensory phenomena are in the main normal throughout the body, though she has impaired tactile sensibility with hyperalgesia on right side of face, the outline of which is as follows: Beginning at the external angular process extending upward to about the bregma, then downward and apparently dividing the ear—i. e., the tragus is more hyperalgesic than the concha and pinna—continuing downward to about one-half inch below the angle of lower jaw and forward to about the median line. Touching the skin at this area is not perceived as readily as on the opposite, and when perceived is painful. Pressure over the mental foramen causes severe pain and sometimes excites a paroxysm, the same being true at the infra-orbital opening. The supra-orbital is seemingly no more sensitive on one side than on the other. The submaxillary gland is also very tender and excites an attack readily. The mucous membrane of the right nostril, as well as the same side of the tongue, was markedly hyperesthetic.

Special Senses.—Salt placed on the right half of the tongue in the usual manner was met by immediate signal, but the tongue was washed and wiped, avoiding the left side, without allowing her to withdraw it, and she was unable to recognize and name the substance, even after several attempts with sugar, vinegar, etc.; near the base of the tongue, however, i. e., at about the anterior pillar, she recognized readily. Taste was present throughout the entire left side of the tongue. When asked why she signaled, she stated that she felt it as rough and thought she could taste it, but it was the pain she perceived. There can be no evidence of interference with smell detected.

Diagnosis.—The foregoing was strongly indicative of some mischief along the two lower divisions of the fifth nerve, and especially the third, and that it was undoubtedly irritative in character, a cause was consequently sought.

It was noticed that on depression of the lower jaw the condyle became quite prominent on the left side of the face, about three-quarters of an inch in front of and slightly below the external auditory canal. The right side revealed a corresponding depression, the jaw then rotated laterally and with considerable force, and by examination it was found that when depressed the internal pterygoid muscle was in a state of tonic contraction.

It then seemed but fair to assume that the condyle of the right side was irritating the inferior maxillary nerve at or about the site of the otic ganglion, which assumption was cleared on a study of its communications and branches, viz.:

The second division was affected through the connection by means of the vidian and petrosal, with Meckel's ganglion.

The sympathetic connection afforded an explanation of the flushing, sweating and pupillary symptoms.

The branch of the tensor tympani explained the increased hearing on the affected side.

The chorda tympani accounted for the loss of taste in the anterior portion of the same side of tongue.

It will be seen that the area of the first division was exempt.

Treatment.—Just how long the displacement had existed, or when it occurred, could not be learned, and the patient was even ignorant of its existence.

On account of the probable filling of the glenoid cavity by fibrous tissue a brace was advised, which would constantly exert an upward and backward pressure on the chin and still allow movement of the jaw, also with spring compresses to avoid the lateral movement of the condyles. She was ac-

cordingly referred with the above suggestions. A brace was devised which met the indication, and iodids and small doses of mercury given for a time.

Course.—After 4 or 5 days the pain began to subside and in about 10 days disappeared entirely. The sense of taste returned rapidly and the hearing on the affected side diminished until equal to that of the other side. She continued wearing the brace until about June 22, when she developed pain in the tongue and was unable to protrude the organ. Whether it was due to irritation of the hypoglossal by the chin rest or not I can not say, though it disappeared in two or three days after removal of the brace.

About July 1, having removed the brace, she began to experience very severe paroxysms of pain, which she described as boring in character, and localized constantly at a point a little below the condyle of the same side. The attacks were therefore quite different from her former ones, there being no area of hyperalgesia, nor was the tenderness of the mental foramen as marked. There was no swelling or tenderness along the process at this time.

The possibility of neuroma at or about the point of irritation was considered, and Drs. Hopkins and Freeman were consulted regarding the advisability of operative interference,

The brace in position.

which now seemed to be the only means of affording her relief. During this time she was given three grains of codein in divided doses, in the course of three hours, which gave her more relief than three-eighth grain morphin. A small cyst was noticed at the angle of the jaw, internally, and opened without perceptibly influencing the symptoms.

The brace was reapplied and adjusted to increase the pressure on the chin, and at first had little effect. The potassium iodid and mercury (red iodid 1/50 grain) were resumed.

The pain gradually left her and at the present time she scarcely notices it. The brace has been kept in place steadily since, and she is still taking the iodids, mercury and codein, while enjoying her escape from the surgeon for the present.

DEATH BY LIGHTNING.

B. KELLY, M.D.

CLIO, IOWA.

A powerful man, weighing 212 pounds, 36 years old, was struck by lightning. In about 10 or 15 minutes I found him deeply cyanosed, skin relaxed and cool and the heart beating feebly, but there was no visible respiration. I gave him hypodermic injections of strychnin, whisky and finally the normal salt solution, keeping up artificial respiration for ten hours, when the heart failed. The face brightened slightly for a few moments under the artificial respiration and the heart at short periods beat stronger, but no farther reaction was obtained, the temperature steadily falling.

Finding literature scarce on the subject I will be glad to have some one state what would have been the prospects for a better result by the use of an electric battery.