ALUMINA.

This term has been adopted in modern nomenclature to signify the earthy basis of alum or pure clay. This earth is one of the most extensively diffused in nature; its properties (at least the most important of them) have been familiar to mankind from the earliest antiquity, and on every account it deserves particular attention.

Alumina is nowhere found native in a state of purity, and it is a little remarkable that in all the natural clays, in which some of its characteristic properties are the most conspicuous, it never forms even half the composition, and generally no more than a third or fourth; and, on the other hand, it is found nearly uncombined in the hardest gems and the corundum, minerals in which its existence was for ages unobserved, till the experiments of the illustrious Bergman, and where it exhibits properties widely different from those of clay. The term alumina, therefore, is well chosen, both as it suggests the most important saline combination of this earth, and as it prevents those erroneous ideas of its properties which might attach to the ancient name of clay.

Alumina, when pure, is white, and without taste or smell. The natural clays, indeed, and the other earths that contain much alumina, have a peculiar earthy smell when breathed upon, which is a test of some use in mineralogy, but this appears to be owing to the combination of clay with the oxide of iron, and certainly does not belong to pure alumina. This earth is incapable of artificial crystallization, nor is it, like siliceous, found native in a state of absolute purity and crystallized. However, in one or two varieties of the corundum gems, alumina forms by far the greater part of their composition, and thus circumstanced, this earth is regularly crystallized and intensely hard. A peculiar difficulty lies in the way of ascertaining the true specific gravity of this earth; in the gems it is considerable, being from about 3.99 to 4.28.

Alumina, either as it exists in the natural clays or when prepared by art, till it has been exposed to a red heat, unites readily with water into a very tenacious, unctuous-feeling mass, which differs from all other wetted earths in being so plastic that it may be turned on a wheel, or moulded into any form. On adding more water it becomes gradually and equally diffused, making the liquor uniformly milky, and remaining very long suspended before it subsides. It is not, however, in any degree soluble in this fluid. Plastic clay retains water in greater quantity, and for a longer time than any other earth; on drying, if heated very gradually, it shrinks in every direction, and at last cracks into a number of pieces, and returns to its original state. When thus dried at a moderate heat, it is soft, greasy, brittle, and crumbly; when rubbed, it takes a good polish, and remains equally plastic as before, on the addition of fresh water. But if it is dried at a full red heat, it becomes extremely hard, so as to strike fire with steel, is harsh to the touch, and now no longer mixable with water into a plastic mass, nor does it absorb any sensible quantity of this fluid. Its original properties, however, are regained on solution in an acid, and precipitation by an alkali, or *vice versa*.

Alumina in many instances gives its characteristic properties with respect to union with water, to a very large proportion of any other earth with which it is mixed. In the natural clays, as we have mentioned, it never forms quite half the mixture. The very pure aluminous earth of Silmanitz, according to Klaproth, contains only 45 per cent. of alumina, and in the best porcelain clays the alumina is seldom more than 25 to 35.

The action of heat on moistened alumina is very striking. The first effect is to expel most of the large quantity of water which it has absorbed in becoming plastic. The greater portion of this fluid is expelled at a very low red heat, or at the point at which the earth assumes that entire alteration in its texture as to lose the power of again uniting with the water which it has lost. In consequence of this it shrinks in every direction. But the loss of weight and diminution of bulk do not stop here, for they both continue (the former in a diminishing ratio, the latter tolerably uniformly) up to as intense a heat as has ever been produced in furnaces. Hence it is that clay by its contraction affords a certain indication of increase of heat as all other bodies do by their expansion; not that this earth, however, is incapable of expansion by heat, for, like every other substance in nature, alumina dilates when heated and contracts when cold; but this is only observable when the heat applied is not more than the highest to which it has been exposed since its plastic state; above this point the contraction peculiar to this earth (produced either by actual loss of water or a different arrangement of its component particles) begins again to operate, and the extreme limit of its shrinking has never yet been precisely determined. A scientific manufacturer, Mr. Wedgwood, very ingeniously applied this property of clay to the construction of a thermometer for measuring the more intense degrees of heat, where the common instruments were inapplicable, but, though a valuable instrument, it is not susceptible of that degree of accuracy which has often been attributed to it.

M. Saussure observes that when alumina is precipitated from alum by ammonia or by carbonated ammonia, it is capable of existing in two forms, according to the manner in which the solution of alum was made; if no more water was used than necessary to dissolve this salt, the precipitated earth when dried at about 80° is a white, friable, light, very spongy earth, sticking to the tongue, which the author calls spongy alumina. But, on the other hand, if the alum is dissolved in a very large quantity of water, the precipitate dried at the same moderate warmth will be transparent, yellow, brittle, cracking in the heat of the hand, like rolls of sulphur, with a smooth conchoidal fracture, not appearing like an earth, nor adhering to the tongue, and not again mixable with water. This the author terms gelatinous alumina. At the above temperature of about 80°, both species contain an equal quantity of water, though the spongy occupies ten or twelve times the bulk of the gelatinous. But at a higher heat they differ much in the power of retaining water. The spongy alumina loses at a full red heat, rather less than is sufficient to melt silver, all the water which it had absorbed amounting to 58 per cent.; so that 100 parts dried at 80° are hereby reduced to 42, and lose no further weight by the highest intensity of heat that can be given. But the gelatinous alumina dried at 80° is said by M. Saussure to lose only 43 per cent. of its weight at a silver-melting heat, 48¼ at a heat of 130° of Wedgwood (or sufficient to melt cast iron), and appears never at any higher heat to alter further in its weight.

M. Saussure further remarks that contraction of clay (in the higher heats at least) cannot be exclusively owing to the loss of water, or to any loss sensible to the balance, since the diminution in weight from the thirteenth degree of Wedgwood to the highest intensity of heat is very trifling, whereas the contraction in the bulk of the clay, within these terms, is more than a fourth of its bulk.

Pure alumina is infusible in any heat that furnaces can procure, if not in contact with an alkali, or with any other earth which may operate as a flux. However, when exposed to oxygen gas from the blow-pipe, supported on a piece of charcoal, both Ehlman and Lavoisier found it converted into a milk-white globule, imperfectly fused, and intensely hard, so as to scratch glass nearly as the diamond. Some doubt, however, arises whether the alumina might not still contain a minute portion of potash, as it was prepared by precipitation from alum, the difficulty of procuring perfectly pure alumina in this method being very great. Besides, as all charcoal contains some alkali, this might have assisted in the fusion, so that it is still doubtful whether pure alumina is capable of melting without addition, by any known heat. The other earths promote the fusibility of alumina when mixed with it. Gerhard found this earth fusible in a chalk crucible, but not in one of clay; Klaproth found lime to melt in a clay crucible, and not in one of chalk; but it is certain that alumina is one of the most refractory of all bodies in nature, and hence its use in its natural clayey mixtures in forming crucibles, furnaces, and vessels for resisting fire, and containing other bodies to be exposed to heat.

Fused with alkalis, alumina does not run into a clear glass like the other earths, but forms an opaque, puffy, white semi-vitreous, pasty enamel. So Klaproth found the corundum, in which the alumina is about 84 per cent., fuse with its own weight of carbonated soda in an intense heat, into an opalescent, greenish-white, and glass.

Alumina is soluble in most of the acids, with more or less facility, and without effervescence, unless newly precipitated from a solution, and still retaining a portion of carbonated alkali, which it does with considerable force.

For a complete solution, the assistance of heat is generally required, unless it be effected by double decomposition, as where acetated alumina is formed by adding acetated lead to alum. This earth is much more soluble when in its plastic state than after it has been heated, and, most of all, immediately after precipitation from its solution either in an alkali or an acid.

Alumina dissolves with ease in the pure fixed alkalies, and even in ammonia if newly precipitated. If alumina in this fine state of division be put into three or four times its weight of a solution of potash, and heat applied, it dissolves like wax in oil, and by slow evaporation dries into rhomboidal pieces. Even the hardest aluminous stones, such as the corundum, yield to fusion with alkali with particular management. This solution lets fall the alumina when saturated with any acid, an excess of which redissolves the precipitate. Muriated ammonia also separates the earth from the alkaline solution by a complicated affinity, the potash uniting with the muriatic acid, and the ammonia thus expelled determining the precipitation of the alumina, which when properlyedulcorated and dried is now perfectly pure.

In many respects alumina resembles siliceous so strongly as to have been frequently mistaken for it; in hardness the aluminous gems surpass even that of rock crystal; both earths are soluble in fixed alkali, precipitable thence by an acid, resolvable in a slight excess of acid; and when in strong aggregation, the aluminous earth resists the solvent power of acids as completely as the silicious. The most conspicuous difference between them, as they occur in analysis, is the following: Siliceous when redissolved by excess of acid, after being first precipitated by it from its alkaline solution, is entirely again separated by the application of heat, and by evaporation nearly to dryness, and then the earth remains insoluble on any future application of acid. Alumina, on the other hand, though the solution has been evaporated to dryness (if not otherwise decomposed by heat), remains again soluble on affusing fresh acid, and thus may be completely separated from the siliceous.

From its strong affinity for magnesia, lime, barytes, strontian and other substances, as well as siliceous, it may be imagined how difficult it is to procure this earth absolutely in a state of purity.

Pure alumina may be prepared either by precipitation from alum, or from the finer clays or aluminous earths. Each method has some particular advantages, but in each some precautions must be taken to insure its efficiency.

In alum, the only earth is that which is intended to be obtained, which is a considerable advantage, and the affinity of the earth both for sulphuric acid and for potash, is at least as great as that which it shows for other earths, and accurate experiments have shown that a single precipitation from alum can scarcely ever be depended on to obtain the earth absolutely pure. Berthollet found that alumina precipitated from alum, even by pure ammonia, and welledulcorated, still contained a small portion of sulphuric acid, as was proved by affording a sulphuret when strongly heated with combustible matter. He, therefore, recommends that the first precipitate from alum, by ammonia, be redissolved in nitric acid, purified from sulphuric acid by nitrated barytes, and again precipitated by the same alkali, and, after dueedulcoration, dried in a heat sufficient to expel the ammonia, whereby the earth is now obtained pure.

Alumina has a strong affinity for the vegetable and animal oils. Alum in solution, mixed with soap water, curdles it completely, and the coagulum is an intimate union of this earth with the oil of the soap, forming a soft, flexible mass, which, when dry, remains supple, is insoluble in alcohol, and, after fusion, becomes transparent.

The affinities of alumina for the acids are in the following order: Sulphuric, nitric, muriatic, oxalic, arsenic, fluoric, sebatic, tartareous, succinic, mucous, citric, phosphoric, formic, lactic, benzoic, acetic, boracic, sulphurous, nitrous, carbonic, prussic.—*Pottery and Glassware Reporter*.

UTILIZATION OF BLOOD.

By P. MARGUERITE-DELACHARLONNY.

The author gives the proportion of nitrogen in dried blood as 12.15 to 18.73 per cent., and the phosphoric acid at 1.63 to 2 per cent. The value of the blood of the 43,000,000 animals yearly slaughtered in France he estimates at 21,000,000 francs. The author describes various methods proposed for testing blood so as to prevent nuisance. A mixture recommended for this purpose by the Conseil d'Hygiène de la Seine is as follows:

Sodium sulphate	0.60 kilo.
Crude phenol	0.15 "
Common vinegar	0.15 "
Sulphuric acid	0.025 "
Water	2.500 "
	3.425 "

This quantity is to serve for 100 liters of blood.

QUASSIIN.*

By A. CHRISTENSEN.

The literature upon the chemical constituents of quassia wood is very inconsiderable. In 1806 Thomson† mentioned that by digesting quassia wood with water and evaporating he obtained a very bitter substance, which when warm was soft, and became brittle upon cooling, and which, in his opinion, consisted principally of the pure bitter constituent; it gave a slight precipitate with tincture of galls. In 1811 it was stated by Pfaff‡ that the bitterness was completely extracted by cold water. In 1826 Buchner introduced 1 grain of the alcoholic extract into a wound in a rabbit, which was followed by the death of the animal. Another investigation of the extract by Keller§ contains nothing of importance.

Quassia was first prepared by Winckler|| in 1835 as follows: Three ounces of quassia wood were extracted with 2 pounds of 80 per cent. alcohol. After evaporation in a water-bath the residue was dissolved in water, and the solution filtered and evaporated to the consistence of an extract. The extract was exhausted with absolute alcohol, evaporated almost to dryness, and then again exhausted with boiling water. The light yellow solution, decolorized by animal charcoal, left after slowly evaporating crystals of quassia. Winckler described this substance as crystalline, difficultly soluble in water and freely in alcohol, faintly alkaline, giving a precipitate with tannic acid, and as being obtained crystalline most easily from water.

Subsequently Wiggers¶ prepared quassia by repeatedly boiling 8 pounds of quassia wood with water, evaporating the filtered decoction to 6 pounds, cooling, adding lime, and allowing the mixture to stand a day with frequent shaking, filtering, evaporating nearly to dryness, and then boiling with 90 per cent. alcohol. From the solution, which, according to Wiggers, contained, besides quassia and coloring matter, some sodium chloride and potassium nitrate, the alcohol was distilled off and it was then evaporated to dryness. The residue was dissolved in the least possible quantity of alcohol and mixed with a considerable quantity of ether, which separated much coloring matter. The filtrate was now again evaporated, and dissolved in absolute alcohol and ether, and this repeated until the residue was colorless and free from saline matter. Lastly the ethereal solution was poured into a small quantity of water and left to evaporate spontaneously.

Wiggers found for quassia the formula $C_{22}H_{20}O_6$. According to him it loses 1.3 per cent. of water at 100° C., and 0.37 per cent. more upon fusing. The melting point, he says, lies a little above that of the resin. He found it dissolve in 220 parts of water at 12° C., and it did not neutralize acids. It crystallized by spontaneous evaporation from a solution in alcohol and ether which was mixed with water, and also when a warm saturated alcoholic solution was mixed with boiling water and allowed to cool. The quassia of Wiggers was precipitated by tannic acid, but not by metallic salts.

In commencing the present investigation, Herr Christensen first directed his attention to find a new method of preparation, since the methods of Winckler and Wiggers yielded it in a very impure condition, and the operations for its purification would involve considerable loss. The author tried two methods that are available in the extraction of several bitter substances: (1) treatment of a solution of the watery extract with bone black and exhaustion of this with alcohol; and (2) precipitation of the watery extract with tannic acid, decomposition of the washed precipitate, extraction with alcohol, etc. In both cases, by evaporating slowly, he obtained tabular crystals, having an intensely bitter taste, leaving on combustion no ash, difficultly soluble in water and fairly soluble in alcohol. The solutions gave with tannic acid a dense white precipitate.

Notwithstanding that the quassia is tolerably completely removed by bone black, as is shown by the fact that the filtrate tastes only slightly bitter, the tannic acid method has the advantage that it yields the purer product. The following method was adopted by the author for the preparation of quassia:

Five kilos of quassia were boiled several hours with two successive quantities of water. The united filtrates were evaporated, at first over an open fire, subsequently on a water-bath, to 1½ liters. After cooling this was filtered and the filtrate precipitated with tannic acid, taking care to keep the reaction neutral or faintly acid by the addition of carbonate of soda, since the compound of quassia with tannic acid, like many similar compounds of bitter substances with that acid, is very soluble in it. The precipitate, after the addition of well-washed pipe-clay, is easily collected and removed by filtration. The precipitate after being repeatedly washed was mixed while moist with a considerable quantity of freshly precipitated lead carbonate, and evaporated to dryness in small portions, with frequent stirring, upon a water-bath. The mixture was extracted four times with alcohol, the alcohol distilled off, and the residual liquid evaporated until the separation of crystals of quassia, which were washed and dried between filter-paper. The filtrate yielded more crystals upon further evaporation, and some more quassia was separated by shaking it with chloroform. Lastly, the quassia was recrystallized from ether-alcohol and water.

In the place of lead carbonate, calcium hydrate also was used for the decomposition of the tannic acid compound, and an alcoholic solution obtained that was not so much colored, but the yield was smaller.

Upon evaporating the watery extract of quassia wood, a considerable precipitate separated as a smeary mass. This was treated with freshly precipitated lead carbonate, evaporated almost to dryness, and extracted with 84 per cent. alcohol. After the alcohol had been distilled off, and the liquid sufficiently evaporated, some quassia separated, but in an impure condition.

Quassia is very freely soluble in chloroform, and by shaking an aqueous solution of it with chloroform it can be nearly entirely removed. The author therefore tried to obtain the quassia from the aqueous extract of quassia wood by simply shaking this with chloroform; but the practical difficulty was met with that the chloroform does not completely separate it, and a pasty, slimy mass was formed, the consistence of which was not altered even upon warming. After the chloroform had been distilled off a very impure residue was left, from which the quassia was only

* Abstract of a communication from the Strassburg Pharmaceutical Institute (*Archiv*, xx., 431).

† *System der Chemie*, iv., 47.

‡ *System der Materia Medica*.

§ Buchner's "Repertorium," ii.

¶ Buchner's "Repertorium," liv. (1837), 85, and lxxv. (1839), 74.

|| *Annalen der Pharmacie*, xxi. (1837), 40.

obtained by repeated recrystallizations from ether-alcohol mixed with a little water. Neither can direct exhaustion of the quassia wood by chloroform be adopted in the preparation of the quassia, since much resin is taken up with it, and this can only be separated by the ether-alcohol method, which involves considerable loss. The last two experiments show, however, that quassia exists as such in the wood, and is not a product formed during the treatment.

The quassia, obtained as above described with tannic acid, was purified by recrystallization from the smallest possible quantity of warm alcohol, in which it is so much more soluble than in cold that the solution solidifies into a paste upon cooling. After the crystals had been washed with dilute alcohol and dried between filter-paper and then over sulphuric acid they were white and gave a colorless solution. They were recrystallized once more by the addition of boiling water to a hot concentrated alcoholic solution; the crystals that separated on cooling showed the same melting point as those of the first crystallization, and were used in the further experiments.

With respect to the proportion in which quassia occurs in the wood, the author obtained from 18 kilos of Jamaica wood (*Picramnia excelsa*) 12 grammes of impure substance, equal to 0.6 per 1,000. But in treating a subsequent consignment, both of Jamaica and Surinam wood, he found to his surprise that in repeated experiments upon considerable quantities of both kinds he could obtain scarcely any quassia. A sample of *Quassia amara* wood, that yielded 2.8 per cent. of extract, and one of *Picramnia excelsa* wood, that yielded 2.6 per cent., yielded no quassia. The sample of *Picramnia excelsa* wood that gave 0.6 per 1,000 of quassia yielded 3.2 per cent. of extract. The author is unable to explain the cause of this difference; but if the value of quassia wood is dependent upon the proportion of quassia it contains, the results do not afford much justification for the exclusion of the Jamaica wood from some pharmacopœias.

Pure quassia forms very thin rectangular double refracting crystalline scales. It has an intensely bitter taste, is without smell, permanent in air, and gives neutral solutions with alcohol, water, etc. It melts at 205° with slight puffing—possibly due to the giving off of water—to a yellow resin-like mass, which upon solution and recrystallization is recognized as unaltered quassia.

Water, left in contact with finely powdered quassia several days, with frequent shaking, at 15° C., dissolved 1 part in 1,230. A warm saturated solution left forty-eight hours at 15° C., with frequent shaking, contained 1 part in 735. Boiling water dissolves quassia with difficulty, but much more freely than cold. In alkalies it dissolves with a yellow color very readily upon boiling, and separates unaltered on the addition of acids, becoming again colorless. It is dissolved by alkalies in the cold somewhat more easily than by water, turbidity resulting upon neutralizing the liquid with acids. Boiling alcohol dissolves it very freely, but cold alcohol much less so, 1 part of quassia requiring 30 parts of 84 per cent. alcohol at 15° C. In ether and petroleum spirit it dissolves with difficulty, but very freely in chloroform, of which 2.1 parts dissolve 1 part of quassia. A solution in chloroform is dextrogyre, the specific rotation being $\alpha_D + 37.8$.

Quassia contains no nitrogen. After being dried to a constant weight over sulphuric acid, it did not alter in weight when exposed for an hour to a current of dry air at 100° C., and gave upon analysis results corresponding with the formula $C_{21}H_{42}O_8$.

	Calculated.	Found.
C.....	66.66	66.90
H.....	7.52	7.44
O.....	25.82	25.65

An aqueous solution of quassia is not rendered turbid by metallic salts. A saturated alcoholic solution gives no precipitate with a similar solution of neutral lead acetate, and upon evaporation the quassia separates unaltered. Other saturated alcoholic solutions of metallic salts behaved similarly. Tannic acid produces in aqueous and alcoholic solutions of quassia a white precipitate which is soluble in acids and in caustic and carbonated alkalies.

Dilute hydrochloric and sulphuric acids dissolve quassia in the cold somewhat more freely than water, but 3 to 10 per cent. solutions, even after being heated in sealed tubes to 125° for an hour, did not reduce alkaline cupric tartrate, showing that quassia is not a glucoside.

By treatment of quassia with a 3 per cent. solution of sulphuric acid a new compound was obtained, crystallizing in fine needles, and having only a slightly bitter taste. It melted at 237° C., did not lose all its water of crystallization (four molecules) till 110° C., and when anhydrous gave upon analysis results corresponding with the formula $C_{21}H_{40}O_8$. It was very sparingly soluble in water, required 180 parts of alcohol at 15°, dissolved very freely in caustic potash and caustic soda, and gave no precipitate with tannic acid. There was also a formation of resinous matter that could not be crystallized and possessed an intensely bitter taste.

A strong solution of quassia in chloroform decolorizes bromine, with formation of hydrobromic acid. The author obtained a bromine compound as a white amorphous mass, with a taste even more bitter than that of quassia; this compound dissolved in alkalies with an intense yellow color, distinctly produced by the smallest traces, and at the same time lost its bitter taste almost completely.

An experiment made to test statements as to the occurrence of an essential oil in quassia wood gave negative results, a small quantity of solid white substance that collected on the surface of the aqueous distillate proving to consist of free fatty acid.

MODERN THEORIES OF COLOR.

A good résumé of the opinions at present held in regard to color is given by M. J. Soury, in the *Revue des Cours Scientifiques*. The theory, which may now be called old, though it replaced that of Newton, is the Young-Helmholtz theory. According to this, the retina of man possesses three distinct kinds of nerve fibrils, each of which is capable of distinguishing one of three colors supposed to be fundamental or elementary. One of these sets of fibers is sensible to red alone, another to green, and the third to violet or perhaps to blue. Our entire chromatic scale is made up of combinations of these three primary colors. Every sensation is due to the excitation of the fibers of the retina by these tints, each set of fibers being strongly stimulated by its own special excitant, and feebly stimulated by the others. In the case of red, for example, the terminal apparatus of the fibers adapted for conducting red are powerfully stimulated by the large waves of the less refrangible extremity of the spectrum, while the terminal apparatus of the nerves adapted for green and violet are very feebly stimulated, and the sensation of red is the result. In

like manner, when the sensation of green is perceived, the terminal apparatus of the nerves adapted for conducting the impression of green are stimulated by the waves of moderate amplitude in the middle of the spectrum, while the red and violet fibers respond but slightly to such waves, and the sensation of green results. And so, too, the fibers adapted for conducting violet respond to the short and rapid waves of the more refrangible rays which have only moderate action on the green, and still less on the red conducting fibers. The intermediate colors, such as orange, are due to the simultaneous stimulation of the several sets of fibers to a greater or less extent, and when all these sets are equally excited the impression of white light results.

Lately, a new theory—that of Hering—has been advanced. According to this physicist there are six simple sensations of vision, and the various colors seen result from their different association and blending. These elementary sensations are disposed in couples or antagonistic pairs; black and white, green and red, yellow and blue. On this theory violet is a compound color formed by the blending of blue and red, and in an ordinary perception of color four primary sensations are excited; black and white, and two colors; but the blue and the yellow are never simultaneously excited, nor are the red and the green. When the blue sensation is experienced the yellow is virtual, and, though present, is not perceived by the mind. Black and white are capable of mingling, and produce all the shades of gray, but red and green, or blue and yellow, have no disposition to blend, but rather behave like the poles of an electric battery, and it is in this sense that the school of Hering, of which Happe is the chief exponent, speak of the polarity of colors. The organic substratum of color sensation, according to Hering, is not the terminal apparatus of the nerve fiber itself, but certain hypothetical substances, such as the visual purple, which undergo chemical change. On looking at "white" light, the visual purple, by a photochemical process, is constantly undergoing decomposition and destruction, and is as rapidly restored. Red light decomposes the visual purple slowly, violet light decomposes it more rapidly. Further, on Hering's theory, processes both of disassimilation and of assimilation are supposed to take place, so that when exposure to red light has exhausted or partially destroyed the visual purple, or its mother substance, purpurogene, green light aids in the reparation of this material, and fits the eye again for perceiving red light. Black does the same office for a retina exhausted with white light, and blue for yellow. This theory affords a good explanation of the phenomena of contrast, and of complementary colors. A red wafer placed on a sheet of white paper gives the sensation of a vivid red disk. As long as it is looked at it causes disassimilation of the retinal purple, which is as constantly renewed; but if the eyes be closed, the sensation of red vanishes, and is replaced by green. Why? Because assimilation is now taking place in excess of disassimilation. When one of these states is equal to the other two, the two antagonistic sensations are reciprocally destructive. Hence red and green do not blend to form white. The third theory, originally suggested by Draper, and more recently developed by Preyer and by Charpentier, refers the color sensations we perceive to modifications of temperature. Preyer distinguishes the quantity or intensity, the quality or tone, and the temperature. The intensity depends on the force with which the retina is impressed by the ethereal waves; the quality on the nature of the excitation caused by the vibrations of the particles of ether constituting the waves, the vibrations the rapidity of which increases with the diminution in the size of the waves. In regard to temperature, Preyer differentiates the colors into hot and cold—the warm colors are red, orange, yellow, and yellow-green; the cool colors are green, blue, and violet. Charpentier, who has just completed a series of researches with Landolt, finds that the statements recently made to the effect that certain zones of the retina are specially adapted for the perception of certain colors have been founded on imperfect observations, and that even the most peripheral portions of the retina can differentiate colors providing they are sufficiently intense. Aware of the observations of Munk and Samelson, the former of whom seems to have demonstrated that the seat of chromatic sensation is in the cortical layers of the occipital lobes, while the latter has published a case of hemianopsia, in which the sense of space was preserved in the affected half of the retina while the chromatic sense was lost, Charpentier differentiates in every normal eye a sensibility for light, a sensibility for color, and a visual sensibility or sensibility for form. In every chromatic impression made on the peripheral regions of the retina he further distinguishes three phases: a simple impression of light, a vague indeterminate impression of color, and a distinct impression of the true color. The first phase he thinks clearly proves that the luminous sensation is distinct from that of color, that it is the result of a simple excitation of the optic nerve, but in the second phase the cerebral centers for color are excited, and the reason that a quicker and surer recognition of color is made by the macula and central parts of the retina is only because they are more used. The sense of color and of light being a general function of the retina, he regards neither the rods nor the cones as the retinal elements which are stimulated by chromatic stimuli. He rather looks to the layer of ganglionic cells which lie on the anterior part of the retina as the functionally active elements. It is clear that there is still a wide field for research in this subject open to the scientific explorer.—*Lancet*.

MALTOSE.

This sugar has been very fully investigated since O'Sullivan first defined its principal chemical and physical properties, but unfortunately the results obtained differ much from each other. The latest experiments in this direction are by E. Meissl, whose researches are published in the German *Journal of Practical Chemistry*. On making experiments with various preparations of maltose and using a sodium flame, the author found that its specific rotatory power varies inversely with the temperature and concentration. As a mean of several experiments he gives the following equation for the specific rotatory power of maltose with the sodium flame: $[\alpha]_D = 140.375 - 0.01837 P - 0.095 T$, where T represents the temperature and P the percentage of maltose in the aqueous solution. As previously shown, maltose possesses birotatory powers, 15° to 20° difference being observable between freshly prepared and stale or heated solutions, and for solutions lying between 5 to 40 per cent. The percentage of maltose may be accurately found by multiplying by the factor 0.362 the degree of rotation obtained at 17.5°, a tube 200 mm. (nearly 8 in.) long being employed. The assimilation of the elements of water to maltose is attended with much greater difficulty than in the case of cane-sugar,

and its conversion into dextrose is best attained by three hours' heating with a 3 per cent. solution of sulphuric acid. Under the most favorable conditions, however, only 98.5 per cent. of the maltose can be converted into dextrose, as a point is arrived at where the destruction of the ready formed dextrose proceeds with greater rapidity than the formation of fresh dextrose.

A SIMPLE METHOD OF RECOVERING THE SILVER FROM WASHED EMULSION.

AN easy, efficient, and, at the same time, inexpensive method of recovering the precious metal from washed emulsion is a great desideratum—like to the manufacturer of gelatine plates on a commercial scale and to the amateur, who may possibly have but, comparatively, a few ounces of silver in the accumulated residues of many months' working, yet too large a quantity to sacrifice if it could be easily recovered. Of course, if the water in the emulsion be evaporated, the gelatine can be burnt away in a crucible, leaving an ash and the bromide behind, to be afterward dealt with in the furnace like ordinary residues.

Those who have adopted this course know full well the difficulty there is of getting rid of the gelatine by this means; for, unlike the cuttings of paper, it will not burn of itself, and, when heat is applied to assist the combustion, the gelatine frizzles up and expands, then chars, but is not consumed unless a very strong heat be used, and then only after a somewhat long application. Moreover, the fumes given off are by no means pleasant. Even after the gelatine has been so far destroyed a bulky ash always remains, which takes up considerable room in the crucible, and thereby increases the cost of reducing by necessitating a longer time in the fire, and consequently entailing the consumption of more fuel, as well as requiring a larger proportion of flux.

To obviate this difficulty, various methods have at times been suggested. Captain Abney recommends two plans—one being to boil the emulsion with caustic potash, and the other to boil it with hydrochloric acid. By each of these methods the setting property of the gelatine is destroyed, and the silver subsides. These plans answer the purpose admirably, but many complain of the trouble of the boiling operation when large bulks have to be treated, as in the case of the residues accumulated in manufactories where a large business is conducted.

Another plan suggested is that of mixing sawdust with the waste, and then drying and burning it. By this means the gelatine is more completely consumed than when it is burnt by itself; but in this case there is a large amount of ash produced, which is a disadvantage. In order to overcome this, and to obtain the ash as "concentrated" as possible, after the sawdust is dried it is mixed with more emulsion and again dried. After saturating and drying several times, the mixture is finally wetted with a strong solution of nitrate of potash, and dried again. It will then burn with great energy, evolving sufficient heat in some instances to reduce part of the silver to the metallic state, the carbonate of potash formed by the combustion of the nitrate acting as a flux. This plan, although very efficient, has the disadvantage of involving a great deal of trouble.

About twelve months since, M. Ferdinand Braun, of Dornach, gave, in our columns, his method, which is to precipitate the gelatine with tannic acid, and this, as it subsides, carries down the bromide with it. The precipitate is then dried and burnt in a crucible, and afterward reduced by the refiner. In a recent number of the *Correspondenz Dr. Schlessner* advises a modification of M. Braun's method, by employing a mixture of gallic and tannic acids dissolved in alcohol as a precipitant, and then to add sawdust to the precipitated gelatine and bromide before burning. By each of these plans also we have the ashes from the sawdust, as the charred gelatine, to deal with, which, of course, increases the bulk to some extent, while, moreover, the plan is somewhat costly. It is manifest that any method like that of Captain Abney (by which the bromide is separated from the gelatine) will be a considerable advantage in practice. This, it is true, may be accomplished by the employment of Mr. Plener's apparatus, but few are yet provided with that.

In a recent conversation, Mr. E. W. Foxlee mentioned to us a very simple plan by which he had recovered the silver from a large quantity of emulsion residues by first separating the bromide. It was simply to dissolve the emulsion and then add a considerable quantity of the waste acids from the manufacture of pyroxyline. By this means the gelatine was decomposed, and the pure bromide then subsided in an hour or two, when the fluid portion could be decanted perfectly free from silver. After washing, to free it from acid, the bromide was dried and fused with the carbonate of potash and soda in the ordinary manner. This plan appeared to us so simple and inexpensive that we determined to try some experiments with it ourselves, the result of which we now give for the benefit of our readers.

We took some old emulsion, and, after melting it, poured in some mixed acids that had been used for pyroxyline-making some years ago, and had been reserved for cleansing dirty bottles, dishes, and the like. A somewhat violent action ensued after the addition, red fumes being given off for a short time. In about an hour the bromide had subsided to the bottom of the vessel, and the supernatant liquor could be poured off, which, on testing, was found to be perfectly free from silver. As the waste acids from the manufacture of pyroxyline are not at the command of every one, and to prepare them specially would make the process rather more costly, we determined to try the effect of sulphuric acid alone, using the common kind usually retailed at about twopence per pound. We took six ounces of waste emulsion, which had been prepared at different times when experimenting, and added to it three fluid ounces of the acid, and after well stirring, the jar was filled up with hot water. In an hour the bromide had all subsided in a compact mass at the bottom. The fluid portion was then poured off and tested, but no silver was present. The emulsion we employed in this experiment was somewhat decomposed and had become rather thin; so, to put the method to a crucial test, we took six ounces more of the emulsion and added to it half an ounce of Coignet's "gold medal" gelatine. This made it somewhat thicker than ordinary emulsion. Three ounces of sulphuric acid were then stirred in and the jar filled up with hot water as before; but the bromide subsided as quickly as it had done in the previous experiment. From the manner in which the silver went down in these experiments, we have little doubt that less acid would suffice; but we had no opportunity of determining this, as we had already exhausted our stock of waste emulsion. When the bromide is obtained in an isolated state the silver may be recovered from it in precisely the same way as from the chloride.—*Br. Jour. of Phot.*