any of the extrinsic ocular muscles, nor the levator palpebræ superioris; then, by traction, I tore the tumor from its mooring, which was somewhat adherent to the recti muscles. There was some hemorrhage, but there was no trouble in controlling it with the hemostats. I finally detached the tumor in its entirety.

In incising it, the pathologist pronounced it an adenoma, thinking it to be the lachrymal gland luxated from its bed in a hyperplastic condition, having increased in size and weight. The dimensions of the tumor taken by the calipers were 28 mm. horizontally, 38 mm. vertically, and 22 mm. transversely. Its weight was 19.5 grams. Figures 1 and 2 show the appearance two days before the operation, and Figure 6, ten days after the operation; Figures 3 and 4 show the bulk and shape of the tumor, the illustration being natural size. Figure 5 shows the tumor cut open. A photograph taken four days after the operation (Fig. 7) shows the swelling of the lids not all gone, with complete ptosis and ankylosis of the eyeball, but it has receded into the orbit, and the vision is even better than before the operation. There is a decided convergence of the eye with ankylosis, indicating a paralysis of all the extrinsic ocular muscles except the internal rectus, but these conditions gradually improved as the swelling subsided. A photograph taken two weeks after the operation (Fig. 8) shows the swelling of the lids and the chemosis of the conjunctiva nearly gone and the eyeball receded into the orbit. There is complete paralysis of the levator palpebræ superioris and of all the extrinsic ocular muscles, the patient not being able to lift the lid nor to move the eyeball in any direction. The tearing of the tumor loose from its attachment evidently disturbed the nerves innervating these muscles, besides there has been some intraorbital cellulitis, which accounts for the ankylosis. The patient says that the whole left temporal and supraorbital region half way back is numb and void of all feeling, but the paralysis and anesthesia were transitory and in time passed away. He handed me several photographs taken at different periods of time for the past sixteen or eighteen years. These pictures establish the fact beyond doubt that the tumor did not start until about five years ago, and that it was of slow growth.

The special thing this case teaches us is not to depend too much on the history of the case from a layman's point of view, but to examine closely and to arrive if possible at a correct diagnosis before outlining the treatment.

I was inclined to rest content and to agree with my colleague, and to call the tumor a dislodged lachrymal gland until I submitted it to our pathologists, one of whom called it a fibrosarcoma, and the other an angio-myoma. After this controversy, the tumor was sent to Chicago. Dr. Emil Ries and Dr. Zeit pronounced it an endothelioma or a perithelioma.

This patient last called on me June 1, 1905, and there was no evidence of the tumor returning either locally or metastatically. The muscles had regained in part their function. His vision is now sufficient to read newspaper print. Figure 9 shows the appearance of the eye at this time.

Microscopic examination showed principally that the blood vessels had no definite walls. (Figs. 10 and 11.)

Patent Medicines in Norway.—It is reported that a new law in Norway forbids the newspapers to publish advertisements or in any way to further the sale of any and all foreign patent medicines.

ONE OF THE CAUSES OF IMPERFECTION IN THE RADIOGRAPHIC IMAGE.*

HENRY POWER, M.D.

SPOKANE, WASH.

Two recent articles on the *x*-ray in *THE JOURNAL* of the American Medical Association have interested me much. One in particular, in which the author, Dr. Stover of Denver, touches on the subject of the formation of *x*-rays at parts of the Crookes tube remote from the target. The belief in the formation of such rays is widespread and, no doubt, has some origin in fact. I wish in this paper to show that in some cases at least these rays are not thus produced, and can not, therefore, be held responsible for the bad effects noticed in radiography which are usually laid at their door. That there is, nevertheless, some obscure cause at work which tends to reduce the sharpness of the radiograph or of the fluoroscopic image is undoubted and it may be stated further that the *x*-rays do reach the photographic plate from directions other than that in which the target of the tube lies. In order to demonstrate the true origin of these interfering rays, I have performed three experiments, the results of two of which appear in the illustrations. The reproduction shown in Figure 1 is taken from a negative in which different parts of the plate were treated in three different ways. The part marked A was used to receive an exposure of the metacarpal bones in the ordinary manner. The part faintly showing the fingers of the hand, marked B, was exposed at the same time, but a sheet of lead 1/16 of an inch thick was suspended between the target and the hand exposed and at a distance of twelve inches from the hand and plate. Over the two narrow strips marked C, strips of lead were placed having the same thickness as the first, but in apposition to the plate. It will be noted in this illustration that the shadow C is much more dense than the shadow B, though the lead in each case was 1/16 inch thick. To account for this difference, we must assume that the rays which produced the image seen at B must have passed around the edge of the lead and not through it.

If they passed around the edge, we may assume three possible methods by which they accomplished their results:

1. They may have come from a different source of light in the tube, accessory rays.
2. They may have bent from their normal straight line, refracted rays.
3. They may have been produced by the action of the well-known rays from the target, striking the air and causing it to give off new rays in new directions, induced rays.

I am unable at present to devise an experiment which will distinguish between refracted and induced rays, but in regard to the first form, the accessory rays, that they, on this occasion, do not exist, is proved by the following experiment made with the same tube, coil, etc.

A completely closed lead box, which may be considered as a photograph camera for *x*-ray work, was placed fifteen inches below the target of the tube. The camera itself being five inches deep, with the photograph plate at the bottom, it will readily be seen that the plate was twenty inches from the target. As is well known, it would be impossible to use a lens by means of which to form an image of the source of *x*-rays in the tube, so

* These experiments were performed in February and March, 1905; the paper was read at the April meeting of the Spokane County Medical Society.

that for this purpose resort was had to what is known as pin-hole photography. In other words, the front of the camera was pierced with a minute round hole, which would allow the x-rays to pass through and which at the same time was so small that they would be forced to form an image on the plate at the bottom of the camera. The plate was now prepared in the ordinary way for x-ray work by enclosing it within two light-tight paper envelopes, and then placed at the bottom of the camera. The x-ray was now turned on and a long exposure

Fig. 1.—Printed lightly to show the image of the fingers produced by the lawless rays which have been induced by the lawful rays which affected the plate at A.

Fig. 2.—Print fully exposed from part A of negative from which Fig. 1 was taken.

Fig. 3.—An image of the source of x-rays in an active tube taken by pinhole photography with a lead camera and a plate protected from ordinary light.

given. In examining the resulting negative, it is noted that it is perfectly blank with the exception of one spot (Fig. 3). When one examines this spot one sees that it represents a picture or image of the target of the tube, and, on measuring it exactly, I find that its diameters are exactly one-third of the diameters of the target. In the center of this spot there is a far more dense spot, having a slightly oval shape, and the diameter of about $\frac{1}{8}$ of an inch.

This dense spot represents an image of the active central part of the target, where the cathode rays come to a focus; this measurement multiplied by three gives the diameter of the focus as about $\frac{3}{8}$ of an inch. Over all the rest of the plate no effect has been produced by the action of the rays, though the various parts of the plate must have been exposed respectively to the effect of any x-rays which emanated from the various parts of the tube other than the target. I feel that this negative, therefore, proves that in this particular instance, at least, no x-rays worthy of notice were produced in any art of the tube, with the exception of the target. In the illustration the faint image of the target has been lost and only the image of the central or focus point is shown.

I am therefore forced to the conclusion that the rays which caused the trouble in this work, as illustrated at B in Figure 1, are produced in one of the two other ways, and probably by the regular rays inducing new rays by their effect on the air through which they pass.

OPERATION FOR CANCER OF THE MOUTH.*

W. W. GRANT, M.D.

DENVER.

The advantages claimed for the new operation which I have recommended¹ for carcinoma of the lip are: a better, more useful mouth, and, with less tension, a fuller and more natural lip and expression. There is also the distinct esthetic advantage that the lines of incision are less conspicuous, and in men, the usual victims of carcinoma, are soon covered entirely by beard. Plastic surgery means the free use of the knife in liberal incisions, and infinite patience and attention to details. In some respects this is more important in the surgery of the face than of any other part of the body. I know of no department of surgery which taxes more severely the skill, nice sense of proportion, harmony and delicate technic of the surgeon. In the unusually bad cases I report, the operation is tedious, bloody, of rather extensive proportions, and formidable in appearance during its progress; but results justify every step and effort in its consummation. The disease usually occurs in the lower lip of males. The operation is equally applicable to the upper lip, and to partial or complete excision of the lip.

The ordinary V operation, in common use so long that it seems heterodox to question its standing, is most inappropriate, except in small growths, with large mouths. Though the lips are very elastic, they can not stand the excision of the vermilion border with its broad base and apex on the chin. The sucker mouth results, and the tension and contraction are too great to make a useful mouth or a desirable appearance.

METHOD.

The method of procedure consists of two perpendicular incisions on each side of the growth, connected by a straight transverse incision at the base, which is usually about the crease between chin and lip. This leaves a quadrangular space to be filled by flaps. An incision is now made from each inferior angle of the wound, obliquely downward and backward beneath the maxilla, on a line about equidistant between its angle and apex of chin or symphysis. Its further extension is measured

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-sixth Annual Session, July, 1905.
1. Medical Record, March, 1899.