It is seen that the pepsin has completely destroyed the trypsin, even when the acidity was very slight.

EXPERIMENTS ON PEPSIN:

The mixture is then brought to an acidity of 0.2 per cent. HCl and egg-albumin solution added. The quantity of the coagulable albumin remaining after digestion is:

Composition of the mixture during the preliminary digestion of six hours.	
(a) Pepsin and boiled trypsin	0.1070
(b) Pepsin and boiled trypsin and 0.01 per cent NaOH ..	0.1308
(c) Pepsin and unboiled trypsin	0.1410
(d. The albumin solution, without digestion, contained ..	0.1395

Comparison between (a) and (b) shows that the very low alkalinity of 0.01 per cent. NaOH has destroyed most of the pepsin; while the trypsin (c) has destroyed it completely.

EXPERIMENTS ON DIASTASE:

The preliminary digestion lasted 9 hours. The mixtures contained:

Starch was then added and digested for 8½ hours. The digestion is estimated by copper reduced by the solution:	
(a) Diastase and water	0.2854
(b) Diastase and NaOH 0.03 per cent.	0.1945
(c) Diastase and NaOH 0.1 per cent.	0.0010
(d) Diastase and HCl 0.009 per cent.	0.0121
(e) Diastase and HCl 0.02 per cent.	0.0089
(f) Diastase and trypsin and 0.03 per cent NaOH	0.1511
(g) Diastase and pepsin and water	0.1836
(h) Diastase and pepsin and 0.009 per cent HCl	0.0074
(i) Diastase and invertin and water	0.3134

The comparison of (a), (b) and (c) shows that 0.03 per cent sodium hydrate produces considerable destruction of the diastase; with 0.1 per cent. of sodium hydrate the destruction is practically complete; (d) and (e) show the very deleterious effect of acids; even with 0.009 per cent. (equal to 1/20 that of the gastric juice), the destruction is almost complete; (g) and (h) show the severe destructive action of pepsin; invertin (i) has no effect; comparison of (b) and (f) shows that trypsin also is not very deleterious. Of other digestive ferments, rennin is not destroyed by pepsin; invertin is not injured by pepsin, trypsin and diastase, but is weakened by acids and alkalies.

To recapitulate these results relating to the ferments which are most commonly administered for therapeutic purposes:

At a temperature of 40° C. and in six hours:

Trypsin is greatly injured by 0.056 per cent. HCl, and destroyed by 0.56 per cent. It is completely destroyed by pepsin in 0.112 per cent. HCl.

Pepsin is largely destroyed by 0.01 per cent. NaOH, and completely destroyed by trypsin.

Diastase (in nine hours) is considerably injured by 0.03 per cent. and destroyed by 0.1 per cent. NaOH. It is almost completely destroyed by 0.009 per cent. HCl; it is greatly weakened by pepsin even in neutral solution, but it resists trypsin fairly well.

MIXED SOLUTIONS OF FERMENTS VALUELESS.

These conclusions illustrate strikingly the absurdity of mixing these ferments in solution: If the solution is acid (as in the Elixir Digestivum Compositum of the National Formulary and most of the proprietary digestant mixtures), the trypsin and diastase will be destroyed; if it is alkaline, the pepsin and diastase will disappear, and if, as a last resort, it is made neutral, the pepsin will destroy the diastase, and the pepsin, in its turn, will be digested by the trypsin. At room temperature the process will be somewhat slower than in the thermostat, but the final result, in a very short time, will be the same.

This must be the case from the nature of the ferments, and the actual analytical results of the Council, as quoted in its report, only furnish an added proof of these firmly established facts. The busy practitioner may be excused for not having these data always in mind, but ignorance of these matters on the part of manufacturers is scarcely conceivable. Indeed, the letters received by the Council show that the manufacturers are familiar with the worthlessness of these mixtures,

and the most reputable houses only manufacture them in response to the popular demand. Reform must, therefore, come through the medical profession, whose members should refrain from prescribing these mixtures.

(To be continued.)

Clinical Notes

FURTHER STUDY OF THE NEW INFECTIOUS DISEASE.

LEONARD K. HIRSHBERG, A.B., M.D. (Johns Hopkins).
BALTIMORE.

I have already¹ reported 8 cases of an infectious disease hitherto undescribed in text-books or the literature. Since contributing that study, through the kindness of the authors, I have found two articles which report an analogous fever.² These reports, however, are very obscure and incomplete. They make no mention of important tests in order to exclude the various well known fevers, typhoid, etc. The cases of Dr. Sheffield occurred among children and in epidemic form. There are many points in common both in Dr. Sheffield's and Dr. Happel's cases, yet the data are so very much obscured and faulty that more complete reports must be made before these cases can be all identified as the disease reported here.

Through the kindness of many of my colleagues, I am enabled in the following study to report on eleven histories of similar cases in addition to the nine of my own.

This disease may be defined as an acute infection of sudden onset and unknown bacterial origin. It is characterized clinically by fever, constipation, absence of blood parasites, agglutination with the usual bacteria, rose spots, enlarged spleen, and tympanites. It lasts from ten to twenty days. Its pathology is unknown; there have been no deaths. The history of analogous fevers has been called to my attention by Dr. T. J. Acker of Croton-on-the-Hudson. He says that in forty years' practice he has frequently found this fever and "together with Wilson, Da Costa, John Mason Good, and other great writers" attributes "this simple continued fever to solar heat, great fatigue, excitement, surf bathing, eating crabs, etc."

This disease differs from febricula or ephemeral fever as described by Osler. He mentions this as lasting from two to four days; "in some instances it may continue for a week." Weil's disease has as its striking feature a marked catarrhal jaundice, enlarged spleen and liver. None of these features is present in my cases or their analogues. There is absolutely no similarity between these cases and miliary fever, which occurs with sweats and is very fatal.

According to the correspondence I have received on the subject the disease seems to prevail throughout the United States. Members of the American Medical Association have written to me and described similar cases from Missouri, New York, Connecticut, Virginia, California, Georgia, Kentucky, Pennsylvania and New Jersey.

ETIOLOGY.

Of the 20 cases of which I have notes, 8 occurred in girls and 12 in men. The ages ranged from 16 in the

1. THE JOURNAL A. M. A., 1906, Aug. 11, p. 433.

2. Sheffield, H. B.: Am. Med. Surg. Jour., Dec. 26, 1896; also Happel, T. J.: THE JOURNAL A. M. A., July, 1906.

youngest to 36 in the oldest; 18 were under 28; 12 between 16 and 25. Only three of the twenty had been out of the city on vacations. The others had not left the city within six months.

This disease seems to prevail in the same months as typhoid, malaria and hay fever. No cases have been reported to me, however, for December, January, February, March, April or May.

Cultures made from the blood of three patients yielded no growths. Agglutination tests with young cultures of typhoid and paratyphoid bacilli were negative. Cultures made from the stools have so far proved unsatisfactory.

Very little light could be obtained on the origin of the infection. Some of the patients had never drunk unboiled water or milk. None of the cases could be positively traced to any definite source. They seem to be endemic, but there is so far no evidence that any epidemics occur. No banquets, picnics, or sausage luncheons preceded them.

Although there is no morbid anatomy to describe, the mode of infection seems evidently to arise by way of the intestinal tract. The thick white, slightly moist fur on the tongue, the bad odor to the breath, the rigid constipation, all indicate this. The spleen and liver were at no time palpable in any of the cases. This is important in excluding paratyphoid and typhoid.

SYMPTOMS.

There was no way to ascertain the incubation period from the histories. The onset is markedly acute in all cases. From twelve hours in the shortest to three days in the longest; most of them were twenty-four hours. Epistaxis was not present in any of this series. Constipation was a conspicuous symptom in every case. There are no records of chills or sweating. Six of the twenty patients complained of headaches; none of them had abdominal tenderness at any time. Three of the nine walked into my office with temperatures of 104. They felt perfectly well the day before.

The pulse exhibits a striking peculiarity. It is, if anything, even slower than that of typhoid. The rate varies from 60 to 90; in seven of my cases it never rose above 80. The volume is full, rate regular, quality strong, and tension usually fair. Three exhibited diastolic murmurs.

The tongue is covered with a rough, white fur, in most cases moist. A fetid odor was noticeable in several patients. The fever is a rapidly ascending one and may reach 105.4 without delirium or nervous symptoms. No characteristic eruption whatever appears in the course of the trouble. Herpes was present in two patients.

Even in the second and third weeks of the disease no change occurs in the pulse, character or rate. The temperature works gradually downward, and the morning remissions finally become permanent. Tympanites and abdominal tenderness are never present. In this series, cough and expectoration occurred only once. It disappeared a week before the afebrile period began.

In Case 6 Dr. T. R., after a long struggle with the infective agent, finally some three weeks afterwards passed through a typical typhoid attack. Whether the preceding illness was something else or only a battle between his native immune bodies and the typhoid bacilli is impossible to determine.

Meningeal or nervous symptoms were entirely absent. The only pulmonary signs were those of bronchitis in one case.

One case began with an acute attack of vomiting. This lasted three days, after which the course was just

like the others. Throughout the attack the eyes remain bright and the face flushed. There were no relapses, recrudescences, or chills. Erythema appeared in two patients. Perspiration was absent. Desquamation in fairly large flakes attracted notice in one, and occurred in small amounts in another.

The blood showed no marked changes in 9 cases. The hemoglobin was not reduced and the red count remained materially unchanged. The lowest count was 4,300,000. There was no leucopenia or leucocytosis. The count ranged from 4,500 to 8,500.

The Riva-Rocci systolic pressure was never below 125. Jaundice, hemorrhage, perforation or neuritis were never met with. Renal complications are also rare. Ehrlich's diazo test appeared in 3 of the 7 cases so examined. Pyuria may be present as in one of the series. Albuminuria occurred without casts in 4 cases.

DIAGNOSIS.

From typhoid and paratyphoid fevers the diagnosis should be simple after the tenth day; failure of agglutination tests, entire absence of spleen, characteristic eruption, want of a lengthy incubation period, tympanites and cultures of the bacilli from the blood.

Malaria should give no trouble. The parasites are now so readily recognized by even a second-year medical student that even the hidden parasites of the estivo-autumnal form can no longer escape observation for more than a few days. Ulcerative endocarditis and other forms of pyemia will only confuse for a few days.

There really is no chance to confuse this fever with acute miliary tuberculosis, appendicitis, or the diseases mentioned under history, such as febricula or Weil's disease.

There is really no chance to confuse this fever with acute miliary tuberculosis, appendicitis, or the diseases mentioned under history, such as febricula or Weil's disease.

The condition can thus be considered a distinct clinical entity. It seems really to enter a well-recognized place from the numerous letters received.

PROGNOSIS.

The patients all recovered with no apparent sequelae except the patient who, some weeks afterward, ran a typical typhoid course with a Widal reaction present. Twenty cases, however, are not enough on which to generalize, and prognosis, as well as the other data, must be left necessarily incomplete until additional cases are studied.

The general management of these patients must be left to the hands of skilled nurses just as in the case of typhoid. In my series nothing unforeseen developed to require much interference with Nature. The constipation is very stubborn and must be closely watched.

A typhoid diet of albumin, milk and water, as well as hydrotherapy, seems to be the *sine qua non*.³

3. For another article on this disease the reader is referred to Mellette, U. N.: Med. World, July, 1906.

Acetanilid in Solution of Hydrogen Dioxid.—According to C. H. Wall, in the *American Journal of Pharmacy*, several samples of hydrogen peroxid solution were found to contain nitrobenzol, which results from the addition to the solution of small quantities of acetanilid for the purpose of preserving it. The preservation is effectually secured, and it is probable that the small amount of acetanilid added is not objectionable, but its presence and amount must be declared according to the new law.