medies before mentioned I continued to alternate with calomel for a few days longer, until *sixty-four* worms, the least of them measuring above ten inches were voided *per anum*.—The febrile and othersymptoms having subsided, the before mentioned vermifuges were superseded by chalybeates, barks, bitters, wine, &c. which completely restored my patient's health and vivacity.—Miss M. B. is eighteen years old, of the middle size, of dark complexion, and naturally healthy.

Cases of Disease of the Lungs. By JONATHAN SHERWOOD, M.D.

[Communicated for the New-England Journal of Medicine and Surgery.]

THERE has been so much written on the efficacy of a salivation in pulmonary consumption, that you may deem it superfluous to publish the following observations on that subject. But as there cannot be too much testimony in favour of a remedy that has already rescued from premature death hundreds of the human species who must otherwise at this time have been mouldering in the dust, I am induced to transmit to you a brief history of three cases out of many others that have occurred in my practice in which the utility of mercury has been demonstrated beyond the possibility of a doubt. I am further induced to make this communication in consequence of noticing that mercury has fallen into disrepute among many of my medical brethren as an efficacious remedy in that formidable disease. I therefore, consider myself in duty bound, to exert my feeble efforts to convince them of their mistake and shew them wherein they have erred. The following according to the late Doctor Rush are some of the reasons why they have failed to cure consumption with mercury. He says, It was probably given before the morbid action of the blood vessels was reduced by blood-letting and low diet. 2dly, after the suppurative action in the lungs was so far advanced as to prevent the action of the mercury predominating over it. And perhaps a third cause of our want of success may be derived from our having given the mercury after all the excitability of the system (on which medicine act) had been wholly expended ; or 4thly, from our not giving it with some other medicine calculated by its previous or accompanying stimulus to prepare the system for its impression upon the mouth and salivary glands ; or lastly, from our not giving it in the form as in the antimonial powder in which it is diffused through the mouth in the act of swallowing it, which powder is composed of

Nitre - - - 15 grs.
 Tart. antimon. - - $\frac{1}{6}$ "
 Calomel - - - $\frac{1}{2}$ "

CASE 1. A Miss E. R. aged about 25 years, was attacked with phthisis pulmonalis in the year 1813—the disease was hereditary in her family, she having lost in the course a few years a number of brothers with consumption. When she came under my care she was labouring under the following symptoms—viz: a fever of the hectic kind, accompanied with chilly fits, a severe cough, and copious expectoration from the lungs, pain in the breast, and difficult respiration, night sweats, a full, frequent, and hard pulse, a regular accession of fever in the afternoon with a circumscribed flush on the cheek, and she was so weak as to be confined to her bed.

After reducing the morbid action of the blood-vessels by blood-letting and low diet, I commenced giving the above antimonial powder with an increase of the calomel to three grs. every six hours—after continuing its use for about three weeks, I with much difficulty succeeded in producing a ptyalism. Immediately after a mercurial action was excited in the salivary glands, her fever, cough, expectoration, pain in the breast, and every other symptom of her disease subsided—she became convalescent and was soon restored to her former health and vigour. She married some years afterwards and emigrated to the State of Ohio, a distance of upwards of five hundred miles—since her residence in that country I have not heard from her.

CASE 2. On the 5th December, 1820, I was requested to visit a Mrs. F. aged about fifty years, with the following symptoms—a fever accompanied with cough, and expectoration from the lungs, pain in the breast and side, difficult respiration, a burning dryness in the palms of the hand, and pulse small, hard and tense, and in frequency about one hundred in a minute; night sweats, considerable emaciation, great debility, a regular exacerbation of fever in the afternoon preceded by chilly fits, and there was a circumscribed flush on the cheek; tongue foul, and loss of appetite.

I drew from her arm about ten ounces of blood, which was covered with the inflammatory buff, and as there were symptoms of a morbid state of the stomach and bowels, I gave her a cathartic of jalap and calomel, and commenced giving the tincture of digitalis and laudanum, equal parts according to the directions of Doctor Beddoes of which she took from 20 to 30 drops every eight hours.

6th. Felt somewhat easier in her breast and side—less difficulty of breathing. From this time to the 20th, the tincture of

digitalis and laudanum was continued, and blisters were repeatedly applied to the chest, this treatment afforded much relief, but did not eradicate the disease, and I found it indispensably necessary to adopt some more decisive plan in order to check the progress of this fatal malady—after stating to her the probable fatal termination of her disorder unless she took mercury—she after much persuasion consented to submit to my directions. 3 grs. of calomel was given every six hours, and continued ten days, at the end of which time a slight ptyalism came on when every unfavourable symptom subsided—her fever, cough, pain in the breast, and difficult respiration left her, and she appeared to be convalescent until the 13th of March, when she informed me that the pain in her breast had returned the day previous. At this time her pulse was about 80, small and hard. She was again bled to the amount of 10 ounces, which was still buffy, and the tincture of digitalis with the omission of the laudanum, was again ordered, and another epispastic was applied to her breast.

17th. Found her with less pain in her breast, pulse small, hard and frequent—8 ounces of blood were again taken from her arm, and the tincture of digitalis continued.

23d. I was highly gratified to learn, that the pain was removed, and a great number of boils had made their appearance on different parts of the thorax, and it was a singular circumstance to me, that the ptyalism continued although she had not taken any mercury for ten weeks.

4th April. The salivation has ceased and she requires no more attendance.

17th February 1822. It is now upwards of a year, that she has remained in her usual health, and without a single symptom of the complaint.

CASE 3. May 27th, 1821. I was desired to visit a Mrs. D. aged 36 years, of a florid complexion—She informed me she had been indisposed four years with pain and soreness in her left side, accompanied with cough at intervals, and discharge from the lungs.

I found her labouring under the following symptoms, viz.—A fever of the hectic kind with chilly fits, and expectoration of pus-like matter from the lungs, and respiration was so much impeded that she could not make an inspiration without greatly distressing her, neither could she lie on the affected side. She raised from her lungs every morning a considerable quantity of pus mixed with blood; and it is to me singular, that it came up without cough—she had the hectic flush on her cheek in the afternoon and was much debilitated.

From the above symptoms, I had no doubt of her disease being phthisis pulmonalis—her pulse was not hard or frequent, but there was so great an oppression at her breast and difficulty of breathing that I thought proper to take from her arm a few ounces of blood which gave considerable relief to the pain in her side and difficulty of breathing. By the time the blood had done flowing a large epispastic was applied over the left side of the chest, and I commenced giving her 3 grs. of calomel every four hours.

May 30th. There was very little pain in her side or difficulty of breathing, but the expectoration of pus and fever continued.

June 2d. A slight ptyalism had commenced, and she was so weak as to be unable to rise from her bed. Being determined to give the mercury a fair trial, I continued the calomel with directions to discontinue it whenever the salivation and soreness of her mouth became much increased.

June 8th. Found her with a profuse salivation, and I was highly gratified to find that the more copious the discharge of saliva the stronger she became, for at this time she was able to leave her bed and walk into the garden.

June 12th. The expectoration of pus had nearly ceased and she was without cough, pain in her side, chilly fits or fever, and the mercury was discontinued.

June 18th. I learned that the salivation had suddenly subsided and every unfavourable symptom had returned—she expectorated pus tinged with blood every morning, the pain had returned in her side, and her pulse increased in frequency to between 90 and 100 in a minute, and she was again confined to her bed. The calomel powders were again left for her.

June 20th. The salivation had returned, and the pain in her side was again greatly relieved, and expectoration of pus much lessened—the mercury was ordered to be continued so often as to continue the ptyalism without suffering it to diminish—from 3 to 6 grs. a day were found sufficient for that purpose.

June 25th. The pain in her side and expectoration of pus had nearly subsided, and her pulse was reduced to 70 in a minute, and she was again able to leave her bed and walk about her room. It is here worthy of remark, that when the salivation was suffered to diminish, she not only failed in strength but the pain and other unfavourable symptoms would immediately return, for when the discharge of saliva was so profuse as to wet a small towel every hour, her strength would increase, the pain and expectoration of pus would diminish, and her pulse would be rendered soft and slow—in this situation she continued until the 20th July, being constantly under a ptyalism, which seemed

to do much towards mitigating her disease, but did not effect a complete cure. Being extremely anxious to save so valuable a member of the community as I conceived this woman to be, I began to look about for other remedies, and having noticed in the Philadelphia Medical Museum, vol. 6, page 270, a case of phthisis pulmonalis successfully treated by Doctor Samuel Agnew, with acetate of lead and digitalis, I determined as a last resort to try the lead in this case. I therefore commenced giving her on the 21st of July, a pill composed of

Sacc. Saturni	-	2 grs.
Opium	-	1 "
Ipecacuanha	-	2 "

This pill was given once in eight hours, and continued ten days, when she was attacked with distressing pains in various parts of her body, especially the extremities, accompanied with considerable febrile commotion, which subsided in about 8 days, under the antiphlogistic mode of treatment, when the expectoration of pus, pain in her side, and every symptom of her complaint were removed, and she became convalescent, and in the course of a few weeks returned to her ordinary business.

Champion, Jefferson County, N. Y.

Case by Dr. CHADBOURNE.

[Communicated for the New-England Journal of Medicine and Surgery.]

THE following case forms one of those interesting anomalies in pathology, which, although sufficiently instructive to be made known and recorded, will not enable us to deduce any useful principles that can be applied in practice. In short, it is so unlike any diseases of the uterine system hitherto described, that it is presumed a parallel case cannot be produced from the annals of medicine.

Miss S—y W—, aged thirty-six, who now resides in Andover, N.H., fell at the age of fourteen, and received a considerable injury in the region of the kidneys. Her menstrual health continued good till the age of twenty-seven, when she began to experience severe pain at each period, and small scales resembling bone came from the urethra, attended with much pain in voiding the urine. From that time to the present period, pieces of bone continually increasing in magnitude have been discharged from the urethra at different intervals. Four years since they had become so large that it was necessary to dilate the urethra be-