less. But respiration still goes on; the capillary circulation is unusually active; and although the muscular system generally is in a state of relaxation, the involuntary muscles seem preternaturally excited; both the urine and the fæces may be suddenly voided. This stage of complete insensibility is followed, after a time, by symptoms of congestion of the brain; the respiratory act becomes less frequent; the blood, of a dark-blue colour, stagnates in the capillaries; the extremities feel cold; the lips are blue and livid; and if the animal be not speedily removed into pure air, respiration first stops—then the contractions of the heart become fainter and fainter—lastly, the peristaltic action of the intestinal tube ceases. Upon examining the bodies of those destroyed by the inhalation of ether, there is found great congestion of the vessels of the pia mater, and of the sinuses at the base of the brain; the vessels of the medulla oblongata and pons Varolii are especially turgid with dark-coloured blood; both sides of the heart are distended by fluid black blood. The air-sacs of a frog (exp. 1) were full of air; the lungs of the guinea-pig (exp. 6) were in a state of collapse.

Let us, in the first place, see what inferences of practical importance may be deduced from the foregoing remarks:—

1. The vapour of ether, when inhaled continuously, and in considerable quantity, is capable of destroying life, by producing an accumulation of dark blood in the vessels at the base of the brain.
2. The commencement of this state is indicated by slow and laborious respiration—by failing pulse, cold extremities, and blue, livid lips.
3. The arterial blood is, under these circumstances, of the same dark colour as that contained in the veins.*
4. The slow circulation, or the stagnation, of such a fluid in the cerebral vessels may produce effects which the inspiration of pure air will be unable to rectify.
5. Great judgment, therefore, should be exercised in compelling patients struggling for pure air to inhale through the ether apparatus.
6. But a state of insensibility, produced by the proper influence of the ether, may be safely prolonged for a very considerable time, by permitting the occasional inhalation of pure air, and by readministering the ether only when symptoms of consciousness are beginning to manifest themselves.
7. The involuntary muscles, so far from being in a state of relaxation, as are the voluntary muscles, seem to be preternaturally excited, and to act with increased power.

(To be continued.)

ON

THE PRECAUTIONARY MEASURES WHICH MAY BE EMPLOYED IN DISSECTIONS AND POST-MORTEM EXAMINATIONS.

By THOMAS CATTELL, M.D., M.R.C.S.E.

THE very lamentable death of Mr. Potter, from the consequences of a dissection wound, sufficiently indicates the importance of every precautionary or corrective means being familiar to the mind of the medical functionary on occasions like the present. It is with this view I refer,—

I. To those generally known or ordinarily adopted.

1. Dr. Duncan has suggested the anointing of the hands with camphorated oil, or with simple axunge, before handling the viscera.
2. The oil, or spirits of turpentine, (commercial,) is frequently used in wounds of this kind. Having once severely suffered from a dissection wound, I took the precaution, ever afterwards, of carrying a small bottle of it in my pocket whenever I visited the dissecting-room; and from the opportunities I had of estimating its value, there is no doubt of its being an efficient prophylactic, if immediately applied.
3. Nitrate of silver. The sudden formation of chloride of silver, whenever the nitrate is brought into contact with abraded parts, may perhaps prevent the absorption of an animal poison; or, by decomposing it, render it inert.
4. Nitric acid, holding in solution a due proportion of camphor, stands recommended by Dr. Copland as very efficient.
5. Spirits of ammonia is sometimes used.
6. A strong solution of alum has been advised by Macartney.
7. Adhesive plaster should be applied to old abrasions.

* During the performance of amputation of the forearm, upon a patient under the influence of ether, at St. Bartholomew's Hospital, in the month of March, it was observed that the blood, flowing from the radial, ulnar, and interosseous arteries, was of as dark a colour as that contained in the veins.

II. To others of a more original kind.

1. The chloride of antimony is a powerful caustic.
2. Chloride of zinc, in a deliquescent or liquid state. From my experience in caustic applications, I am not aware that I could point to one that exerts a more penetrating influence than this does. Should a case of hydrophobia fall under my care, immediately after the wound had been inflicted by the animal, the first application should be, the deliquescent solution of chloride of zinc.
3. Creasote may be found deserving of trial.
4. Concentrated solution of the chloride of soda, or of lime. A diluted solution of chloride of lime ought always to be on the dissecting-table, as well as by the side of the post-mortem examiner, if the motto of prevention be better than cure.

Lastly, I shall refer to the necessity of the use of gloves made of vulcanized caoutchouc, or Indian-rubber, which should be kept on finger or hand-blocks when not in use.

Braunston, Northamptonshire, June, 1847.

Original Contributions to The Lancet in Abstract.

MEDICINE, SURGERY, MIDWIFERY.

A New Stethoscope.

J. S., of Chelsea, states that the stethoscope which he uses, consists of a cylinder of light porous wood, about ten inches long, two inches and a half in diameter, and half an inch thick. Over one end there is a piece of parchment tightly stretched, with the means of keeping up the requisite amount of tension; the other end is fitted with a projecting rim sufficiently large to enclose the whole of the ear. J. S. is of opinion that an instrument thus constructed is better adapted for the transmission of sound than is the stethoscope in ordinary use.

Congenital Wry Neck.

Mr. LORY MARSH, of Nottingham, was consulted in the case of a girl, eleven years of age, who was afflicted with congenital wry neck. When the patient was seen by Mr. Marsh, the neck presented a completely twisted appearance, the head nearly resting on the right shoulder, with the face turned outwards and downwards, the sterno-mastoid muscle being rigid, tendinous, and shorter by nearly two inches than the muscle on the left side, and the left scapula projecting considerably. The sterno-cleido mastoideus was divided soon afterwards by the subcutaneous section, about two or two inches and a half above the sternum; a pasteboard collar, softened and padded, was placed round the neck, and the whole supported by a figure-of-8 bandage. All going on well, the apparatus was removed at the end of a fortnight; the muscle then presented a thickened, irregular feel, which has since gradually disappeared, and the head is now exactly in the mesian line.

Strangulated Congenital Hernia.

Mr. HARVEY, of Castle Hedingham, describes, in a communication which he forwarded to us some time since, the case of a labourer, forty years of age, to whom he was summoned in consequence of his having injured, as he supposed, one of his testicles, by a fall with a weight on his shoulders. Mr. Harvey saw him the day after the accident, and on examination, found the right side of the scrotum distended, not very tender, and having somewhat the appearance of a hydrocele. The cord could be distinctly felt, as it passed through the abdominal ring, and with it a portion of intestine; but it was neither tense nor tender. The act of coughing produced the usual impulse communicated to hernial swellings, but it was not perceptible in the lower and larger portion of the tumour, which evidently contained a fluid, through which the light of a candle was dimly transmitted. The man at this time asserted that the scrotum was not enlarged prior to his fall, although he afterwards acknowledged that he had observed that the right testicle was larger than the other, but as it did not cause him pain, he paid no attention to it. He had had, he said, a small rupture for several years, which always returned readily on his lying down. He used to wear a truss, but had lately left it off. Mr. Harvey endeavoured to return the protrusion, but in this he could not succeed, although the opening in the obliquus externus was large enough to admit the point of the finger. The next day, the swelling being larger, and tender, accompanied with vomiting of dark but not fæculent matter, and with troublesome hiccup, as well as tenderness of the abdomen, the taxis having again failed to reduce the hernia, Mr. Harvey determined to operate. After cutting