Annals and Magazine of Natural History: Series 4

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tnah10>

XXIV.—Catalogue of Zygopinæ, a subfamily of Curculonidæ, found by Mr. Wallace in the Eastern Archipelago

Francis P. Pascoe F.L.S. ^a

^a Pres. Ent. Soc.

Published online: 19 Oct 2009.

To cite this article: Francis P. Pascoe F.L.S. (1871) XXIV.—Catalogue of Zygopinæ, a subfamily of Curculonidæ, found by Mr. Wallace in the Eastern Archipelago , Annals and Magazine of Natural History: Series 4, 7:39, 198-222, DOI: [10.1080/00222937108696350](https://doi.org/10.1080/00222937108696350)

To link to this article: <http://dx.doi.org/10.1080/00222937108696350>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied

upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

dred of them might therefore have been expected to occur; not one has been found.

This peculiar character alone would seem sufficient to separate generically the so-called *Ceratodus Forsteri* from *Dipterus* and *Ctenodus*, and shows very clearly the relationship of the former to *Lepidosiren*, which is provided with two small pointed teeth in front of the upper dental plates*, which latter do not differ much from those of this interesting Australian fish.

EXPLANATION OF THE PLATES.

PLATE XIII.

- Fig. 1.* Mandible, natural size, with the dental plate attached, of *Ctenodus imbricatus*: *a*, dental plate; *b*, glenoid notch.
Fig. 2. Outside view, natural size, of the right ramus, with the dental plate attached, of *Ctenodus obliquus*: *a*, dental plate; *b*, symphysial margin; *c*, glenoid notch; *d*, channel or cavity overhanging by the dental plate.
Fig. 3. Scale, much enlarged, of *Ctenodus elegans*: *a*, posterior or imbricated extremity.

PLATE XIV.

The palato-pterygoid bones, natural size, with dental plate attached, of *Ctenodus tuberculatus*: *a*, anterior extremity of the bone; *b*, dental plate; *c*, palatal side of the bone; *d*, pterygoid side of ditto.

XXIV.—*Catalogue of Zygotipinae, a Subfamily of Curculionidae, found by Mr. Wallace in the Eastern Archipelago.* By FRANCIS P. PASCOE, F.L.S. &c., late Pres. Ent. Soc.

[Plates XV. & XVI.]

FOR the Zygotipinae of America and of the Old World Lacordaire sought to establish two sections depending on the, as a rule, greater breadth of the episterna of the metathorax in the former, and their narrowness in the latter, or, when in the latter the episterna are broad, on there being a funicle of only six joints, and the pectoral canal being absent. But, even with these limitations, the distinction will not now hold good, as several genera have broad episterna, either with a pectoral canal or with a seven-jointed funicle.

Lacordaire, however, was acquainted with but three of these Malasian genera, only one of which (*Arachnopus*) has narrow episterna; and this, with *Sphadasmus*, *Sympiezopus*, and their allies, it seems to me, had better be excluded from the subfamily. It is not at present my intention to swerve from Lacordaire's arrangement; still it may be desirable to show how the New-World Zygotipinae may be differentiated from the

* "Description of the *Lepidosiren annectens*," by Richard Owen, Esq. Trans. Linn. Soc. vol. xviii. p. 341, tab. 27. fig. 2.

normal species of the Old World. Having regard to the aberrant portion, these differences may perhaps be best shown in a tabular form :—

True Zygopinæ.—Episterna of the metathorax broad, parallel, separating the posterior coxæ from the elytra; eyes always large, generally occupying the whole of the head or nearly so.

A pectoral canal; funicle of seven joints. *All the New-World Zygopinæ.*

Without a pectoral canal, or, if with one, then with a funicle of six joints; or with a funicle of seven joints without a pectoral canal. *Old-World Zygopinæ.*

Aberrant Zygopinæ.—Episterna of the metathorax narrow (*Nauphæus*, ex.), often indistinct, leaving the posterior coxæ in contact with the elytra; eyes of moderate size.

Perhaps entomologists only can appreciate the really wonderful collections made by Mr. Wallace, especially when we consider how poor the collections are, even the Dutch ones, which are occasionally sent to Europe. In this family Curculionidæ, to which the Zygopinæ belong, exclusive of the other Rhynchophorous groups, it is probably within the mark to say that he obtained during his travels among the eastern islands not less than a thousand species; and I think I am not far out of the way in estimating the number of new ones at nine-tenths of the whole. But even the knowledge we have thus obtained is insufficient to enable us to place any great dependence on the distribution or limitation of their beetle-fauna. If we suppose that Mr. Wallace has collected even half the species inhabiting these vast regions, extending for more than 4000 miles in length and about 1300 in breadth, it is obvious that in doubling the number results might be obtained which would very considerably modify any conclusions we may arrive at now. For instance, in the great island of Borneo, Sarawak, a small district on the north-west was the only part visited by Mr. Wallace; it is quite possible that on the south-east coast, opposite to and in one part not more than seventy miles from Celebes, the fauna might be found as allied to the "Australian region" of the archipelago as the north-western or, rather, the Sarawak fauna undoubtedly is to the "Indian region"—that is, supposing the Malayan-peninsular fauna is to be called Indian.

Wallace's Straits (as that portion of the sea has been called which separates Borneo, Java, &c. on the west from Celebes and the islands to the east of it as far as New Guinea) may in other branches of the fauna mark off the two regions alluded to; but, it seems to me, for the Coleoptera they do not show any stronger line of demarcation than would probably be found in taking other parts of the archipelago of about equal

extent. Taking, for instance, the 63 species noticed in this paper as belonging to the Malasian fauna, we find 5 genera and 17 species confined to the "Indian," 10 genera and 40 species to the "Australian," and 6 species, belonging to 4 genera, common to both regions, besides 4 other genera (*Dædania*, *Phylaitis*, *Pempheres*, and *Osphilia*) represented in both. They may be tabulated thus:—

	"Indian."	"Australian."	Common to both.
<i>Mecopus</i>	2	4	2
<i>Talanthia</i>	1		
<i>Agametis</i>	2	1	2
<i>Ganyopis</i>	1		
<i>Odoacis</i>	2		
<i>Chirozetes</i>	2	3	1
<i>Dædania</i>	1	1	
<i>Phylaitis</i>	2	2	
<i>Pempheres</i>	1	1	
<i>Emexaure</i>	1		
<i>Heurippa</i>	..	1	
<i>Metialma</i> *	..	1	1
<i>Osphilia</i>	1	3	
<i>Brimoda</i>	1		
<i>Nauphæus</i>	..	1	
<i>Arachnopus</i>	..	10	
<i>Thyestetha</i>	..	1	
<i>Telaugia</i>	..	1	
<i>Idotasia</i>	..	5	
<i>Semiathæ</i>	..	2	
<i>Elichora</i>	..	1	
<i>Xychusa</i>	..	1	
<i>Nyphæba</i>	..	1	
	17	40	6

Again, if we divide this "Australian region" into two groups, the one consisting of the shallow-sea islands of the Papuan group, New Guinea, Salwatty, Aru, Mysol, and Waigiou, the other the deep-sea islands between these and Wallace's Straits, comprising, among others, Celebes, Gilolo, Morty, Batchian, Ceram, and Amboyna, we find the 46 "Australian-region" species, including the 6 common to the two "regions," to be thus distributed:—5 genera and 21 species confined to the shallow sea, 5 genera and 17 species to the deep-sea islands, 8 species belonging to 5 genera common to both groups, and 3 other genera (*Chirozetes*, *Metialma*, and *Osphilia*) represented in both. The table below will show their special distribution:—

* But this genus has species found in India and in China.

	Shallow-sea Islands.	Deep-sea Islands.	Common.
<i>Mecopus</i>	2	2	2
<i>Agametis</i>	1	1	1
<i>Chirozetes</i>	3	1	
<i>Pempheres</i>	..	1	
<i>Dædania</i>	1		
<i>Phylaitis</i>	..	1	1
<i>Heurippa</i>	..	1	
<i>Metialma</i>	1	1	
<i>Osphilia</i>	1	2	
<i>Nauphæus</i>	1		
<i>Arachnopus</i>	5	3	2
<i>Thyestetha</i>	1		
<i>Telaugia</i>	..	1	
<i>Idotasia</i>	2	1	2
<i>Semiathe</i>	2		
<i>Xychusa</i>	1		
<i>Elichora</i>	..	1	
<i>Nyphæba</i>	..	1	
	<hr/> 21	<hr/> 17	<hr/> 8

I do not attach much importance to these tables, or to any of the same character which may be drawn up on the strength of our present materials; but it would not be difficult to show that similar differences would be found to a greater or less extent in other cases. How very few species, for instance, are as yet known to be common to Dorey and Aru! and yet their faunas are probably almost identical.

Of the Australian *Zygopinae* only two species have been described, *Ilacuris laticollis* and *Mecopus tipularius*. Another *Mecopus* and two *Idotasiæ* are in my collection; and these, as far as I know, are all that have at present been found in Australia—as when there are genera common to Malasia and Australia, they are frequently found in regions beyond the two: *Mecopus* and *Idotasia* have representatives in the Fiji Islands and New Caledonia respectively.

As to the habits of the species, Mr. Wallace tells me that the *Mecopi* are always found on dead trees in the forest, love the sunshine, and take flight rapidly. The members of the allied genera appear to have similar habits; but with the *Arachnopodes* it is different; they look like spiders and have the motion of spiders, frequent the edges of leaves, moving rapidly beneath when approached. *Idotasiæ* and their allies are also found on leaves, but are sluggish. The species observed by Lacordaire in Cayenne and Brazil “live exclusively on the trunks of trees, and are usually quiescent. At the approach of danger, they run rapidly round the tree; and when one attempts to seize them, they fall suddenly as though dead. But instead of

dropping to the ground, they take flight in the middle of their fall, and go back to the point from which they departed."

In the following table, which is drawn up on the principle of taking the most obvious characters rather than the most important, it is thought desirable to include two genera not occurring in the Malayan archipelago—one (*Sphadasmus*) represented by an Indian species, the other the Australian genus *Ilacuris*. Thus all the known Asiatic and Australian genera are here enumerated.

a. No pectoral canal, or, if present, not passing the anterior coxæ.	
b. Eyes closely approximate, occupying the whole head or nearly so.	
c. Funicle six-jointed.	
d. Anterior femora linear, elongate.	
e. Scape not nearly reaching the eye	<i>Mecopus</i> , Schön.
ee. Scape extending beyond the eye	<i>Talanthia</i> , n. g.
dd. Anterior femora thickened, not, or only moderately, elongate.	
f. Propectus canaliculate.	
g. Second joint of the funicle very long.	<i>Agametis</i> , Pasc.
gg. Second joint of the funicle scarcely longer than the others	<i>Ganyopis</i> , n. g.
ff. Propectus not canaliculate.	
h. Second abdominal segment as long as the third and fourth together.	
i. Anterior coxæ contiguous.	
j. Anterior tibiæ compressed, curved, or bisinuate .	<i>Dædania</i> , n. g.
jj. Anterior tibiæ straight or nearly so.	
k. Second joint of the funicle twice as long as the first	<i>Pempheres</i> , n. g.
kk. Second joint of the funicle not longer than the first	<i>Phylaitis</i> , n. g.
ii. Anterior coxæ not contiguous.	<i>Chirozetes</i> , Pasc.
hh. Second abdominal segment shorter than the third and fourth together	<i>Odoacis</i> , Pasc.
cc. Funicle seven-jointed.	
l. Tarsi elongate, linear.	<i>Emezaure</i> , n. g.
ll. Tarsi less elongate, the three basal joints gradually broader.	
m. Body rhomboidal or elliptic.	
n. Three intermediate segments of abdomen equal .	<i>Brimoda</i> , n. g.
nn. Second segment of the abdomen as long as the two next together.	
o. Rostrum triangular at the base	<i>Osphilia</i> , n. g.
oo. Rostrum cylindrical at the base.	<i>Metialma</i> , n. g.
mm. Body oblong, parallel	<i>Heurippa</i> , n. g.
bb. Eyes not closely approximate, nor occupying the whole head.	
p. With ocular lobes.	
q. Intercoxal process narrowed	<i>Nauphæus</i> , n. g.
qq. Intercoxal process very broad	<i>Arachnopus</i> , Guér.
pp. Without ocular lobes.	
r. Intermediate segments of abdomen of equal length	<i>Sphadasmus</i> *, Schön.
rr. Second abdominal segment longer than the next two together	<i>Ilacuris</i> , Pasc.

* *Sphadasmus* has been considered to be confined to South Africa ;

- aa.* A pectoral canal passing behind the anterior coxæ.
s. Pectoral canal gradually effaced behind.
t. Rostrum in repose extending to the abdomen *Thyestetha*, Pasc.
tt. Rostrum in repose not passing beyond the intermediate coxæ *Telaugia*, n. g.
ss. Pectoral canal limited behind.
u. Intercoxal process very broad, the posterior coxæ close to the elytra.
v. Metasternum of moderate length, intermediate and posterior coxæ on each side not approximate.
w. Femora canaliculate beneath, receiving the tibiæ in repose *Idotasia*, n. g.
ww. Femora not canaliculate beneath *Semiathæ*, n. g.
vv. Metasternum very short; intermediate and posterior coxæ on each side nearly contiguous.
x. Pectoral canal not extending to intermediate coxæ . . . *Xychusa*, n. g.
xx. Pectoral canal extending to the posterior border of the intermediate coxæ *Elichora*, n. g.
uu. Intercoxal process narrower; posterior coxæ not near the elytra *Nymphæba*, n. g.

MECOPUS.

Schönherr, Disp. Meth. p. 304; Lacordaire, Gen. vii. p. 157.

The new species of this genus described below are all very distinct, and cannot possibly be confounded with any others. This is not the case with *M. bispinosus**, Weber, on which the genus is founded, and from which I cannot separate *M. Audineti*, Rld. Of the numerous examples in my collection, from seventeen localities, extending from Java and Singapore to New Guinea, I am unable satisfactorily to separate apparently different forms which, if taken without the intermediate individuals, would seem to represent good species. Here "natural selection" may be said to have failed to strike out connecting varieties, without which any such graduated series of forms can be considered only to represent a single spe-

but the following species, not uncommon in Indian collections, brings it into our table:—

Sphadasmus brahminus.

S. breviter ovalis, fuscus, infra griseo-squamosus; rostro prothorace brevius; funiculi articulo primo secundo duplo longiore, tertio secundo fere æquali; prothorace transverso, subconico, regulariter convexo, medio et utrinque griseo-squamoso; elytris profunde striato-punctatis, interstitiis latis, convexis, basi figura transversa triloba, plaga laterali et maculis conjunctis in medio et pone medium sordide griseis notatis; pedibus squamis filiformibus vestitis. Long. 4½ lin.

Hab. Bengal.

* *Rhynchænus bispinosus*, Fab. (Syst. El. ii. 475), quoted in Schönherr, is surely very different from this. Fabricius connects it with *Ctenus*.

cies. It may be added that there is no ground for believing these varieties to be local subspecies. The descriptions below are all made from males; and it will be seen that two of them are without pectoral spines. There are a few other species in Mr. Wallace's collection; one of them is *M. trilineatus*, Guér., which was found in the islands of Aru and Waigiou as well as in New Guinea.

Mecopus spinicollis. Pl. XV. fig. 8.

M. sat breviter obovatus, ater, supra maculatim albo-squamosus, infra squamis albis dispersis; rostro prothorace duplo longiore, supra quinquecarinulato, basi silaceo-squamoso; antennis nigris, funiculo articulo secundo quam tertio duplo longiore; prothorace transverso, ampliato, medio antice paulo gibboso, in utroque latere cavitate sat profunda impresso et extrorsum spinis duabus erectis instructo, supra granulis opacis plurimis sat vage inter squamas detectis; elytris basi prothorace angustioribus, dorso planatis et versus scutellum paulo excavatis, sulcato-punctatis, interstitiis 4°, 5°, 6° spinis brevibus uniseriatim remote armatis; cornibus rectis, subulatis; pedibus minus elongatis, albo-squamosis, concinne nigro annulatis. Long. 5 lin.

Hab. Java.

In my specimen there are only two, but in the British Museum there is one with four or five shorter spines on each side. In the female they are replaced by tubercles.

Mecopus cuneiformis.

M. oblongus, postice gradatim attenuatus, nigro-piceus, silaceo-squamosus, femoribus elytrisque aliquando squamis niveis irratis; rostro prothorace duplo longiore, quinquecarinulato, basi genisque silaceo-squamosis; antennis nigris, funiculo niveo-piloso, articulo secundo quam tertio triplo longiore; prothorace plus minusve ampliato, utrinque fortiter rotundato, supra subvage granulato; elytris anguste cuneatis, dorso planatis, striato-punctatis, interstitio quinto uniseriatim spinoso, reliquis exterioribus asperato-elevatis, apicibus truncatis, angulo externo tuberculato-productis; corpore infra dense albo-squamoso, maculis fuscis dispersis; femoribus supra silaceo-squamosis, reliquis albo sparsis. Long. 4-5 lin.

Hab. Sarawak.

In one or two of my specimens there are brown spots on the middle of the elytra. What I take to be the female has the apices of the elytra rounded. Another species very close to this, from Java, has the elytra mucronate in both sexes; but my male specimen is in too poor a condition for description.

Mecopus pulvereus. Pl. XV. fig. 3.

M. oblongus, postice gradatim attenuatus, prothorace squamulis, subtus elytrisque indumento, griseis dense tectus; rostro capite cum prothorace vix longiore, supra quinquecarinulato, squamulis griseis fere usque ad apicem irrorato; antennis dense griseo-pilosis, clava, basi excepta, nigra, funiculo articulo secundo quam tertio triplo longiore; prothorace sat amplo, apice angustato, squamulis rotundatis tecto; elytris cuneato-cordiformibus, apicibus rotundatis, striato-punctatis, interstitiis omnibus granulato-spinosis, marginibus exterioribus nigris; cornibus nullis; pectore longe lanoso; pedibus anticis quam reliquis multo longioribus; femoribus, præsertim posticis, incrassatis, his corpus vix superantibus, griseo-squamosis; tibiis tarsisque minus squamosis. Long. $4\frac{3}{4}$ lin.

Hab. Macassar.

The same wedge-shaped outline as the last, but differently clothed, and the apices of the elytra rounded.

Mecopus tenuipes.

M. obovatus, fuscus, squamis grisescentibus albidisque variegatus; rostro prothorace fere triplo longiore, basi (præsertim lateribus) crebre rugoso-punctato, infra denticulato-granulato; antennis piceis, funiculo modice tenuato, articulis duobus basalibus fere æqualibus; prothorace valde transverso, ampliato, apice abrupte constricto, medio æqualiter fusco-squamoso, vittis tribus margineque basali subalbido-squamosis; scutello oblongo, griseo; elytris subcuneatis, supra subdepressis, apicibus rotundatis, interstitiis valde convexis, tertio postice bi-tridentatis, dorso griseo subvittatis, singulis pone medium macula fusca obliqua notatis; cornibus piceis, longiuseculis, paulo recurvatis; corpore infra atro, albo marginato, abdomine segmento primo pone coxas brevi; pedibus tenuissimis; femoribus squamis griseis irroratis, posticis perparum incrassatis; tibiis posticis dense albo-squamosis. Long. 5 lin.

Hab. Dorey; Aru.

There is a vestige of another stripe on each side of the prothorax, making five in all; there are also a few nearly obsolete brownish spots on the base of the elytra; but in this genus colour is very uncertain*.

* Another species in my collection, from one of the Fiji islands, is very like the above in coloration and outline, but, *inter alia*, is finely granulated on the prothorax; it may be described as follows:—

Mecopus collaris.

M. forma et colore *M. tenuipedis*, sed rostro minus punctato, basi quadrangulari; antennis valde tenuatis, funiculo articulo secundo quam primo sesquolongiore; prothorace supra manifeste punctato, interstitiis tenuiter granulatis; elytris sutura postice subseratis, interstitiis 3°, 5°, 7°, granulis elongatis subapproximatis instructis; abdomine segmento primo longiore; femoribus posticis linearibus, nullomodo incrassatis; tibiis posticis dense nigro-squamosis. Long. 5 lin.

Mecopus serrirostris.

M. subellipticus, ater, squamis luteis, vel aurantiacis, albisque variegatus; rostro prothorace vix duplo longiore, squamoso, quinquecarinulato, carinulis, media excepta, subremote breviter spinosis, squamis carinulisque prope apicem evanescentibus; antennis tenuissimis, piceis, funiculo articulo secundo quam tertio quadruplo longiore; clava longe pedunculata; prothorace modice ampliato, granulis paucis inconspicuis instructo, atro, linea media nivea, lateribus luteis, sed plaga atra plus minusve ampliata signato; scutello niveo; elytris sat late subcuneatis, medio magis convexis, apicibus rotundatis, striato-punctatis, punctis oblongis, interstitiis angustis, acute elevatis, præsertim tertio pone medium, granulatis, granulis (propter squamas, vix conspicuis) atris, sutura vittisque duabus utrinque obliquis albis luteo lavatis; cornibus nullis; pectore integro, aurantiaco-lanoso; metasterno abdomineque atris, lateribus albo-squamosis; pedibus anticis elongatis, atris, femoribus intus aurantiacis, apicem versus apiceque ipso atris; femoribus intermediis et posticis (his modice incrassatis) plerumque aurantiacis. Long. 6 lin.

Hab. Batchian, Dorey, Saylee.

The yellowish tints of this species are more or less of a faded orange, but brightest on the breast, where, in the male, the scales are long and woolly-looking.

Mecopus lituratus.

M. obovatus, ater, squamis albis maculatim variegatus; rostro prothorace duplo longiore, valde arcuato, basi quinquecarinulato, utrinque vage squamoso; antennis subtestaceis, extrorsum infuscatis; funiculo articulo secundo quam tertio duplo longiore; prothorace sat brevi, vix ampliato, apice latiore, subtiliter creberrime punctato, maculis tribus albis basi notato; scutello oblongo, angusto; elytris breviusculis, utrinque modice rotundatis, striato-punctatis, interstitiis alternis elevatis, sat vage granulatis, apicibus rotundatis, basi suturaque, scutello incluso, literam T formantibus, maculis plurimis albo-squamosis decoratis; cornibus subulatis, paulo recurvatis; corpore infra atro, sparse ochraceo-squamoso, ad latera squamis magis condensatis, metasterno utrinque plaga magna atra notato; pedibus anticis posticisque longitudine æqualibus; femoribus, posticis exceptis, modice incrassatis, his infra dente magno armatis. Long. 4 lin.

Hab. Tondano, Macassar, Sarawak.

TALANTHIA.

Rostrum elongatum, tenuatum; *scrobes* subterminales. *Antennæ* longissimæ; *scapus* ultra oculum productus; *funiculus* 6-articulatus, articulis valde elongatis; *clava* ovata, basi pedunculata. *Prothorax* subquadratus, utrinque paulo rotundatus. *Scutellum*

parvum. *Elytra* prothorace paulo angustiora, basi supra planata. *Pedes* longissimi, lineares; *femora* mutica; *tibiæ* fere rectæ; *tarsi* articulo basali longissimo. *Pygidium* obtectum.

With a habit very much like that of *Mecopus*, this genus is trenchantly differentiated by the unusual length of the antennæ—the scape alone, in (so far as I can judge) both sexes, extending to the posterior border of the eye.

Talanthia phalangium. Pl. XV. fig. 4.

T. oblonga, nigra, opaca; rostro nitido, corpore longiore, basi supra bicarinulato, ad latera sulcato, apicem versus arcuato, depresso; antennis piceis, funiculo articulo secundo longiore, primo tertioque æqualibus, cæteris gradatim brevioribus; prothorace subtiliter punctato, punctis majoribus vage interjectis, maculis septem albosquamosis ornato, una basali lineari, tribus utrinque rotundatis; elytris latitudine duplo longioribus, lateribus sensim angustatis et parum rotundatis, sulcato-punctatis, punctis foveiformibus, interstitiis angustis, convexis, plaga communi scutellari, maculisque quatuor vel plurimis posticis, e squamis albis formatis, notatis; corpore infra ad latera albo plagiato; tibiis tarsisque infra concinne ciliatis, illis apice albis. Long. $3\frac{1}{2}$ –5 lin.

Hab. Penang.

AGAMETIS.

Pascoe, Journ. Proc. Linn. Soc. (Zool.) x. p. 473.

The addition of four new species to this genus necessitates a slight modification of its characters as given by me in the work above quoted. The posterior femora of these species do not extend beyond the body as in the type; and all the femora, except in *A. deleta*, are more or less thickened towards the apex; the elytra, too, though still depressed, are not flattened. None of these species has a trace of the bright orange of their congener, *A. festiva*, but are of different greys and browns, with a few spots of whitish, which, as in other instances where these colours predominate, probably vary according to the individual. The forms of the femora and tibiæ seem to afford good characters.

Agametis agrestis.

A. subovata, fusca, squamis griseis sat dense tecta; rostro rufo-piceo, apice nigro, subtilissime punctulato, basi vage squamoso; antennis subtestaceis; prothorace valde transverso, modice punctato, sat parce squamoso, utrinque vitta indeterminata pallida notato; scutello haud conspicuo; elytris prothorace valde latioribus, sulcato-punctatis, punctis quadratis, interstitiis angustis, convexis, postice, prope apicem, subgibbosis maculisque duabus albidis notatis; corpore infra femoribusque dense griseo-squamosis; femori-

bus posticis incrassatis, medio macula nigra signatis; tibiis tarsisque rufo testaceis, squamis filiformibus albidis vestitis. Long. 4 lin.

Hab. Sarawak.

Agametis deleta.

A. subovata, ferruginea vel fusca, squamis grisescentibus vestita; rostro testaceo-piceo, subtilissime punctulato; antennis subtestaceis, funiculo articulo secundo sequentibus conjunctim haud longiore; prothorace valde transverso, sat dense squamoso, utrinque vitta indeterminata ochracea notato; scutello distincto, subquadrato; elytris prothorace paulo latioribus, sulcato-punctatis, punctis quadratis, squamigeris, interstitiis sat latis squamis elongatis remote uniseriatim munitis, sutura antice macula oblonga et pone medium maculis duabus albidis notata; corpore infra, lateribus thoracis, femoribusque dense albido-squamosis; femoribus posticis perparum incrassatis; tibiis posticis extus paulo arcuatis, anticis, apice solo crassiore, teretibus. Long. $2\frac{3}{4}$ lin.

Hab. Singapore, Sarawak, Morty, Saylee.

Agametis morata.

A. subovata, fusca, flavido-squamosa; rostro nigro nitido, apice rufo-piceo, subtilissime punctulato; antennis subtestaceis, funiculo ut in præcedente; prothorace valde transverso, sat dense squamoso, utrinque flavescente maculis indeterminatis notato; elytris prothorace paulo latioribus, sulcato-punctatis, punctis ovatis, interstitiis sat latis, convexis, squamis elongatis uniseriatim remote munitis, sutura antice, scutello incluso, albido-squamosa, pone medium albo binotatis, maculisque indeterminatis fuscis variegatis; corpore infra pedibusque flavescenti-squamosis; femoribus incrassatis, præsertim posticis; tibiis breviusculis, extrorsum sensim incrassatis, posticis valde arcuatis. Long. $2\frac{1}{2}$ lin.

Hab. Sarawak.

Agametis ortyx.

A. late subovalis, fusca, squamis ochraceis maculatim varia; rostro minus elongato, magis robusto, a medio ad apicem sensim latiore; antennis subpiceis, funiculo articulo secundo sequentibus conjunctim manifeste longiore; prothorace fortiter transverso, plaga media fusca, cruciatim separata, ornato, lateribus pallidis, indistincte fusco maculatis; elytris prothorace magis latioribus, sulcato-punctatis, maculis numerosis ochraceis fuscisque irroratis, quasi subtessellatis; corpore infra pedibusque ferrugineis, sat dense ochraceo-squamosis; femoribus incrassatis, posticis medio macula fusca notatis; tibiis anticis basi fortiter arcuatis, apice angustatis. Long. 3 lin.

Hab. Mysol.

GANYOPIS.

Caput inter oculos cristatum; *rostrum* subvalidum, basi incrassatum, supra compressum, inter oculos ascendens; *scrobes* præmedianæ. *Scapus* oculum haud attingens; *funiculus* 6-articulatus, articulis crassis, tribus basalibus paulo longioribus, cæteris breviter obconicis; *clava* parva obovata. *Prothorax* suboblongus, apice tubulatus et supra haud sinuatus. *Elytra* prothorace latiora, elongata, subparallela, postice subito declivia. *Pedes* mediocres; *femora* parum incrassata, infra dentata, postica longiora; *tibiae* subrectæ; *tarsi* mediocres, articulo tertio fortiter dilatato; *coxæ* anticæ contiguæ. *Propectus* leviter canaliculatus, haud cornutum. *Abdomen* normale, suturis fere rectis.

In habit like *Agametis*, but the funicle different, the rostrum raised at the base, running up and forming a crest between the eyes, and the apical margin of the prothorax not sloped or sinuated above, as in that and many of the genera allied to *Mecopus*.

Ganyopis leucura. Pl. XV. fig. 7.

G. elongata, fusca; rostro prothorace brevior, ferrugineo, basi cristaque inter oculos silaceo-squamosis; antennis piceis, funiculo articulo secundo primo vix longiore, omnibus parce pilosis; prothorace latitudine longitudini fere æquali, fortiter sat crebre punctato, medio fusco, lateribus scutelloque dense silaceo-squamosis; elytris supra subplanatis, latitudine fere triplo longioribus, basi singulatim valde rotundatis, lateribus sensim angustatis, postice utrinque tuberculo conico et, apice ipso, tuberculo minore externo, munitis, fortiter seriatim punctatis, punctis approximatis, quadratis, lineis elevatis separatis, dimidio antico supra parteque declivi dense silaceo-squamosis; corpore infra femoribusque dense albo-squamosis; femoribus posticis macula fusca notatis; tibiis tarsisque minus dense squamosis. Long. 6 lin.

Hab. Malacca.

ODOACIS.

Pascoe, Journ. Entom. ii. p. 427 (1865).

Macrobamon, Lacordaire, Gen. vii. p. 158 (1866).

Lacordaire's description of this genus was based on a female from Ceylon, which is doubtless distinct from the species described by me, as well as from the following. The genus is allied to *Mecopus*, and, owing to the length of the hind femora, is of a remarkable appearance. The male in the type (*O. grallarius**) has the anterior coxæ spined, not the breast as in

* See Pl. XVI. fig. 5.

Mecopus. I have another species allied to it, from Singapore, but it is not quite perfect; it has a longer prothorax, with the posterior angles produced, and the sides of the elytra, for more than a third of their length, dilated, or forming a sort of ledge. The following is very distinct from both.

Odoacis pedestris.

O. ovalis, obscure fuscus, squamis sordide ochraceis sparse et submaculatum indutus; rostro nigro, basi quinquecarinulato; antennis piceis, griseo-pilosis; funiculo articulo secundo quam primo duplo longiore; clava subcylindrica; prothorace transverso, medio carinulato, angulis posticis producto-rotundatis, dorso reticulato foveato; elytris sulcato-punctatis, interstitiis alternis magis elevatis, quarto sextoque granulatis; corpore infra pedibusque squamis albidis irroratis; tibiis, praesertim posticis, valde compressis, his extrorsum nigris; tarsis posticis articulo basali valde compresso. Long. 4 lin.

Hab. Sarawak (and Labuan).

CHIROZETES.

Pascoe, Journ. Proc. Linn. Soc. Zool. x. p. 447.

From a communication lately received from M. Chevrolat, I find that the species described by me, and from which I drew up the characters of this genus, had been long ago published by Wiedemann*, under the name of *Rhynchænus sphaerops*, in a work which, at the moment, I had unfortunately overlooked. Of the species here described, the first three have a certain general resemblance to the type (*C. sphaerops*); so have also, but in a less degree, the last two with one another. The following table will facilitate their recognition:—

Derm brown.

Prothorax longer than broad *sphaerops* (Wiedm.).

Prothorax transverse.

Elytra (♂) scarcely more than half as long again
as broad.

Prothorax granulate *auguralis*.

Prothorax not granulate *junix*.

Elytra nearly twice as long as broad *seclator*.

Derm black.

Apex of the prothorax entire *nervosus*.

Apex of the prothorax sinuate *grammicus*.

* Zool. Mag. Bd. ii. Stück i. p. 129 (1825).

Chirozetes sectator.

C. subellipticus, fuscus, squamis silaceis conspersus; capite inter oculos silaceo-squamoso; rostro nitido, sat fortiter punctato; antennis piceis; prothorace transverso, apice supra modice sinuato, granulato-punctato, squamis silaceis irrorato, in medio et in utroque latere condensatis, vittas tres formantibus; elytris sulcato-punctatis, interstitiis alternis elevatis, reliquis complanatis, maculis silaceis notatis, pone scutellum in singulo elythro macula curvata, postice divergente, signatis; cornibus gracilibus, acutis; corpore infra dense albido-squamoso; pedibus minus squamosis. Long. $5\frac{1}{2}$ lin.

Hab. Sarawak, Penang, Singapore.

Chirozetes junix.

C. breviusculus, fuscus; capite inter oculos silaceo-piloso; rostro sat fortiter punctato, basi vage silaceo-squamoso, apicem versus rufo; antennis testaceo-piceis; prothorace transverso, basi manifeste bisinuato, apice modice sinuato, haud granulato suberebre punctato, squamis silaceis irrorato, in medio carinulato; elytris piceis, squamis silaceis flavescentibusque variegatim indutis, sulcato-punctatis, interstitiis 1°, 3°, 5° elevatis, hoc confertim punctato, reliquis subasperatis, plaga fusca pone medium notatis; cornibus gracilibus, acutis; corpore infra pedibusque albido-squamosis. Long. 4 lin.

Hab. Sarawak, Singapore, Mysol.

Chirozetes auguralis.

C. subovalis, fuscus; capite inter oculos genisque albidis; rostro ferrugineo, nitido, leviter confertim punctato; antennis testaceis, funiculo articulo primo secundo brevior; prothorace magis transverso, basi fortiter bisinuato, subgranulato-punctato, medio obsolete carinulato, squamis albidis irrorato plagisque quatuor notato, duabus apicalibus, duabus basalibus; elytris sulcato-punctatis, interstitiis plus minusve elevatis, praesertim prope basin, et interstitiis 1°, 3°, 5°, his asperatis, albido-squamosis, plaga in medio, ad suturam interrupta, nonnullis minoribus transversis ad latera notatis; corpore infra dense albido-squamoso, flavescente lavato; pedibus squamis albidis vestitis; tibiis posticis intus basi apiceque fuscis. Long. 4 lin.

Hab. Aru.

Chirozetes nervosus. Pl. XV. fig. 9.

C. ellipticus, niger; capite inter oculos genisque niveo-squamosis; rostro sat fortiter punctato; antennis piceis; prothorace latitudine vix longiore, apice supra perparum sinuato, confertim granulato-punctato, vittis tribus albidis notato, una mediana, una

utrinque, infra vitta laterali fuliginea, leviter punctato; elytris striato-punctatis, interstitiis alternis asperatis, nitidis, convexis, lineis albis ornatis, una, scutellum includente, margine basali sita, in singulo elytro una obliqua abbreviata ante medium et prope suturam, una post eam interstitio quinto limitata et fere ad apicem protensa, alteraque exteriore interstitio octavo locata; cornibus pectoris apice dilatatis; corpore infra squamis albidis sat dense vestito, lateribus sterni abdomineque in medio fuligineis; pedibus squamis piliformibus albis minus dense tectis. Long. 6 lin.

Hab. Amboyna.

Scales on the median stripe of the prothorax oval, with a central longitudinal depression, and placed, in the middle, obliquely or almost transversely.

Chirozetes grammicus.

C. subellipticus, fusco-niger; capite inter oculos genisque pallide ochraceo squamosis; rostro sat fortiter punctato, basi valde incrassato; antennis piceis; prothorace transverso, apice supra valde sinuato, granulato-punctato, squamis silaceis irroratis, in medio et ad latera condensatis vittas tres formantibus; elytris ut in præcedente, sed lineis obliquis a scutello incipientibus; cornibus pectoris apice acutis; corpore infra fuligineo, sterno segmentisque abdominis marginibus ochraceo-squamosis; pedibus ut in præcedente. Long. $3\frac{1}{2}$ lin.

Hab. Mysol.

Scales on the median stripe of the prothorax narrowly wedge-shaped anteriorly, towards the base more oval, and without a central depression, arranged as in the last.

DÆDANIA.

Rostrum sat robustum, basi cylindricum, lineatum, apice depressum: *scrobes* præmedianæ. *Oculi* magni, contigui. *Scapus* basin rostri haud attingens; *funiculus* 6-articulatus, articulis duobus basalibus longioribus, cæteris submoniliformibus; *clava* pedunculata. *Prothorax* transversus, ampliatus, basi bisinuatus. *Elytra* prothorace haud latiora, subeuneiformia, supra subplanata, apicibus rotundata. *Pedes* mediocres, intermedii minores; *femora* incrassata, infra fortiter dentata; *tibiæ* compressæ, intus bisinuatae; *tarsi* modice elongati, articulo tertio dilatato. *Coxæ* anticæ approximatae. *Propectus* foveatum, bicornutum. *Epimera* mesothoracis haud ascendente. *Abdomen* normale.

The anterior tibiæ in this genus have the inner edge bisinuate, owing to the tooth-like process at or near the middle;

the anterior femora also are much stouter, and have a large triangular tooth beneath.

Dædania mesoleuca. Pl. XV. fig. 1.

D. obovata, nigra; rostro prothorace paulo longiore, castaneo, fortiter crebre punctato, basi quinquelineato; antennis piceis; prothorace utrinque fere parallelo, subtiliter granulato-punctato, maculis duabus, una apicali, una basali, lateribusque niveo-squamosis; elytris striato-punctatis, interstitiis convexis, uniseriatim granulatis, plaga magna basali subtriangulari, scutellam includente, niveo-squamosa; corpore infra, medio excepto, femoribus basi, tibiisque medio niveo-squamosis, reliquis tarsisque nigris; femoribus posticis corpus superantibus; tarsis anticis dilatatis, ciliatis. Long. 3 lin.

Hab. Mysol.

One of my specimens, apparently a female, has the anterior tarsi also dilated, but to a less extent.

Dædania meleagris.

D. oblongo-obovata, nigra; rostro basi lineis abbreviatis minus elevatis; antennis testaceis; prothorace utrinque rotundato, crebre punctato, squamis silaceis irrorato, maculis duabus, una apicali, una basali, lateribusque etiam duabus, sed majoribus, niveo-squamosis; elytris striato-punctatis, interstitiis convexis angustis, uniseriatim granulatis, squamulis silaceis maculatim dispositis, macula communi pone scutellum, singulatim maculisque tribus, 2 juxta suturam, altera exteriore notatis; corpore infra toto albo-squamoso; femoribus tibiisque minus dense squamosis; femoribus posticis corpus haud superantibus. Long. 2½ lin.

Hab. Sarawak, Singapore, Penang (and Cambodia).

The silaceous scales on the elytra are much smaller than those on the prothorax. A specimen, also from Sarawak, differs in having its markings pure ochre-yellow, and in its longer rostrum less rugose at the base.

PHYLAITIS.

Rostrum tenuatum, basi cylindricum haud compressum, lineatum, apice subdepressum; *scrobes* præmedianæ. *Oculi* permagni, contigui. *Scapus* basin rostri vix attingens; *funiculus* 6-articulatus, articulis duabus basalibus, longioribus, æqualibus, cæteris transversis vel subtransversis; *clava* basi elongata, vix pedunculata. *Prothorax* transversus, basi bisinuatus. *Elytra* prothorace vix latiora, subeuneiformia, apicibus rotundata. *Pedes* elongati; *femora* modice incrassata, infra dentata, postica corpus longe superantia; *tibiæ* graciles, longiusculæ, anticæ rectæ, reliquis

flexuosis; *tarsi* elongati, articulo tertio dilatato. *Propectus* foveatum, bicornutum. *Episterna* metathoracis haud ascendens. *Abdomen* normale.

In this genus the anterior tibiæ are slender, and of equal breadth throughout; the anterior femora are slightly thickened, and have a slender spine-like tooth beneath.

Phylaitis V-alba. Pl. XV. fig. 6.

P. obovata, nigra, albido-squamosa; rostro prothorace manifeste longiore, nigro-piceo, basi quinquelineato, squamis albidis adperso; antennis fusco-testaceis, funiculo articulis duobus basalibus longiusculis, cæteris breviter obconicis; prothorace fortiter transverso, ampliato, squamis albidis irrorato, apice nigro marginato; scutello lanciformi; elytris striato-punctatis, interstitiis latis, sordide albido variegatis, basi squamis majoribus albido-coloratis, literam V formantibus; corpore infra dense albido-squamoso; pedibus squamis filiformibus albis minus dense vestitis. Long. $2\frac{1}{4}$ lin.

Hab. Macassar, Celebes, Ternate, Mysol, Dorey.

Phylaitis lineata.

P. subelliptica, squamis angustis albidis conspersa; rostro prothorace cum capite manifeste longiore, basi quinquelineato; antennis fusco-testaceis; funiculo articulis duobus basalibus longiusculis, cæteris transversis; prothorace subtransverso, vix ampliato, utrinque infra dense albido-squamoso, dorso vittis tribus indistinctis notato; scutello punctiformi; elytris striato-punctatis, interstitiis vage squamosis, duobus juxta suturam octavoque magis dense vestitis, inde lineatis; corpore infra dense albido-squamoso; pedibus fuscis, femoribus tibiisque squamis albis angustis conspersis. Long. $2\frac{1}{2}$ lin.

Hab. Mysol.

Phylaitis pusio.

P. oblonga, rufo-fusca, variegatim niveo-squamosa; rostro nigro, extrorsum rufo-testaceo, prothorace vix longiore, æquilato, basi obsolete trilineato; antennis testaceis; funiculo articulis duobus basalibus breviusculis, cæteris modice transversis; prothorace valde transverso, basi omnino niveo-squamoso; scutello parallelogrammico; elytris minus cuneiformibus, striato-punctatis, basi sutura, fascia obliqua medio, maculisque indistinctis posticis sparse niveo-squamosis; pedibus testaceis, squamis niveis sparse vestitis; femoribus posticis valde ampliatis, infra dente magno instructis. Long. 1 lin.

Hab. Sarawak.

This is the smallest species of the Malasian Zygopinæ, and is a narrower form than its congeners; it has also a

shorter funicle, and the rostrum is a little flattened at the sides at the base.

Phylaitis cyclops.

P. oblonga, fusco- vel testaceo-fusca, griseo-squamosa; rostro testaceo, prothorace manifeste longiore, fere æquilato, basi obsolete lineato; oculis peramplis, toto contiguus; antennis testaceis, scapo rostri basin fere attingente; prothorace valde transverso, ampliato, basi vix bisinuato, dorso sat dense griseo-squamoso, maculis quatuor brunneis transversim locatis; scutello orbiculari; elytris striato-punctatis, griseo-squamosis, singulatim maculis 3 v. 4 ornatis, 2 v. 3 antice ad latera, una pone medio sitis; corpore infra sat dense albido-squamoso; pedibus testaceis, minus dense squamosis; femoribus posticis valde ampliatis, infra dente magno instructis. Long. $1\frac{1}{2}$ – $1\frac{3}{4}$ lin.

Hab. Sarawak.

A slightly aberrant form, owing to the greater length of its scape.

PEMPHERES.

Characteres *Phylaiti* approximantes, sed *scapo* basin rostri fere attingente; articulo secundo *funiculi* quam primo duplo longiore; cæteris obconicis; *femoribus* posticis corpus haud superantibus; *tibiis* anticis flexuosis.

The relative length of the two basal joints of the funicle in the species of this and the preceding genus seems, from its persistence, to be a good primary character for both genera; the form of the anterior tibiæ, however, is decisive of their generic difference.

Pempheres trilineata. Pl. XV. fig. 2.

P. anguste obovata, nigra, lineis albo-squamosis ornata; rostro prothoraci cum capite longitudine æquali, ferrugineo, basi sat fortiter crebre punctato, et medio subcarinulato, apicem versus punctis evanescentibus; antennis testaceo-piceis; prothorace valde transverso, ampliato, apice supra sulcato, leviter granulato-punctato, vittis septem albis ornato, una mediana, tribus in utroque latere sitis; elytris striato-punctatis, interstitiis latis, vix convexis, vittis tribus albis ornatis, una suturali communi, scutellum includente, una dorsali utrinque sitis; corpore infra dense sordide albido-squamoso; pedibus squamis filiformibus albidis vestitis. Long. $3\frac{1}{2}$ lin.

Hab. Batchian, Morty, Amboyna.

Pempheres habena.

P. anguste obovata, nigra, lineis abbreviatis læte ochraceo-squamosis

ornata; rostro prothoraci cum capite longitudine æquali, fusco, basi leviter tricarinato, crebre sat fortiter punctato, punctis apicem versus evanescentibus; antennis testaceo-piceis; prothorace transverso, paulo ampliato, leviter sat crebre punctato, vittis septem, ut in præcedente, sed ochraceis ornato; elytris breviusculis, striato-punctatis, interstitiis latis, regione scutellari, scutellum includente, vitta dorsali utrinque, medio interrupta, ochraceis ornatis, reliquo suturæ pilis albis instructis; corpore infra dense concinne ochraceo-squamoso; pedibus squamis filiformibus albis subvage vestitis. Long. $3\frac{1}{2}$ lin.

Hab. Singapore, Sumatra.

EMEXAURE.

Rostrum longum, tenue, basi incrassatum, compressum, apice depressum; *scrobes* præmedianæ. *Oculi* magni, contigui. *Scapus* oculum vix attingens; *funiculus* 7-articulatus, articulo primo breviusculo, crassiore, secundo elongato, tertio quartoque sensim brevioribus, tribus ultimis obeonicis; *clava* oblongo-ovata, distincta. *Prothorax* oblongus, basi bisinuatus, medio lobo productus. *Elytra* breviuscula, obovata, prothorace paulo latiora. *Pedes* graciles; *femora* in medio incrassata, infra dentata; *tibiæ* subrectæ; *tarsi* tenuati, articulo basali valde elongato, tertio haud dilatato. *Epimera* mesothoracis ascendente. *Pygidium* obtectum. *Abdomen* breve, segmento secundo ampliato, sutura lateraliter valde arcuato.

The diagnostic characters of this genus are the seven-jointed funicle and long linear tarsi; the scales on the species described below are narrow and hair-like. The male has a manifestly shorter rostrum.

Emexaure gallinula. Pl. XVI. fig. 1.

E. elliptico-rhomboidalis, fusca, griseo-squamosa; rostro arcuato, prothorace duplo longiore, basi vage silaceo-squamoso, extrorsum ferrugineo; antennis læte testaceis; prothorace latitudine paulo longiore, sat crebre punctato, medio sparse squamoso, ad latera vitta curvata e squamis formata; scutello orbiculari; elytris striato-punctatis, interstitiis latis, planatis, interstitio suturali squamis longioribus oblique locatis, reliquis squamis griseis, nonnullis saturatioribus variis; corpore infra pedibusque fusco-testaceis, illo sat dense, his sparse albido-squamosis. Long. 2 lin.

Hab. Sarawak.

HEURIPPA.

Rostrum longiuseculum, tenue, basi compressum; *scrobes* medianæ. *Scapus* oculum haud attingens; *funiculus* 7-articulatus, articulis duobus basalibus modice elongatis, cæteris transversis; *clava* ad-

nata, longitudine funiculo æquali. *Oculi* magni, contigui. *Prothorax* suboblongus, basi rotundatus. *Elytra* parallela, prothorace haud latiora. *Pedes* mediocres; *femora* incrassata, subtus dentata; *tibiæ* basi arcuatæ; *tarsi* breves, articulo tertio haud dilatato. *Epimera* mesothoracis ascendentia. *Pygidium* obtectum. *Abdomen* segmento secundo vix ampliato.

This is the only genus with an oblong cylindrical or sub-cylindrical body among all those with a seven-jointed funicle. The figure on the plate is far too broad, and gives a very inadequate idea of the insect.

Heurippa amæna. Pl. XV. fig. 5.

H. oblonga, angusta, nigra; rostro modice arcuato, ferrugineo, longitudine prothoracis, basi utrinque niveo-squamoso; antennis testaceis; prothorace basi longitudini æquali, antice angustiore, crebre punctato, maculis septem niveo-squamosis ornato, una basali lineari, tribus utrinque sitis; scutello ovato; elytris pone humeros paulo incurvatis, sulcato-punctatis, interstitiis subplanatis, cervino-squamosis, vitta suturali postice ampliata lateribusque nigris; corpore infra sat dense niveo-squamoso; pedibus rufo-testaceis, leviter albo-squamosis; femoribus posticis linea nigra obliqua notatis. Long. 2 lin.

Hab. Macassar.

METIALMA.

Rostrum tenuatum, basi cylindricum, squamosum, supra lineatum; *scrobes* medianæ. *Scapus* oculum haud attingens; *funiculus* 7-articulatus, articulo primo crassiore, secundo longiore, cæteris brevioribus et gradatim latioribus; *clava* ovata, adnata. *Prothorax* transversus, subconicus, basi lobo mediano productus. *Elytra* cordiformia, supra planata, apice late rotundata. *Pygidium* detectum, transversum. *Femora* incrassata, præsertim antica, dente valido infra armata; *tibiæ* arcuatæ, intermediæ posticæque flexuosæ, extrorsum incrassatæ, apice oblique truncatæ; *tarsi* mediocres, articulo tertio dilatato. *Epimera* mesothoracis ascendentia. *Abdomen* normale. *Corpus* rhomboideum.

The species on which this genus is founded have a very uniform and distinctive appearance, and apparently are very widely distributed, two of them being found in Bombay and Hong Kong* respectively.

* I take this opportunity to describe them: the first probably marks the range of the subfamily to the north-west; and the other has perhaps very nearly attained the limit to the north or north-east.

Metialma scenica.

M. nigra, supra pedibusque squamis filiformibus flavidis albisque variegata; rostro fusco, nitido, quinquelineato; antennis testaceis, funiculo

Metialma naevia. Pl. XVI. fig. 4.

M. nigra, squamis filiformibus albis variegata; rostro fusco, nitido, apice luteo, basi crassiore, (♀?) supra quinquelineato parce squamoso; antennis subtestaceis, funiculo articulis quinque ultimis transversis; prothorace supra indistincte albo varia, basi macula majuscula alba, margine nigra, quadrifida circumdata; scutello oblongo, distincto; elytris striato-punctatis, interstitiis latis, planatis, basi, linea obliqua media maculisque incertis albo-squamosis, notatis; corpore infra dense albido-squamoso; pedibus minus squamosis; femoribus anticis dente triangulari maximo infra armatis, posticis extus in medio macula fusca signatis. Long. $2\frac{1}{4}$ lin.

Hab. Macassar, Gilolo, Tondano, Java.

There are slight differences among my specimens, but nothing that can be regarded as specific; that from Java has a rostrum not thickened at the base, and, judging from *Mecopus*, probably represents a male.

Metialma novata.

M. præcedenti affinis, rostro magis attenuato, basi in utroque latere obsolete lineato; funiculo articulis quinque ultimis longioribus; prothorace antice minus angusto, medio nigro plagiato, basi macula albida literam T formante; scutello orbiculari, distincto; elytris brevibus, lateribus magis gradatim angustatis, supra albido-

articulis duobus basalibus minus longiusculis; clava magna, ovali; prothorace manifeste transverso, lobo medio triangulari, flavido squamoso, maculis fuscis sex, quatuor ante medium transversim sitis, duabus basalibus, ornato; elytris subcordatis, striato-punctatis, apice late rotundatis et macula nivea notatis, squamis albidis flavidisque intermixtis, maculis nigris concinne dispersis; corpore infra albido-squamoso, segmentis tertio quartoque medio nigris; pedibus nigro annulatis. Long. 3 lin.

Hab. Bombay.

Metialma signifera.

M. nigra, squamis filiformibus flavidis albisque interjectis subvariegata; rostro fusco, nitido, quinquelineato; antennis subtestaceis, funiculo articulis duobus basalibus longiusculis; clava breviter ovata; prothorace modice transverso, basi lobo medio producto, hoc maculis duabus notato, una postica alba, altera flavida, duabus conjunctis linea nigra fere circumdatis; elytris breviter subcordatis, apice latis, leviter emarginatis, striato-punctatis, flavido-squamosis, squamis albis parce intermixtis, maculisque nigris vage dispositis, apice macula læte flavida ornatis; corpore infra albido-squamoso; pedibus flavidis, femoribus apicem versus nigro annulatis. Long. $1\frac{3}{8}$ lin.

Hab. Hong Kong.

squamosis irregulariter nigro maculatis; pygidio fere oblecto; corpore infra dense albido-squamoso, epimeris mesothoracis nigro notatis; pedibus ut in præcedente. Long. 2 lin.

Hab. Aru.

From Saylee there is another species of this genus, probably allied to this, but very much worn.

BRIMODA.

Rostrum modice robustum, prothorace vix longius, basi cylindricum, lineatum; *scrobes* medianæ. *Funiculus* 7-articulatus, articulis duobus basalibus longioribus, cæteris brevibus, gradatim crassioribus, ultimo clavæ adnato. *Prothorax* transversus, utrinque rotundatus, basi perparum bisinuatus. *Elytra* trigona, apice rotundata. *Pedes* breves; *femora* subincrassata, infra leviter dentata; *tibiæ* validæ, compressæ, arcuatæ, vel flexuosæ, apicem versus crassiores; *tarsi* breviusculi. *Epimera* mesothoracis parum ascendentia. *Propectus* canaliculatus. *Coxæ* antiçæ sejunctæ. *Abdomen* segmentis tribus intermediis æqualibus.

A small dull-looking insect at present is the sole representative of this genus, which, however, is one of the most distinct of the group.

Brimoda pagana.

B. subelliptica, fusca, parce rude griseo-squamosa; rostro piceo, basi paulo squamoso; antennis subtestaceis; prothorace modice transverso, apice lato, utrinque manifeste rotundato, parce griseo-squamoso, in medio carina abbreviata valida munito; elytris fortiter striato-punctatis, interstitiis convexis, tuberculato-asperatis, squamis suberectis vestitis; corpore infra femoribusque piceis, dense albido-squamosis; tibiis parce squamosis. Long. $1\frac{3}{4}$ lin.

Hab. Singapore.

OSPHELIA.

Metialma fere congruens, sed *rostrum* basi triangulariter compresso; *tibiis* intermediis posticisque rectis, sublinearibus; et *corpore* magis elliptico.

Whilst *Metialma* contains species nearly homogeneous, in this genus they are more diversified, although there are several which are so alike that, with my present materials, I have not attempted to describe them. Three of these are allied to *O. flavirostris* (two from Sula and one from Sarawak); and two to *O. undata* (one from Makian, the other from Batchian). They are all thinly clothed with narrow scales, so narrow in some as scarcely to be distinguished from hairs. A transverse section of the rostrum at the base would be nearly represented

by the letter V reversed, = Λ ; in *Metialma*, by an O. The antennæ also are situated much nearer the base of the rostrum; the scape is therefore much shorter.

Osphilia flavirostris.

O. elliptica, fuscescens, supra squamulis angustis griseis variegata, subtus æqualiter vestita; rostro fortiter arcuato, subtestaceo, nitido, basi genisque dense flavo-squamosis; antennis subtestaceis, scapo brevi; funiculo articulo secundo quam primo sesquolongiore, cæteris gradatim brevioribus, ultimis transversis; prothorace sat valde transverso, apice haud angusto, basi fortiter bisinuato, lobo medio rotundato, perparum producto, disco plagis quatuor fuscescentibus, cruciatim separatis, signato; elytris subcordatis, striato-punctatis, basi suturaque plus minusve maculisque plurimis griseo-squamulosis; pedibus subtestaceis, griseo-pilosis; femoribus anticis inerassatis, subtus dente valido instructis, reliquis minus robustis, posticis corpus longe superantibus; tibiis anticis gracilibus, modice arcuatis, intermediis posticisque parum compressis; tarsis articulo primo sat elongato. Long. 2 lin.

Hab. Mysol.

Osphilia onca.

O. subrhomboidalis, fuscescens, supra squamulis angustis griseis variegata; rostro ut in præcedente, sed longiore; prothorace etiam simillimo, sed lobo medio truncato; elytris magis late subcordatis, griseo-squamulosis, maculis fusciscentibus plus minusve conjunctis subfasciatim dispositis; corpore infra dense flavido-squamoso; pedibus ut in præcedente, sed femoribus anticis crassissimis, dente magno triangulari apice acute spinoso instructis, posticis corpus haud superantibus; tibiis anticis fortiter arcuatis, ad apicem excurvatis. Long. $1\frac{2}{3}$ –2 lin.

Hab. Morty, Ceram.

Osphilia apicalis.

O. subrhomboidalis, nigro-fusca, pilis flavidis parce vestita; rostro nigro, nitido, dimidio apicali subferrugineo, basi genisque parce silaceo-pilosis; antennis subtestaceis; funiculo articulo secundo quam primo vix sesquolongiore, tribus ultimis ovalibus; clava subcylindrica; prothorace latitudine longitudini in medio æquali, subtiliter confertim punctulato, basi vix bisinuato, lobo medio producto, truncato, lateribus margineque apicali flavido-pilosis; elytris oblongo-subcordatis, striato-punctatis, sutura postice, apice maculisque lateralibus flavido-pilosis; corpore infra nigro, lateraliter flavido-piloso; femoribus anticis valde inerassatis, infra dente oblongo-triangulari armatis; tibiis anticis fortiter arcuatis, prope apicem excurvatis; pedibus reliquis sat tenuatis, omnibus subferrugineis, flavido-pilosis. Long. 3 lin.

Hab. Sarawak.

Osphilia undata. Pl. XVI. fig. 6.

O. subrhomboidalis, fusca, variegatim griseo-pilosa; rostro nigro, nitido, prope apicem ferrugineo, basi parce flavido-piloso; antennis subferrugineis, funiculo articulo secundo quam primo fere duplo longiore, tribus ultimis transversis; clava elongata, subcylindrica; prothorace latitudine longitudini in medio æquali, lobo medio anguste producto, truncato; disco plagis quatuor fuscis, cruciatim separatis, signato; elytris subcordatis, striato-punctatis, griseo-pilosis, maculis fuscis fasciatim dispositis; corpore infra vix dense piloso; pedibus magis elongatis, pilosis; femoribus anticis dente triangulari magno armatis; tibiis anticis fortiter arcuatis, prope apicem excurvatis; tibiis intermediis posticisque tenuatis. Long. 3-3½ lin.

Hab. Batchian.

NAUPHÆUS.

Caput sphaericum; *rostrum* modice arcuatum, depressum; *scrobes* præmedianæ, obliquæ. *Scapus* brevis; *funiculus* 7-articulatus, articulo secundo quam primo multo longiore, cæteris gradatim brevioribus et crassioribus; *clava* ovata, distincta. *Oculi* magni, verticem haud occupantes. *Prothorax* transversus, apice tubulatus, antice utrinque rotundatus, basi medio lobatus, lobis ocularibus manifestis. *Elytra* basi prothoracis haud latiora, subnavicularia, utrinque sensim angustiora, apicibus rotundatis. *Pedes* intermediî minores, antici longiores; *femora* compressa, incrassata, infra dentata; *tibice* compressæ, arcuatæ; *tarsi* modice elongati, articulo tertio dilatato. *Rima* pectoralis profunda, inter coxas anticas terminata, apice (in mesosterno) fornicato. *Episterna* metathoracis lata. *Epimera* mesothoracis haud ascendentia. *Abdomen* normale.

There is some doubt respecting the position of this genus, which, except for its large contiguous eyes, not covered by the ocular lobes in repose, I should have placed with *Cryptorhynchinae*; but, assuming it to be a *Zygopine*, it would, according to Lacordaire's arrangement, take its place with the New-World forms, owing to the breadth of the metathoracic episterna conjointly with the presence of a pectoral canal and a seven-jointed funiculus. For the present I place it after *Sphadasmus*, partly on account of the eyes not occupying the whole of the head, and partly because it leads to *Sympiezopus*, which has a deep pectoral canal. The remarkable insect described below is closely covered above with black opaque spots, in the middle of each of which is a little shining granule: the effect of these among the pure grey scales is to give the upper surface an ashy colour; the under part is of a pure ivory-white.

Nauphæus miliaris. Pl. XVI. fig. 3.

N. oblongo-obovatus, niger, supra granulis nitidis nigro circumdati confertim maculatus, interstitiis griseo-squamosis, medio prothoracis elytrorumque antice linea grisea notatis; capite supra oculos dense squamoso, maculis duabus fuscis decorato; rostro prothorace multo brevior, testaceo, subtiliter remote punctulato, basi squamoso; antennis testaceis; prothorace pone apicem utrinque macula obliqua nigra; scutello distincto, orbiculari; elytris lateribus modice rotundatis, basi pone scutellum paulo gibbosis, in medio dorsi utrinque perparum longitudinaliter excavatis, striato-punctatis, interstitiis latis, vix convexis, confertim uniseriatim granulatis, medio macula rotundata nigra ad suturam approximata, plagiisque duabus oblongis margine externo, ornatis; corpore infra femoribusque densissime eburneo-squamosis; tibiis tarsisque squamis filiformibus minus dense vestitis, his infra flavo-tomentosis. Long. $4\frac{1}{2}$ lin.

Hab. Waigiou, Mysol.

[To be continued.]

XXV.—*Ceratodus*, and its Place in the System.

By Dr. ALBERT GÜNTHER, F.R.S.

THE general external appearance of this most remarkable fish has been described by Mr. Krefft in Proc. Zool. Soc. 1870, p. 221. My observations* are based on three specimens, viz. one without intestines, one fully developed male, and one female which does not appear to have attained to maturity. Differences in the number and microscopical structure of the scales seem to indicate the existence of a second species beside that described by Mr. Krefft as *Ceratodus Forsteri*. Its scales are considerably smaller and more numerous; and it may be named *Ceratodus miolepis*.

The skeleton represents the type which is so well known from Owen's, Bischoff's, and Hyrtl's descriptions of *Lepidosiren* and *Protopterus*. In certain points of detail, such as the arrangement of the bones of the skull, the form of the cerebral and acoustic cavities, the development of the first rib and apophyses generally, the structure of the scapular arch and pelvis, the resemblance of the genera named is perfect; and from an examination of the skeleton alone the conclusion might have been drawn that they belong to the same natural

* The following notes are a short *résumé* of a memoir presented to the Royal Society at the beginning of last month, and containing a detailed description of the entire organization of *Ceratodus*, with the exception of the nervous system.