morning and three hours in the afternoon, coughed when awake, nursed well.

December 28th. Hoarseness and cough continue.

January 8, 1897. Improvement has continued until within a few days. To-day, temperature 104°; much cough; fine, moist râles generally distributed over chest.

January 17th. Convalescence established.

The tube which was used had been made to order several years ago. It is of the same diameter as the one usually used for a child one year old, but is one-fourth of an inch shorter. It was introduced with considerable difficulty, Dr. E. H. Stevens succeeding after several unsuccessful attempts, both by him and myself. As it was feared that extraction of the tube would be difficult, the string was left attached to it, the end being fastened to the cheek by adhesive plaster. The presence of the string in the pharynx probably accounts for the large amount of mucus, which was very troublesome as long as the tube was retained.

The only culture from the throat was taken January 5th, and was negative. The diagnosis of diphtheria is made practically certain, however, by the clinical appearances, and by the facts that during the week previous to this illness a sister had been kept at home because of sore throat, and that on the 2d of January another sister was taken sick, who on the 3d had abundant membrane in the throat, and on the 5th of January gave a culture containing Klebs-Löffler bacilli.

Enemata of peptonized milk and whiskey were given freely. The only medicines used were chloride of ammonium and syrup of ipecac, and these were administered irregularly.

Reports of Societies.

SUFFOLK DISTRICT MEDICAL SOCIETY. SECTION FOR OBSTETRICS AND DISEASES OF WOMEN.

C. H. HARE, M.D., SECRETARY.

REGULAR meeting, Wednesday, March 24, 1897,
Dr. M. Storer, chairman, *pro tem*.

GONORRHEA IN WOMEN.

DR. E. W. CUSHING presented the sociologic side:

I presume this is meant to involve questions not strictly medical; and I feel a little as Cicero may have felt when, in defending the poet Archaus, he asked his legal tribunal to bear with him if he handled the matter in a way not quite customary in the court. So in this reverend body I hope I shall be pardoned if in considering the sociologic aspects I sometimes drift away from the strictly medical side of the question. It is a very broad and important subject, one which within the last five or ten years has begun to be properly appreciated, but which I am afraid is not properly appreciated by the public, or even by the profession at large, at the present time—the immense importance of gonorrhea in women.

Gonorrhea is pre-eminently a disease of civilization, of crowding together. It is probably as old as the aggregation of humanity, and will last as long as humanity does. There are evident references to it in the

Mosaic Law—the segregation of men with a running from the flesh, the pains taken that all women should pass through the period of purification after menstrual discharges, which could have had its origin only in the fact which was observed then, as it is observed now, that immediately after or during the menstrual discharge infectious diseases were most apt to be communicated.

There are many ways in which gonorrhea is distributed in thickly-settled and in rich communities. One of the most important is prostitution, but that is by no means the only method, and in certain respects it is not the most common method. Prostitution has become a business involving a great deal of capital and skill, and commanding medical talent of a certain grade; and the consequence is that in well-regulated houses of prostitution the danger of infection is recognized, the pecuniary loss to the business is appreciated, and strong efforts are made, by the instruction of the unfortunate inmates in the care of their health, against the house getting a bad name. It is the great army of immoral persons outside the houses of prostitution, ignorant girls who have enough modesty not to take the means to secure themselves against infection, and not enough modesty to avoid the opportunity of infection, who are the great spreaders of this disease. In the last ten years I can remember only two cases of actual prostitutes coming from brothels who have required surgical operations, whereas the multitude of other young women thus affected who have passed under my observation has been something painful to contemplate. In Europe the great standing armies are a menace to the health of the community from this disease. I will merely quote a few lines from a paper recently read by Dr. Burr, in which, after referring to the various attempts to limit this disease in India, Holland and Japan, he says:

“The soldiers of the great standing armies of Europe carry this disease to their homes, on furlough or at the end of their term of service, and distribute the mischief to the innocent. Though not afflicted with the attendant evils of large standing armies in America, it is quite within bounds to say that one-half of our male population becomes infected with gonorrhea at some period of life. When we consider what a large percentage remains uncured (Noeggerath believed it to be 90 per cent.), we may know that a vast number of unfortunate wives must become infected also through no fault of their own.”

If we, fortunately, have not great standing armies, we have other evils or methods of rapid distribution in the rapid growth of huge cities and the free communication between town and country, in the immense facility of commercial travel, in the growth of riches and luxury side by side with the spread of poverty and destitution, in the great armies of young women crowded in the cities, living alone and supporting themselves, living on insufficient wages, far removed from all the safeguards which in other times were thrown about women. That is something which has come up in the last ten or fifteen years to a great extent. The cities have increased, and the number of young women living and earning not enough to clothe and feed themselves is something appalling, and they are in contact with other women who have from their male friends clothes and all they wish for. The great aggregation of students in our large cities and great universities, as youths “just free from bar and bolt,”

has increased very much during the last generation. When I was in college 50 to 70 was considered an average number to graduate in a class. Now they number 200, 400, or even more, in the larger colleges, while the whole number of institutions of learning has increased immensely. Of course, out of such numbers of students there is a large proportion given to various forms of dissipation, and about all they really accept and remember of their alleged classical education is the maxim of Horace:

"Nec dulces amores
Sperne puer, neque tu choreas;
Donec virenti canities abest
Morosa. Nunc et campus et aræ,
Lenesque sub noctem susurri
Composita repetantur hora."

It has been argued that gonorrhea is a useful disease and an aid to morality, because it acts in the long run by obstructing the ducts of the epididymis of the too erotic of the men, and by destroying the function of the Fallopian tubes of the unchaste among the women, and thereby acts as an aid to the restraining influences of civilization, by permitting the propagation only of those who are able to restrain their sexual natures according to the requirements of the environment they live in, thereby furthering Herbert Spencer's idea that the future men and women will accommodate themselves to the limitations of civilization. That is probably a long way off. As far as we can see at present, the natural man has not changed very much from the earliest times. As for the ladies, we certainly have the very oldest bit of history of any kind extant to show that there were different kinds of women from the beginning. In the Talmud it is written that before Eve was created Adam had another wife — Lilith, blonde and emotional and sprightly — who would not live with Adam because he was slow and because he was a farmer, and who ran away with an interesting demon. There have been Liliths and Eves ever since. Few men know how women feel on these matters, and they will never tell, and it would be a libel on the sex to tell what women think and say about each other, but we have the authority of poets to think that women are not attracted by men to fall in love with them because they are good. You recognize that Goethe would have made a mistake if he had made Marguerite fall in love with a man who was pious and diligent and good to his mother; what took away her rest and made her heart heavy was:

"Sein hoher Gang, sein edler Gestalt,
Seines Mundes laecheln, seiner Augen Gewalt."

And so it will be forever.

Now, given the fact that gonorrhea is with us to stay, perhaps it would be all very well if it were limited so that it merely affected prostitutes, erotic young women and immoral young women. If it stopped there it would soon purify civilization. It has been said, if we had a jumping toothache fifteen minutes after every sin, there would soon be no more sins. Unhappily it does not stop here; wives are infected, babies are cursed with blindness, little girls become diseased.

A very large proportion of young men acquire gonorrhea. I think that is not subject to dispute. Just how large is the proportion I will not attempt to estimate. What proportion of men once affected with this disease become thoroughly free from it, and after how long a time, is a difficult thing to determine; but so much is known, that a very large proportion of

men who think they are cured are not cured, and infect their wives soon after marriage. Possibly the disease is aggravated by the fact that the natural excitement attending marriage puts an undue strain on the genital apparatus, and an undue congestion, and starts the latent germs into activity. At any rate, there are an immense number of innocent women infected. It is one of those things which has little influence on the people, because nothing is said about it, as the fatality of tuberculosis is not noticed, although if all the tuberculous could be gathered together we should see that it is worse than cholera or the plague. We had a great explosion the other day, and the city and the whole country were horrified. If I could bring together all the ruined lives, and the misery caused by gonorrhea, which I have seen in the last year, and put it together, there would be a worse shock and greater number of miserable victims from this one disease.

I have here very little to do with the pathology of this thing. I will merely run over the way in which it infects the women. There is first the localization in the genito-urinary tract, the glands of Bartholini, vulvitis; it passes into the cervix, into the uterus, into the tubes, makes salpingitis and the conditions which obtain as result of this gonorrhea. I see cases which have resisted treatment, which are brought for operation, and which do not get well under ordinary treatment. I have no doubt from what my friends tell me that a great many cases have it in a lighter way and do get well, and that the tubes may open and discharge, and the patient pass through a period of convalescence similar to that which occurs in epididymitis when the ducts are stopped up. Still, careful observations will have to be made on this subject. There have been great discussions in Germany lately; and Bumm, formerly of Wuertzburg, brought up his observations on treatment. In spite of every kind of treatment, active or expectant, general or local, some 10 per cent. of his cases acquired pelvic disease of a serious nature.

The disease appears to vary in different places and in different communities. I am persuaded that in Philadelphia there is a more virulent form of the disease, with more tendency to suppuration, than here. There is something in the climate, in the race. In China and among the negroes they have a particularly virulent form; and I think what makes the Philadelphia gonorrheal infection so serious is the large amount of negro element in the city, bringing the worst forms of what the darkies call "lady fever" into cultivation, and thereby increasing the virulence of the whole lot. Also, they have had, until lately, no opportunity in the clinics or hospitals of Philadelphia to treat the disease.

What can we, as a profession, do in regard to gonorrhea? The first thing which reformists who wish to cure everything by law attempt is the regulation of prostitution; and that is the one thing which is bound to be a failure in this country. Evils of all kinds would immediately supervene if we tried to invent laws for the thorough regulation of prostitution. It has been tried again and again. Popes and emperors and kings and queens have attempted it, but without avail. Think of the evils which would arise if any attempt were made to regulate this thing by law! There would be a certain amount of false security which would only aggravate the disease. It is not possible to eradicate it in this way, even in houses of prostitu-

tion. I don't think any one who has followed the clinics in Paris and seen the style of doctor, or style of police supervisor, it breeds, would wish to have a similar state of affairs here. There would be demoralization of the police. Prostitution is a great business, with great capital involved, with unlimited opportunities; and from the slight uncovering of the thing in New York you can see what a stew the police would be in, in our government, if they had the system of arresting, confining and examining young women; the hideous chances of blackmail, arresting of innocent women and putting them with prostitutes. The mixing up of doctors with this is a dirty business, and it would drive the prostitutes into clandestine prostitution, which is the worst of any. Then comes the manifest injustice of punishing women and letting men go free. It would not be tolerated. It is not tolerated in England. To undertake any such system all over this country is, in my opinion, impossible, injudicious and unrighteous.

We cannot expect much from laws, for they rest upon public opinion, and until the public opinion is enlightened the laws are inoperative, and when that opinion is sufficiently enlightened the laws will follow of themselves. We have lived to see what a few years ago would have seemed impossible, the public opinion awakened to the question of tuberculosis, so that tuberculous meat and milk are condemned, and the food-supply of cities is cut off except as far as it can come from properly examined animals, and the milk unless it comes from cows that are exempt from tuberculosis. We have seen also an ordinance against spitting in the public cars. The office of the medical profession is to enlighten public opinion and not to propose laws.

Second, it has been proposed to make it incumbent on physicians to report to the board of health all cases of venereal disease under treatment. It is not the business of physicians to do the work of the police without remuneration, and it is against their first principles to go around declaring diseases which their patients do not wish declared. In certain cases where there is no moral stigma involved, as in diphtheria, and where actual danger overlies the community, the profession can be asked to do it, but they cannot be asked to carry out any such regulation as that in regard to venereal diseases.

Third, to make infection with venereal disease a ground for divorce and damages, even if latent at the time of marriage. That is a just provision, but at present it cannot be carried out; and it is a very delicate matter for the profession to take the initiative in saying anything about such things. Their position is to be dragged into the thing if necessary, but not to volunteer information.

Fourth, to make treatment of venereal diseases by druggists a misdemeanor. I don't think the public is ripe for that yet. Such legislation would be futile at present.

What can the profession do, and what is its duty in the matter? In the first place it can instruct itself so that it realizes the gravity of this thing, knows the means of scientific diagnosis, recognizes these diseases in women, really feels the seriousness of gonorrhea, as it has to treat it in young women and meet it in young wives. Next, it can act directly as an instructor of the community, naturally towards the patients themselves, indirectly through school-teachers and Christian asso-

ciations, by oversight over the police, the militia, firemen, colleges, schools, etc., and in that way the knowledge that gonorrhea is a great and threatening evil can be spread abroad among the community. Secondly, when the average young woman knows, and is instructed, that an unclean husband is going to be an imminent danger to her, she is going to refuse to marry such a man. Lastly, when the mother-in-law knows that the "catching cold" of her daughter immediately after marriage, that the "inflammation" is not catching cold, but an infection from a gonorrheal husband, that young man had better have a millstone hanged about his neck and be drowned in the sea. When the average young woman, her mother, sister, cousin and aunt, know that this disease is infectious, and that this operation is all the fault of the young husband, there will be a scorn arise for a dissolute young man such as is exercised against a dissolute young woman. What more can the profession do? They can take means to instruct themselves so thoroughly in the means of diagnosis which exist at present, that a young man before he marries may be able to get a clean bill of health and be able to know whether he will infect his wife or not. I see at least once a week all kinds of ruined hopes and blasted affections from this disease. It is a sad thing.

The profession can learn to do something to cure this disease in men. Unhappily, they cannot do much towards the cure of gonorrhea in women, but they can act by instructing and impressing on every one that it is a serious business. It used to be considered twenty or thirty years ago that gonorrhea in a man was a thing to be laughed about. All the popular misconceptions of to-day are the result of the erroneous teaching of the last generation, or the last but one of physicians. In the last twenty years we have lived down a host of things our grandfathers taught to their patients. If the profession takes gonorrhea seriously, the public will finally take it seriously. There is a great responsibility resting upon those who make scientific tests. It is no light thing to say a man is free from gonorrhea and may marry, unless a person can say it understandingly. It is no small thing for a man to say a patient has gonorrhea when he is not prepared to prove it; he may get himself into trouble. In these cases where gonorrhea affects women, and the relatives rise in wrath and want to know what is the cause of it, a man may get himself into trouble who knows too much. The easiest way out of it is to let them find out somewhere else by hypothetical questions the general facts of infection and the sources of pelvic inflammation and to settle for themselves the question of delinquency. Terrible to suggest as it may be, it is not always the husband who is the guilty party. The part of prudence and the desire to preserve the peace of families forbid us to say what we think, or even what we know.

"Est et fideli tuta silentio merces."

DR. WRIGHT: The pathology of gonorrhea is largely concerned with the gonococcus, its isolation and its occurrence in pathological processes. The gonococcus, which is the infecting organism in this disease, was discovered in 1879 by Neisser in the pus of gonorrhea and of gonorrheal conjunctivitis; and very soon after that he claimed to have discovered it in the exudate of gonorrheal rheumatism. It was early found the organism would not grow upon ordi-

nary culture media and the first successful attempt at its cultivation was made by Bumm, who used human blood serum as the culture medium. Since the work of Bumm to the present decade the cultivation of the organism made very little advances. In 1892 Wertheim gave us a fairly practical culture medium in the use of a mixture of sterile blood serum and agar-agar.

The bacteriological proof of the gonorrheal nature of various inflammatory processes has been a matter of slow development. The most important contribution to the subject was made by Wertheim in 1892. He put upon a definite scientific basis the teaching that pyosalpinx in the majority of cases was due to infection with the gonococcus. Wertheim showed that in 20 per cent. of all cases of pyosalpinx the gonococcus could be demonstrated by proper culture methods. In his paper he sums up the observations upon the occurrence of this organism in pyosalpinx. In 116 cases the gonococcus was observed in 32 while in 72 cases the material was sterile and in 12 various pus organisms were present. Since that time a number of investigations have been made on this subject and his observations have been confirmed. The interesting thing about the results of Wertheim is the fact that so many of these cases are sterile. That is to be explained on the ground that the gonococcus, produces its effects and then dies out. The mode of invasion of the gonococcus in pyosalpinx is evident. It affects the urethra first, next the cervix and the endometrium and then the Fallopian tubes. In recent years it has been shown that the organism can be the cause of endocarditis and of pleuritis. It has been repeatedly demonstrated in cultures from the joints in gonorrheal arthritis and some Vienna pathologists have claimed to cultivate it from cases of periostitis and of other suppurative conditions which were apparently secondary to gonorrhea.

The diagnosis of this condition from the bacteriological standpoint has been practised since the time of Neisser. It has long been regarded as a very simple matter to diagnose with bacteriological methods the presence of the gonococcus infection. Up to recent years the teaching was, that, if you made a cover-slide preparation of suspected pus, you were to make the diagnosis of gonorrheal infection if you found micrococci, of so-called double biscuit shape, arranged inside the pus-cells. No other organisms invade the pus-cells to that extent, nor did the other pus organisms appear in these flattened hemispheres with clear line of demarcation between them. This is not a sufficient ground for the identification of the gonococcus. If one examines pus from any abscess, one is liable to find appearances which are to all practically identical with the characteristic appearances of gonorrheal pus. Every pus coccus whatever, if examined carefully enough, will show this division into two hemispheres. Under the microscope the only reliable point of distinction between the gonococcus and the other pyogenic cocci, is the fact that the gonococcus is not stained when subjected to Gram's method of staining, while the other cocci are stained by this method. This fact shows that the gonococcus differs chemically from the other cocci. No diagnosis of gonococcus infection should be made until it has been proved that the suspected organism does decolorize with Gram.

The recognition of the presence of the gonococcus in the secretions of the genital tract in women, in cases where there is no special exudation of pus, is usually

very difficult and unsatisfactory owing to the frequent presence of large numbers of other bacteria.

DR. REYNOLDS: I do not know that there is very much to say on so trite a subject as the treatment of gonorrhea. I have been confronted with it a good many more times than I should have chosen. I am interested in what Dr. Wright said about the diagnosis. I had come to the conclusion from the clinical standpoint, that it was all very well to have the discharge under the microscope, but that it did not help one practically at all. You seldom get any positive reports from the pathologist on this point, and when you do it is what you do not believe from the clinical standpoint. It affects the vagina, upper genitalia and urinary organs. As regards the treatment of gonorrheal vaginitis I have tried a good many things, and have found nothing so satisfactory on the whole as the dry treatment in the vagina. I have settled down to the routine of having the vagina thoroughly opened in the Sims position, wiped out and cleaned with dry cotton, insufflated with aristol, then dry absorbent cotton and more aristol until filled up, having that done every other day.

As regards the treatment of gonorrheal endometritis, theoretically the thing to do is to curette it. I have no doubt there is a very transitory stage where the curette would do good. Practically I think it does not amount to much. You don't want to use the curette before the endometrium has been attacked. We used to hear a great deal about curetting during the acute stages of endometritis and salpingitis. I think everybody has fallen away more or less from that and personally of late I have dropped curetting these cases after the endometrium was really attacked. I think by the time you are sure you have an endometritis you have a salpingitis beginning, though perhaps you cannot recognize it. I am inclined to think I have seen cases where a salpingitis could have been avoided if the curette had not been used. When it comes to be a salpingitis, you can curette it, but I don't believe in doing it. You can also remove the tubes, and you can let the woman alone; and of the three I think the letting the woman alone is generally the greater triumph in gonorrheal salpingitis during the early stages. A good many come to operation in the end; a good many of them atrophy. The gonorrheal tubes seldom do very badly; and the indication for operation, it seems to me, rests a good deal on whether one or both tubes are involved, on the age and social condition and desires of the woman. It is a question in my mind, whether in case one tube seems to be involved, you improve the chance of saving the other tube by removing that one. If both tubes are involved, I think the majority of women are better off with atrophied tubes and ovaries than with no tubes and ovaries. It makes a good deal of difference to a woman mentally. There is a difference in the marital relation.

As regards the invasions of the urinary tract, the urethritis does not amount to much in women, as a rule. It is an acute affair that passes off. Cystitis is a little more important; it ordinarily subsides into a chronic sort of affair which is not very troublesome. The uretero-pyelitis which used to be regarded as an extremely rare affection has in the last year been shown to be pretty frequent though ordinarily a mild affair. It may in time break down a woman's health a good deal. I think the acute cystitis is treated best by putting the woman into the knee-chest position with as

empty a bladder as you can get, letting the bladder fill with air through a Kelly speculum and then filling it with a one-half-per-cent. solution of nitrate of silver, letting that stay a few minutes and then washing it out with salt solution. As you look at the bladder you will see two or three such applications as that clear up the bladder and leave only a few localized spots which you can treat with strong applications. There is the most marked difference between filling the bladder with a solution of nitrate of silver in that position, and washing out with nitrate solution in the ordinary way. Get it fully distended, and the solution will attack the whole surface, and there is a wholly different effect. I don't think you can hurry a uretero-pyelitis much when you get it. The case in the course of a year or two may improve a good deal under appropriate medicinal treatment, it may even get well. I think the preventing of these chronic ureteral affections is the most important indication for interfering with a gonorrheal cystitis.

DR. WASHBURN: I can only reiterate what has been said. I think the most important matter in the whole question is the matter of preventing the thing and not treating it after it gets started. The difficulty of treating the conditions is so great and unsatisfactory in their results that the matter comes down to the importance of preventing the attacks primarily, and that I am persuaded will not come until the medical profession as a whole take the step of impressing on the community at large the dangers of the infection and the gravity of the results that come from it. The habit has been of making light of the disease, saying it amounts to very little, may cause discomfort a few days or last a few weeks, but that that is all there is of it, whereas the gravity of the results from the conditions are being realized more and more by the medical profession, and should be by them made known to the community at large, so that the influence of the general public may be brought to the stage of realizing the necessity for stamping out the disease as much as possible.

DR. GARCEAU: I would like to call attention to certain cases of gonorrheal cystitis which in my experience have proved to be most difficult to cure by ordinary means. I refer to long-standing cases in which the inflammation has extended into the interstitial tissue of the bladder. Under these conditions irrigations and medicines exert but little action, because they do not come in contact with the seat of the disease. These cases are very apt to be complicated by extension of the inflammation up into the ureters and the pelves of the kidneys, and the irritation caused thereby aggravates the irritability of the bladder and gives rise to the constant urgent desire to micturate. Yet this extension upwards is not always a necessary sequel of cystitis, for in at least one case now under my care the inflammation was confined wholly to the bladder. She was a woman who had contracted gonorrhea about three years ago. This was followed by cystitis, and inflammation of the ovaries and tubes, necessitating the removal of the latter. Her bladder presented on cystoscopic examination the appearances usually seen in cystitis, but there was no ureteritis nor pyelitis, and care was taken to determine the presence or absence of these affections by physical examination and analysis of urine drawn by ureteral catheters from each ureter. She was finally treated by making an artificial vesico-vaginal fistula. The acute forms of

cystitis are very amenable to treatment provided the treatment is instituted at once. This is because the lesions are superficial and affect only the mucous membrane of the bladder. The so-called "subacute forms" are more difficult to treat; while the chronic forms are often very obstinate, even cystotomy with prolonged drainage sometimes fails to cure these patients; for when the artificial fistula is closed, they may be just as badly off as before.

DR. CUSHING: There is only one point I should like to add, and that is as to the question brought up by Dr. Reynolds, whether to operate on these cases of pus-tubes or leave them alone. I would point out some of the points which guide me in making this decision. The first point is as to whether there is any sign of suppuration, whether there is a chill or decided rise of temperature. It may be replied that that is owing to mixed infection, is not a purely gonorrheal affair. That is a mere academic distinction. The concrete case is a woman who is sick, and we want to know whether to operate or not. The fact that there is fever, especially with a chill, is a strong factor in deciding upon operation. The second point is whether the disease is accompanied by a great deal of pain. No one can judge exactly how much pain there is in another person, but my observation has been that women do not submit to operation merely because they are told to do so or on general principles. If it is left to them to decide whether the amount of pain demands operation, and that the dangers are fully explained, they will immediately divide it into two classes, the women who desire and must have operation because they are suffering, and the women who, as they are not suffering very much, will conclude they do not want operation; and that will be found always to correspond with the condition internally, the amount of inflammatory disease, amount of enlargement, and particularly the amount of adhesion and involvement of the intestine. The mere fact that there is a history of gonorrhea, a mass on one side which is not particularly painful, and does not show fever, is not an indication for operation in my judgment. The fact that the woman's usefulness is destroyed is sufficient to warrant operation. The fever, the pain and the involvement of the intestine as shown by difficulty of defecation, pain or obstruction, those are the main points, or the fact that an abscess is forming which requires evacuation of the pus.

DR. REYNOLDS: I quite agree with all that Dr. Cushing has said. I think the bad cases are rather the exception, but when they occur we have got to operate on them of course. I do not think that Dr. Garceau quoted me fairly, nor do I agree with the position that he takes. The quotations that are furnished us by Skeene and other authors are, I think, rather unfair, because the whole subject of the urinary diseases of women has been so revolutionized in the last two or three years, that things in print are mostly matters of the dark ages. I said, or meant to say, that I thought the cystitis was seldom a very troublesome or important affair except when complicated with uretero-pyelitis. It has been my experience, I think I can say without exception, since I have known the modern methods, that whenever a cystitis does not do reasonably well, the ureters, or one ureter and the corresponding pelvis of the kidney are affected, and I think in almost all cases, practically in all cases, you can with sufficient care and topical treatment get a bladder

well, and so get rid of the majority of the symptoms; then you have a long, slow time with the ureter. I think in the end most of these cases get well. If I had a case where the bladder did not get well, I should suspect the ureters. I see nowadays very few cases of simple cystitis that do not get well, and get well pretty promptly.

PATHOLOGICAL SPECIMENS.

DR. CUSHING: I have brought a tumor which I removed yesterday, which is interesting in that it is a tumor of the broad ligament without any pedicle. A woman of fifty-two had noticed something wrong three or four years, but only for a few months had known it was a tumor. In opening the abdomen no adhesions. Two gallons of fluid drawn off. The tumor had no pedicle, but dipped down into the pelvic cavity. The difficulties of removing a tumor of that kind are sometimes so great that I thought it would be interesting to explain the operation from the specimen. The Fallopian tube starting from the uterus is stretched across the tumor, and just below that the sigmoid flexure and lower part of the colon appear to be inserted directly into the tumor. It is not uncommon for errors to be made, and it is a very serious matter to separate the mesocolon by mistake from adhesions, owing to hemorrhage. I cut around the tumor through the peritoneum. This circle goes all the way around the tumor, and by a process of dissection the tumor was drawn out of the pelvis. It was under the peritoneum, and lay over the iliac vessels. I tied the ovarian artery. As the sac was separated from the uterus the latter was left entirely raw on one side. I thought it was best to remove the uterus and did so. In cutting across the uterus what appeared to be antero-posterior was really lateral as the pressure had flattened the uterus laterally. I had then a vast cavity down there, but I was able to bring the peritoneum over the bladder in apposition with the peritoneum running along just below the colon, leaving a glass drainage-tube in the bottom of the wound. The patient has recovered without accident. For comparison I have brought a tumor which I removed ten days ago; it is a regular multilocular tumor, in which you find daughter cysts, and expect to find a pedicle, whereas in the tumor first shown you must be prepared to find no pedicle.

I have also brought a gonorrheal pus-tube removed from a young woman married six months, and for the last three months an invalid. Mass in the pelvis, and getting larger, causing a good deal of pain, and I removed it. I have every reason to think it was gonorrheal. Her husband is a bar-tender. Recovery.

Sunday I did another case, which at first was supposed to be pus-tubes. A young woman, single, but with a lover, had a mass in the pelvis, high temperature, very ill. On opening the abdominal cavity, I soon got into the mass containing dark grumous blood; and letting this out and some large clots, I tried to bring up what I supposed to be a large tube. I found the upper surface of this was formed by the large intestine, of which one could see the appendices epiploicæ; and reaching down behind and separating, I could reach the ovary below. It was a difficult dissection, had to be done partly with the scissors to separate the bowel from the tube. The tube was removed. The other tube was diseased. That left a cavity the size of the fist. It was cleaned as far as possible, and Mikulicz's gauze dressing put in. Recovery.

THE BOSTON
Medical and Surgical Journal.

THURSDAY, SEPTEMBER 2, 1897.

A Journal of Medicine, Surgery, and Allied Sciences, published at Boston, weekly, by the undersigned.

SUBSCRIPTION TERMS: \$5.00 per year, in advance, postage paid, for the United States, Canada and Mexico; \$6.56 per year for all foreign countries belonging to the Postal Union.

All communications for the Editor, and all books for review, should be addressed to the Editor of the Boston Medical and Surgical Journal 283 Washington Street, Boston.

All letters containing business communications, or referring to the publication, subscription, or advertising department of this Journal should be addressed to the undersigned.

Remittances should be made by money-order, draft or registered letter, payable to

DAMRELL & UPHAM,
283 WASHINGTON STREET, BOSTON, MASS.

THE INTERNATIONAL CONGRESS:
THE BRITISH MEDICAL ASSOCIATION.

THE Twelfth International Medical Congress held at Moscow is over, having occupied the week from August 19th to August 26th; and the Sixty-fifth Annual Meeting of the British Medical Association is now in progress at Montreal.

The Secretary-General announced the attendance of members at Moscow as 7,300, of whom nearly 4,000 were from Russia, 800 from Germany, 800 from Austria, 400 from France, 300 from Italy, 300 from England, 120 from the United States, 20 from Mexico, 10 from Japan, 4 from China, and the rest from various parts of Europe, South America and Asia. The most notable of the addresses seems to have been one by Professor Virchow on "The Continuity of Life," in which he harks back to the Cellular Pathology and suggests that in the cell will be found the key, if it ever is found, to unlock the mystery of the ages, and the united labors of the biologist and the pathologist will be the means of the discovery.

The United States was represented among the honorary presidents by Dr. Senn of Chicago, and Dr. Thayer of the Johns Hopkins.

The next Congress will be held at Paris in 1900 during the French Exhibition. Professor Lannelongue, of Paris, has been appointed President of the Committee on Organization, and Professor Chauffard, Honorary Secretary-General. The French representatives were received with marked enthusiasm by the Russians at the Congress, though we see no mention of their being subjected to the demonstrative hugging on the part of the gentle sex which took so acute a form with the sailors accompanying President Faure as to be embarrassing even to those well-seasoned tars,

A telegraphic summary of the first two days of the Congress is given in the *New York Medical Record*. Our special correspondents will furnish presently their impressions of the meeting.