Northrup: Two Cases of Gonococcal Peritonitis in Young Girls.
Archives of Pediatrics, 1918, xxxvi, 475.

The original histories of these two cases had been reported by the writer in 1903. Now, 16 years later, he places the fact on record that both girls had married and one of them borne a child. In regard to this case it proves interesting to recall the following features of the original report: The infection had been acquired from an adult in the family. Onset was exceedingly abrupt, abdominal pain at once very intense. Localized pain and sensitiveness in the right iliac fossa, moderate abdominal distension, ghastly pallor, temperature 104½ F. Six hours later a laparotomy was performed, revealing an intense infection of the entire peritoneum, the cecum being blood red. Appendix was removed; ovaries and tubes showing the same red color were not touched. Prompt recovery.

Bucura: Gonorrhea in Women. Wiener klinische Wochenschrift 1920, xxxiii, 789.

A long article covering the course and treatment of gonorrhea. The gonococcus grows best at the temperature of the human body. It is not resistant to heat as it dies in cultures kept at 36° to 37° C in 13 to 14 hours and at 40° C in 7 hours. It is resistant to lower temperatures as it can be regained from the heart blood 24 hours after death. This is significant in using hot douches and baths in treatment though perhaps the hyperemia is the effective agent.

The gonococcus manufactures a toxin which causes an inflammation resulting in a serous, hemorrhagic, purulent, or mucopurulent exudate. The germ penetrates undamaged epithelium between the cells, eventually reaching the subepithelial connective tissue, leading to endarteritis, thrombosis and metastasis. This fact explains the slight benefit from a direct application of drugs.

The gonococcus attacks most easily cylindrical epithelium, less easily pavement epithelium. The urethra, ducts of the vulval and urethral glands, cervical canal, uterine cavity, and tubes are specially prone to become infected; the vulva and vagina only at the periods when their epithelium is soft and succulent, i.e. before puberty, during pregnancy, and in old age.

Some individuals and families seem to have an immunity against infection and extension of the process.

The duration of the disease is not known. Persistence for ten years and more might be due to reinfections. It is known that the germ dies quickly in closed cavities such as a pyosalpinx. The author has seen the ascension of the infection to the tubes with simultaneous healing in the cervix and uterus which he thinks may be due to the development of an antitoxin.

The diagnosis is simple when the germ is seen. The chronic cases require a painstaking examination of the discharge from urethra, vulval glands, and crypts, vagina, cervix, and uterine cavity, repeated many times. The author insists upon the necessity of finding the gonococcus for the positive diagnosis because the disease has no typical clinical symptoms and may be confused with any gynecologic disease causing discharge.

The gonococcal etiology of adnexal disease is established only by discovering the germ. The wrong diagnosis is made too often. If prostitutes are left out of consideration, less than 50% of adnexal inflammations are due to the gonococcus.

The treatment consists of rest and abstention from coitus, the use of vaccines, and local applications. Vaccines are used in all cases because they seem to lessen extension of the trouble to the uterus and tubes. A polyvalent vaccine is used at intervals of 3 to 8 days beginning with 5 million, then 25, 50, and 100; then 100 repeated to a total in all of 10 to 12 doses.

There is no specific drug for local application. Silver nitrate, corrosive sublimate, and carbolic acid coagulate the exudate and prevent further penetration of the agent. Protargol, collargol, argentamin, and methylene blue in 1 per cent or stronger solution are the best. The urethritis rarely needs treatment, it gets well if let alone. The treatment of the urethral and vulval glands and crypts is most difficult and may require excision. The vagina is seldom infected. Tampons have no place in the treatment, but douches at least wash out the discharge. In treating the cervix the exudate is removed and the drug applied to a clean surface.

Infection of the uterine cavity frequently causes tubal infection. The latter will not heal until the uterine infection is cured. The uterine cavity is treated by the injection of a few drops of any of the above drugs through a ureteral catheter introduced with the strictest regard for asepsis.

Acute infection of the tubes is treated symptomatically. When the acute symptoms subside vaccines and local treatment are given. The latter consists of douches, tampons, hot air, and later on intrauterine treatment. Operation should not be done in the subacute stage because of the danger of peritonitis, of infection of the wound, and phlebitis. Operation in the chronic stage is determined by the symptoms and such social indications as marriage or sterility.

Definite cure is established as follows: During treatment discharge from the several sites of infection is examined once or twice a month; when clinical symptoms have become stationary or have disappeared for several weeks after the last finding of the gonococcus treatment is stopped; then the patient is seen once or twice a month for three months. She leads a normal life and is allowed coitus with a condom. Postmenstrually secretions from the various sites are examined, if no gonococci are found during these three months Bucura considers the patient as cured.

FRANK A. PEMBERTON.

Stevens and Heppner: Gonorrhea of the Lower Genito-Urinary Tract in Women, with Special Reference to the Glands of Bartholin.
Journal American Medical Association, 1920, lxxv, 1477.

This study is based on 3,439 examinations of prostitutes detained at the San Francisco County Hospital. Chronic gonorrheal infection was determined in 43.5 per cent. At first the diagnosis was based on clinical findings plus either the detection of the organisms in smears or a ++ or +++ positive complement-fixation test, the fallibility of both of these methods being conceded. Later, the diagnosis was based on clinical findings alone, yet in 95 per cent of cases so diagnosed, Gram-