of the medicine in the system in sufficient quantity to be dangerous is hardly possible in that strength.

A medicinal agent to be absorbed by the unbroken skin, must either become volatile during its application, or be incorporated in a fatty base and applied with friction. Occlusion is an absolute requirement in the case of volatile substances. Most of the absorption occurs in the crypts of the sebaceous glands. The effect is much slower, about one-fourth as intense, and longer continued than when the same agent is administered by mouth. A large number of medicinal agents, which when given by mouth are intolerable to many patients, may be administered in this way with scarcely any unpleasant effects.

BIBLIOGRAPHY.

- Arendt: *Ann. et Bull. de la Soc. de Med. d'Auvers*, April, 1898.
 Aubert: *Lyon Méd.*, cxv, 1894.
 Bourget: *Thérapeutische Monatshefte*, 1893, p. 531.
 Bartholow: *Text-Book Therapeutics*, 1899.
 Cullen: *THE JOURNAL*, 1898, p. 1218.
 Credé: *Klinische Therap. Wochenschr.*, 1898, Nos. 14 and 15.
 Cushny: *Text-Book, Pharmacol. and Therapeutics*, 1899.
 Davis, N. S.: *Hare's Practical Therapeutics*, iv.
 Destot: *Lyon Méd.*, Dec., 1894, and p. 81, 1895.
 Fitzgerald: *British Med. Jour.*, July 15, 1899.
 Foster: *System Therapeutics*, vol. i, p. 459.
 Fornaca: *Gaz. degli Osp.*, Dec. 25, 1898.
 Flick, L. F.: *Therapeutic Gazette*, Jan. 15, 1900.
 Friedländer: *Berlin. Klin. Wochenschr.*, 1898, xxxv, p. 470.
 Fürbinger: *Virchow's Archiv*, Bd. lxxii.
 Guinard: *Soc. de Therap. et Pharmacologie*, i, 1896.
 Guttman, P.: *Archiv f. Klin. Medic.*, Sept. 15, 1886.
 Gallard, F.: *Cit. by N. Y. Med. Jour.*, p. 56, 1899.
 Gottheil: *Pharmaceutical Era*, July 1, 1894.
 Inglis, D.: *Physician and Surgeon*, 1897, xix, p. 436.
 Juhl: *Cit. by Guinard*.
 Jones: *Med. Record*, Feb. 11, 1899. (*Transact. N. Y. Acad. Med.*)
 Josephs: *British Med. Jour.*, Dec. 30, 1899.
 Klebs: *N. Y. Med. Monatsch.*, xx.
 LeMoine: *Bull. et mem. de la soc. des Hospitiaux*, No. 17, 1897.
 Liebreich: *Deutsch Med. Wochenschr.*, No. 4, 1885.
 Lannois and Linnossier: *Bull. gén. de Therapie*, 1896: *Mem. et Compt. Rend. Soc. de Med. Lyon*, 1898, xxxvii, p. 119. *Soc. de Biol.*, March, 1894.
 Mackey: *Transactions Brit. Med. Assn.*, *Lancet*, 1886.
 Marx: *Cit. by Jones*.
 Osborn: *Physician and Surgeon*, July, 1898.
 Pribram, A.: *Der Acut. Gelenkrheumatismus*, Nothnagel's "System."
 Pavolsky: *Rushxaia Medizina*, 1886, 12, p. 307.
 Renaud: *Soc. de Therapeut.*, 1894.
 Ruel: *Cit. by Guinard*.
 Shriner, G.: *N. Y. Med. Monatsch.*, 1898, x, No. 11.
 Smith: *Am. Jour. Pharmacy*, 1898, pp. 70-182.
 Shoemaker: *Diseases of the Skin*.
 Sciolla: *Semaine Méd.*, April 8, 1893.
 Unna: *Thérapeutische Monatshefte*, 1890, Nos. 2 and 4.
 Weiss: *Wiener Med. Presse*, G. 48, 1894.
 Winternitz: *Archiv f. Exp. Pathol. u. Pharmacol.*, 1891, xxxviii.
 Waller: *Cit. by Bartholow*.
 Vanni and Guicardi: *Cit. by Guinard*.
 Von Zeissel: *Wien. Med. Presse*, 1898, xxxix, p. 569.
 Vidal: *Cit. by Therapeutic Gazette*, 1898, p. 466.
 314 West One Hundred and Forty-first Street.

IDIOSYNCRASY AS TO MERCURY.

A CASE OF ERYTHEMA MERCURIALE.*

ALBERT BERNHEIM, M.D.

PHILADELPHIA.

In the history of many a great man, we are told that he had a peculiar aversion to some special thing: be this a cat or a rooster, or a flower or some perfume. In some cases of convalescence, we find special aversion to smoke, even in persons who have been addicted to smoking. Such aversions may be temporary or constant. If the aversion is temporary, we should call it a disposition in consequence of some disturbance in individuality; but if it is constant we may call it a peculiar constitution of individuality, in consequence of

which particular agents operate on the individual in a manner different from their ordinary mode of action on persons in general; briefly, we may call it an idiosyncrasy. Even animals are reported to have idiosyncrasies, thus the horse to the camel.

Idiosyncrasies are not uncommon; they occur in regard to various agents. Perhaps the most frequent ones are those in relation to some foods, such as strawberries, and certain fish. Others are connected with drugs and medicines. A well-known German professor of surgery could not stand iodoform; some can not use carbolic acid; others can not take antipyrin without becoming subjected to an eruption of blisters on many parts of the body.

In the literature on the use of mercury, we do not find so frequent an idiosyncrasy to this drug, and extraordinary effects of it, that it would not be worth while to mention such effects when they do occur, especially when such symptoms incline to simulate phenomena of syphilis, erysipelas and scarlet fever.

If we look over the literature of the 19th century, we do not meet many cases of untoward effects of mercury compared with its numberless uses.

One of the first who wrote about the unexpected effects of mercury was G. Alley,¹ of Dublin, who published, in 1804, in a large treatise, several cases of a peculiar eruption arising from mercury. About the same time, McMullen,² in Edinburgh, described similar cases; Spens³ followed, in 1805, with a description of three cases of erythema mercuriale. They considered the correctness of the name—eczema or erythema. Others again called it erysipelas, and as late as 1887 Dr Kreidmann⁴ discarded the bacteriological origin of erysipelas in favor of medicinal causes, such as mercury.

In 1811, A. Ramsay,⁵ and in 1812 J. Frank⁶ and J. Nicholson⁷ related cases of mercurial erythema. H. Becker⁸ published a treatise, "De Erysepelate Mercuriali," in 1817; B. Kahleis⁹ spoke of mercurial rose in 1823; in 1845 Rambant¹⁰ called attention to purpura hemorrhagica caused by mercury; and Larmande¹¹ wrote about a hydrargyro-atrophic erythema in 1880.

S. Snell¹² described in 1882 a peculiar idiosyncrasy as to mercury. Alexander¹³ related in 1884 a case of acute universal mercurial eczema; a year later Descroizilles¹⁴ wrote on mercurial eczema. P. Gaucherand¹⁵ wrote in 1886 about scarlatiniform skin eruptions caused by internal administration of mercury; Blanc¹⁶ and Kreidmann, in the following year, make observations on mercurial scarlatiniform erythema.

In more recent years, Robinson¹⁷ speaks of an individual who was suffering from two attacks of erythema, one after the other; the first after mercurial inunction, and the second after internal use of calomel. G. Lewin¹⁸ exhibits a case of toxic exanthema caused by quicksilver; and Fordyce¹⁹ mentions a case of desquamative exanthema that had been diagnosed as scarlatina.

Mrs. V. G., 50 years of age, mother of seven children, felt intense itching on her hands and arms in the morning of January 20; the following day she had a rash on her throat, flexor sides of her arms, palms of the hands and flexor sides of her legs; the eruption was the most marked in the hollows of her knees, flexor sides of wrist and elbows. An extensive redness was visible all over the skin of these said parts; the rash consisted of dark-red, small elevations over the level of the skin and felt, to the touching hand, like velvet; the elevations corresponded to the hair-follicles; on January 22 the skin was edematous and hardened like in erysipelas; fever did not exist; urine was abundant, and of a water-

* Presented to the Section on Materia Medica, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

like color; no albumin; no sugar. Diagnosis was uncertain. I remembered then that patient had told me of eruptions she had had in 1869, 1875 and 1882, and that she had an idiosyncrasy to the use of any kind of ointment. The eruption appeared to me as a symptom of poisoning.

The very intelligent patient gave me the following history of the previous cases of rash: In October, 1869, she consulted a physician for removal of a red spot of the size of a silver dollar, on her neck. The spot was a reminder of a vesicatory put there when as a little child she was believed to suffer from meningitis. The physician prescribed for this spot a grayish ointment. After a single inunction she got an itching rash over her throat, chest, abdomen, flexor sides of arms and legs, with subsequent furunculosis on several parts of the body, particularly the arm-pits. In 1873 the patient married; in October, 1875, after a pregnancy, the hair began to fall out; she used a hair pomade; after its use for the first time she got an eruption on her throat, hands, neck and back of the head. In May, 1882, after another pregnancy, she was suffering from constipation. The same physician who had treated her in 1869 prescribed a powder of grayish color, and greasy, shining appearance. A few hours after taking the first powder a rash all over the body—throat, chest, flexor sides of arms and legs, hands—occurred, followed by furunculosis again. The new attack of eruption seemed to her to resemble the previous ones.

I learned that the patient had taken on the evening of January 19, two pills, and on January 20, three pills, as a laxative, and on closer investigation I found out that the pills were blue-mass pills of 2 grains each; the diagnosis then was easy; it was a case of effects from mercury.

In the previous times, as well as in the new attack, the skin scaled off in large lamellæ, like in scarlet fever, except that the desquamative process was to a thicker extent. The teeth were greatly affected in the attack of 1882, not so in this eruption. Mercurial diarrhea did not occur, there was rather an inclination to constipation.

The urine was examined for the presence of mercury, according to Almen. I used for the purpose a very fine copper wire, such as taken for induction coils of faradic instruments. The examination was positive.

Patient comes from a healthy family; none of her sisters or brothers is known to be subjected to the same idiosyncrasy, except that father and one sister in younger years could not eat strawberries without getting a rash; later on they were not affected any by eating the berries. The children of the patient are not known to suffer from the same disposition, for one or the other has taken the same blue-mass pill or calomel.

The treatment consisted of administration of two tablespoonfuls of castor-oil and magnesium sulphate; of hot baths, diaphoresis, drinking of egg albumin stirred in boiled and cooled water, of large quantities of hot milk and hot water. The itching was first treated with inunction of vaselin and dusting of borated talcum powder, but with not a great success; it stopped promptly on administering potassium iodid, 8 grains in 150 grains of water three times a day; a second examination of urine did not exhibit mercury any longer. A struma that had existed for several years subsided simultaneously with the administration of potassium iodid.

BIBLIOGRAPHY.

1. G. Alley: An essay on a peculiar eruptive disease arising from the exhibition of mercury. Dublin, 1804.
2. J. McMullen: De erythemate mercuriali. Edinburgh, 1805, and Edinburgh Medical and Surgical Journal, 1806, ii, 25-37.

3. Spens: Three cases of erythema mercuriale. Edin. Med. and Surg. Journ., 1805, i, 7-20.
4. Kreidmann: Hydragryum Erysipelas, Allgemeine Med. Zeitung. Berlin, 1887, lvi, 957.
5. A. Ramsay: Case of erythema mercuriale. Edin. Med. and Surg. Journ., 1811, vii, 269-274.
6. J. Frank: Erythema mercuriale. Acta Instit. clinic. caes. Universitatis Vilmensis, Lipsiæ, 1812, 22-27.
7. J. Nicholson: Case of erythema mercuriale, accompanied by an affection of the cornea. Edin. Med. and Surg. Journ., 1812, viii, 39-41.
8. H. Becker: De erysipellate mercuriali. Duisburgi, 1817.
9. B. Kahleis: Ueber die Mercurialrose (Erythema mercuriale). Journal d. prakt. Heilkunde, Berlin, 1823, lvi, 49-68.
10. Rambant: Case of purpura hemorrhagica caused by Mercury. Dublin Hosp. Gazette, 1845, i, 18-20.
11. Larmande: Un erytheme hydrargyro-atrophique. J. de Med. and Chir. Prat., Paris, 1880, li, 497-499.
12. S. Snell: A peculiar idiosyncrasy (induction of pyrexia) as to mercury. Practitioner, London, 1882, xxix, 18-21.
13. Alexander: Ein Fall von acutem universellen Mercurialeczem. Vierteljahrsschrift f. Dermatologie, Wien, 1884, xi, 105-110.
14. Descroizilles: Eczema hydrargyrique; coïncidant avec une stomatite de même nature à la suite des frictions faites avec l'onguent napolitain. France Méd., Paris, 1885, i, 447-451.
15. P. Gaucherand: Des eruptions cutanées causées par l'administration interne du mercure (et en particulier de la forme scarlatineuse). Paris, 1886.
16. Blanc: Observation d'erytheme mercuriel scarlatiniforme. Provence Méd., Lyon, 1887, ii, 393-396.
17. Robinson: Med. Analectic and Epit., August, 1890, quoted in Sajous' Annual, 1899, iv.
18. G. Lewin: Deutsche Med. Wochenschrift, 1894, S. 259.
19. Fordyce: Journal Cutan. and Genito-Urin. Diseases, Dec., 1895, in Sajous' Annual, i, c.

A NEW OBJECTIVE TEST FOR MASTOIDITIS.

WITH REPORT OF CASE.

ALBERT H. ANDREWS, M.D.

CHICAGO.

The use of the tuning-fork and the stethoscope to determine the comparative density of the two mastoids furnishes a valuable aid, and so far as I am aware, a new method in the diagnosis of mastoid complications. The test is made by placing a stethoscope with a small bell over the tip of the mastoid, and placing the handle of a vibrating tuning-fork over the antrum. It is found that when the mastoid is filled with pus or granulations or when it is dense from obliteration of the air-cells the sound waves are transmitted to the ears of the examiner more distinctly than when the stethoscope and tuning-fork are placed in the same relative position over the opposite or normal mastoid. Care must be used, in making the test, not to stretch the skin between the stethoscope and the handle of the tuning-fork, for then the vibrations are heard better than when the skin is not stretched, even though the distance be less.

The lower border of the middle fossa can be very accurately located by the same method. While the bell of the stethoscope is upon the mastoid the vibrating fork is moved downward over the squama and as soon as the upper border of the mastoid is reached the sound becomes much louder. The line indicating the change from cranial cavity to mastoid is usually well defined. When this line is reached a move of an eighth of an inch makes a marked difference in the sound of the fork. The position of the lateral sinus in relation to the mastoid can be determined in the same manner, though with less accuracy.

Since beginning the use of this test I have examined forty cases without mastoid symptoms and have found no perceptible difference in the resonance of the two sides; while in the examination of four cases of undoubted mastoid disease the affected mastoid transmitted the sound waves to the stethoscope with greater force than did the mastoid of the opposite side. One of the mastoid cases was recently operated on and the following description gives the chief points: