Kernig sign, there were no signs of meningitis. There were, however, a slight erythema and a tache cérébrale, often seen in tuberculous meningitis, over the face and body. The bowels were open. From now on, for the next three days, there was a slight headache, some constipation and irregular pulse, after which the patient was in comparatively good health.

February 28, he experienced severe abdominal pain and vomiting. The temperature was 102, and the pulse 110, being irregular. The abdomen was very rigid and tender, as in peritonitis. These symptoms and signs persisted twenty hours, after which they subsided. The urine contained no sugar nor fat. There were no other metabolic changes present to confirm the suspicion of an acute pancreatitis. From now on the patient made an uneventful recovery. The report on the spinal fluid, made by Dr. Josephine B. Neal of the laboratory of the department of health of New York City, was: Ten c.c. opalescent fluid. Cytology, 99 per cent. Mononuclears, great increase. Bacteriology: Spread apparently negative. Culture, negative. Albumin ++ 1. Globulin ++ 1, Fehling's ++ +.

CASE 2.—H. S., a girl, aged 9 years, was bright, nervous and undersized. Her parents were of nervous temperament. She was fed by bottle from the first month. She had had measles three years previously.

February 24, she had a mild parotitis on the left side. The temperature was 101 F. The parotid gland was normal after two days. There was no fever after the second day.

February 28, intense prostration occurred with headache, vomiting and constipation of twelve hours' duration. There was a rigidity of the neck and spine, also a scaphoid abdomen and a slight Kernig sign. The temperature was 99, the pulse was 110 and irregular, and the respiration 20.

These symptoms and signs were present for the next four days, progressively improving. The gastro-intestinal symptoms disappeared first. The irregular pulse persisted for three weeks after the first appearance of the meningitis.

CASE 3.—M. S., a girl aged 5 years, was bright and undersized. The parents were neurotic. She had been fed by bottle after the fourth month. She had had measles two years previous to the date of consultation, and lobar pneumonia with meningismus eight months previous.

March 5, 1918, a bilateral parotitis, moderately severe, appeared. The temperature was 102.

March 9, a severe headache set in, with vomiting and constipation, marked irritability and insomnia. The temperature, pulse and respiration were normal. The abdomen was retracted. There was a slight rigidity of the neck, with increased knee jerks, and a left Kernig sign was present.

The following day, the symptoms and signs were the same, but there was greater prostration. From now on, the patient progressively improved, except during a recurrence of the parotitis on both sides, March 11.

COMMENT

The last two cases were relatively mild and would ordinarily have passed as gastro-intestinal attacks. The absence of fever and the marked prostration should, however, make one suspicious. Meningeal irritation, or meningismus, is to be thought of in the milder cases even though the necropsies have demonstrated definite inflammation in the severer forms. In Case 1, there was no doubt of the diagnosis from the start, its course being very different from that of the other forms of meningitis. The value of early spinal puncture cannot be too much emphasized in this type of case. It was not indicated as a therapeutic measure in Cases 2 and 3, in which there were no signs of cerebral pressure.

165 Audubon Avenue.

Importance of Health.—With a greater degree of health among all, religion will become more effective for good, mortality will have a deeper significance and a wider application, and knowledge will multiply and distribute its blessings more widely.—Vaughan.

THE ANAMNESIS IN CARDIOVASCULAR DISEASE

WILLIAM BRADY, M.D.

ELMIRA, N. Y.

Cardiovascular disease is now the chief cause of death in the registration area, which covers slightly more than two thirds of the population of the United States. Cardiovascular degeneration, for the most part, is an insidious condition, far more difficult to diagnose in the incipient stages than is tuberculosis, but far more prevalent than tuberculosis and more frequently unrecognized by the practitioner.

As a result, I believe, of the great popularity of nostrums in this country, the millions in capital exploiting nostrums, and the vast propaganda of the nostrum interests that work indefatigably to misinform and mislead the lay public through every available avenue of publicity open to dishonest advertising, the great majority of victims of cardiovascular disease do not consult a physician until some alarming manifestation of the disease drives them to his office. And even then, too often, either the patient is averse to a painstaking examination or the physician is too busy, in his way, to make such an examination, so that the condition drifts on for months and years undiagnosed, nay, unsuspected, until disaster comes and realization bursts on the persons concerned. This is the explanation of the popular notion, a notion sometimes encouraged by physicians, that Bright's disease is generally fatal in a year or two, that apoplexy comes like lightning from a clear sky, that heart failure is a sudden calamity that occurs without warning. This is also the explanation for the shameful abuse of such makeshift terms as "debility," "overwork," "run down condition," "nervous exhaustion" and "neurasthenia" in lieu of a diagnosis of the cardiovascular degeneration of middle age.

It is a well recognized fact that a competent physician can and must make a tentative diagnosis of incipient pulmonary tuberculosis, and institute the proper regimen and treatment, long before bacilli can be recovered from the sputum and long before any serious breakdown of health has occurred. Granted that even the most skilful diagnostician occasionally errs in this tentative diagnosis, no great harm has been done the patient in any case, since the usual treatment employed for incipient tuberculosis can do no harm if the patient happens to have, say, masked exophthalmic goiter instead, and, indeed, may be precisely the treatment required by such disease. We all know that the best of diagnosticians is bound to make such mistakes; one with mediocre ability must confess such mistakes. But does this fact militate against the wisdom of a tentative diagnosis of tuberculosis? Or does it weaken the physician's morale in his procedure in future cases of similar character? Not if he is a good doctor. The only man who never makes a grave diagnostic error is the man who never makes a timely diagnosis.

The average well trained physician nowadays has the ability to diagnose cardiovascular degeneration long before Bright's disease, heart muscle failure, apoplexy or general arteriosclerosis overtakes the patient. He has the necessary instruments of precision to corroborate his tentative opinion. But he lacks only the courage to be blunt. He fears to frighten his patient.

For years the same phobia handicapped good men in the diagnosis of incipient tuberculosis. It led them to rather sorry lengths. They fell into the bad habit of telling patients that there was "a weak spot" in one lung, "a slight catarrhal condition" in the apex, or "a few râles" audible by auscultation. They frightened the patient but withheld a definite opinion; in short, they quibbled. And quibbling only led to loss of confidence in the physician sooner or later.

Today we are passing through a similar era in our effort to grasp the tremendous problem of cardiovascular degeneration. We are quibbling with the patient. We are diagnosing "high blood pressure," "functional heart trouble" or "autointoxication." Patients with cardiovascular disease are drifting along from month to month, from year to year, in ignorance of their condition, and, what is worse, seeking relief wherever a plausible promise is held out — and plausibility is the hallmark of the modern charlatan.

What does a physician mean when he assures a patient the "trouble is only functional?" The truth is he does not exactly know what he means. He is quibbling with himself. Disease is altered function; if every organ is normal there will be no altered function. Then how can there be any distinction between "organic" and "functional" disease? Every organic disease is functional; if there were no functional disturbances there would be no symptoms. Every functional disease is organic, only our knowledge is sometimes not sufficient to make the diagnosis, or our opportunity is limited by circumstances beyond our control. The patient either will not submit to necessary observation and study, or cannot afford to pay for them.

The anamnesis of cardiovascular degeneration is written in the testimonials published by the nostrum makers, in the news columns of the daily papers, and in scattered, sketchy notes in the practitioner's case records. It is a puzzle picture so cut up and confused in medical textbooks that one succeeds in piecing it together into a significant whole only after some years of general practice. Perhaps the reason the picture is commonly overlooked is that it requires years to complete it in an individual case. And again, the patient is prone to lose interest in the effort and go off to sit for another artist. It is a new idea, this cardiovascular degeneration. The very term has an unpleasant sound. Degeneration, to the lay mind, suggests degenerate, and degenerate suggests something revolting. Deterioration is no better, for the victim in the last stage resents deteriorating. Had our friends the fishwives not already spoiled it, the change of life might meet the requirement; but one would be taking some risk telling a man that he was going through the change. General arteritis might suffice, unless some one has a more acceptable term.

Bearing in mind that cardiovascular degeneration is the most widespread disease of persons over 40 years of age and that it occurs probably ten times as frequently as cancer, one must think of it as a not improbable cause of the general ill health of patients of middle age who seek medical advice after a prolonged trial of the various nostrums and plausible remedies so popular with the public.

The man or woman of 40 or more who comes complaining of loss of strength, falling off in efficiency both physical and mental, gradual accumulation of superfluous weight, mild dyspnea on exertion, "stomach trouble" or "gas" not apparently due to abdominal lesion, restless or unrefreshing sleep, "all tired out,"

"run down condition," fatigue headaches, "poor circulation" or "nervousness" certainly requires a careful study of the cardiovascular system. A symptom that has struck me as rather suggestive in the anamnesis is failure of sexual competence in a middle-aged man not suffering with any discoverable genital lesion.

When repeated measurements at short intervals and at varying periods of the day show a constant elevation of the systolic pressure above 140 mm. of mercury, a diagnosis of cardiovascular disease should be considered feasible until definitely ruled out by rigid correction of the patient's habits.

Inability of the patient to hold the breath for at least thirty seconds without preliminary hyperpnea should raise the suspicion of cardiovascular disease.

A pulse rate remaining more rapid than normal for over three minutes after fifteen deep knee flexions should also arouse a suspicion of the cardiovascular integrity, be the patient a neurasthenic or not. "Neurasthenia" developing after 40 is in itself suggestive of cardiovascular degeneration, irrespective of alleged overwork or worry.

The full, florid or tumid face, with slight exophthalmos, in a person over 35 who has accumulated excessive weight in recent years and boasts of a hearty appetite, though troubled with "biliousness" or fullness after meals (a high stomach, some patients call it), is very likely a mark of cardiovascular disease.

It is the practice of physicians to treat acute respiratory infections, wrongly called "colds" and "grip," in a routine way, the expectant way, without attempting any great refinements of diagnosis. And this practice has proved reasonably satisfactory. But it has long been the practice to insist on great refinement of the diagnosis in cardiovascular disease, although the routine treatment that proves most successful is very much the same whether the individual case be considered myocarditis, arteriosclerosis, nephritis or apoplexy. Diet, medication, physical therapy, rest, exercise are all managed according to individual indications, and not according to the particular expression of the degeneration in a given instance. The administration of digitalis, for example, is not indicated by the expression of the degeneration, but by the patient's general condition as determined by examination. The same holds true of most remedial measures employed in cardiovascular disease. In other words, treatment cannot be intelligently prescribed for the disease, but must be suited to the needs of the patient.

Although the fine art of prognosis seems ever more difficult and unattainable in general practice, one harbors the idea that the best prognosis rests on an intimate knowledge of the patient's general condition rather than on any fine spun anatomic diagnosis of the specific area of his vascular system which is most injured.

In view of the common incidence of chronic cardiovascular degeneration as compared with that of tuberculosis, a diagnosis of the condition in its incipency should be more frequently made than the diagnosis of incipient tuberculosis, and may be tentative as is that of the latter, with only benefit to the patient.

If physicians were in the habit of making such a diagnosis as a routine in daily practice, many millions of dollars now squandered by the laity for worthless and fraudulent nostrums and countless plausible brands of treatment exploited by irresponsible healers would probably be diverted to the useful work of preserving efficiency and prolonging life.