termed nerve force, but which will also account for its conduction in the neuron, and its transmission from neuron to neuron. To do this I have sought to reduce to their simplest conceptions the physical and physiologic conditions present in the structure of the essential elements of the nervous system, and to state what appears to me to be the most probable physical results of such conditions consistent with the phenomena we observe. In recapitulation: The nutritive processes of the neuron, presided over by the nucleus, maintain the protoplasmic contents in that state of readiness for action which constitutes the primary or fundamental stage of all nerve function. This creates in the neuron a condition of surface tension and of static electric potential, and is perhaps accompanied by some protoplasmic or ameboid movement.

Each neuron in this state or condition has a restraining or balancing influence while at rest, upon all other neurons within its field of influence by reason of the energy exerted by it in the form of what is known as electro-magnetic induction.

The end-organs of the sensory nerves serve as avenues of ingress for such forms of motion as are attuned to their mechanism, and this added motion serves to create a molecular rearrangement of the nature of a chemic change in the protoplasm of the neuron, and this is accompanied by a change in the static potential.

A change in the static potential is inseparable from a change in the electro-magnetic field surrounding the neuron, and this variation of the electro-magnetic field acts as a stimulus upon every neuron associated with the one whose end-organ has been excited.

Thus conduction and transference of nerve force are, according to this theory, to be regarded as of the nature of electro-static phenomena.

DISCUSSION.

Dr. COE of Oregon—As I came in a friend of mine in another Section wanted to know where the Section on Neurology was to hold its meetings. I told him in the Baptist church. He said, "I think if there is any Section that ought to meet in the Baptist church it is the Neurological Section, because if any one gets into deep water it is the neurologists;" and I think that this is entirely too deep for me to discuss at this time.

Dr. D. R. BROWER of Chicago—I am not equal to the discussion of this important question. Some of the statements in the paper are facts, and these we all admit. As to the theories therein, they are plausible and reasonable theories, and I thank Dr. Herdman for the wonderful suggestiveness of his paper.

Dr. SEARCY—I have been much interested in the paper on the dynamics of the nervous system. It is one of the opening fields of medicine, and psychologic medicine especially. The unit of life is the cell, and the unit of the nervous system is the nerve cell. We are learning probably more of the action of the nerve-cell in the neuron of the cortex than in any other part of the nervous system, because we find those cells more isolated and more easy of access in their structure, their function and their site. As worked out of late, possibly we can say the dynamics of the neuron consist of the received motions, and adjusting its emitted motion to those received—which is a vital phenomena. The way I have criticized nearly all philosophic theories along this line is that we attempt to describe animate phenomena by inanimate phenomena; for instance, describing the functions of the neuron by a reference to electricity. We would simplify our work if we would describe vital motion by simpler vital motion, describing, for instance, the extremely complex nervous system of man by studying the movements in the lower organisms, in ameboids and protozoa, showing the movements in their simpler forms. We can grade up the vital phenomena from that basis much better than we can by attempting to explain them by inanimate phenomena.

Dr. ADAM A. FORD of Denver—One suggestion came to me in listening to this very able paper, and that is with regard to the ultimate action of the molecule, or rather the atom in the molecule, affecting the neuron extremity. The paper indicated

a variety of different actions due to vibratory action up to a number of trillions. Where can we differentiate chemic action from other action? Can we not trace it to the various forms of vorticular action in the action of these ultimate atoms? I believe it is one of the theories of scientists, now accepted, that this co-ordinated form of action in the atom is vorticular action, and owing to the different formation of this vortex, together with its affinities, we have certain manifestations of power, of force. Might not the author of this paper explain this theory, so as to include vorticular action?

Dr. MEYER—I should like to ask Dr. Herdman with regard to the conduction time. The conduction time has been used in a great many of the du Beromo phenomena in nerve elements. It seems, if I may trust my memory of physics, that the difficulty of meeting the objections rising from the conduction time become rather greater in the theory suggested, than even in the idea that was originally held by du Beromo.

Dr. PETERSON of New York—The author has given a great deal of thought to this subject, and the critic of his paper should at least give as much time as he has to it. The paper is purely speculative, but speculative papers, I think, always have a certain charm. The theory advanced, in my opinion, is as reasonable a theory as ever given in neural dynamics. If I understood Dr. Herdman rightly, it was based upon the installation and independence of the neuron. That is by no means proved, and the most recent researches would tend to show that it is probably untrue. However, theories like these are the tentacles of the scientific octopus, reaching out in every direction for truths, and therefore, are extremely useful.

Dr. W. J. HERDMAN—While, as has been said, such papers as this are largely speculative, my attempt has been to give a solid foundation of fact for the building up of those speculative ideas. The question of vitality has been raised. It is, as we all know, a very much debated question among physiologists at the present time as to whether all manifestations of vitality are not physical and chemic; that molecular arrangements have simply reached a certain stage when these are the natural outcome of the molecular and anatomic arrangement; and I must say that, from the recent investigations I have made on that subject, covering the last five years, the evidence is largely on the side of the man who holds what is called the mechanic view. It is with relation to that view, and to making it bear on the side of neural dynamics, that this paper has been prepared. I have not yet subjected the theory to the tests of nerve condition, as has been asked by Dr. Meyer. I have not conceived how it could be, or how the static condition of an individual neuron could be subjected to the test which has been made of nerve conductivity in the physiologic laboratory method, but some device, perhaps, can be arranged for bringing that under observation as well. It has seemed to me that we must take into consideration what we know in physics and physiology, and bring this knowledge together, and see how the two will dovetail.

UREMIC FOCAL LESIONS.

Presented to the Section on Neurology and Medical Jurisprudence, at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June 7-10, 1898.

BY H. H. HOPPE, M.D.

Lecturer on Nervous Diseases, University of Cincinnati Medical Department; Neurologist to Cincinnati Hospital.
CINCINNATI, OHIO.

The literature on localized lesions of the central nervous system occurring as a result of uremic attacks is very scanty. A diligent perusal of many American text-books on general internal medicine, as well as the text-books on nervous diseases, fails to find mention of the subject at all.

It is true that compared with the multifiform manifestations of uremia, especially compared with the frequency of convulsive attacks, the paralytic attacks occur very rarely; or, perhaps, it were better to say, are rarely recorded, and this is probably due to the peculiar character of the paralytic attacks and to the rapidity with which the manifestations disappear. The study becomes all the more interesting because of the light which the subject seems to throw upon the nature of other central disturbances of the nervous system. If the advance in our knowledge of medicine has been marked in the past quarter of a

century in any one particular direction, it is in that of the somatic origin of the various forms of insanity and of the ordinary diseases of the nervous system, in other words, that the intellect or the soul itself is never affected, but that its disordered manifestations are due to a diseased condition of the media through which it acts. The first successes, among which we may mention Mendel's classic demonstration of the pathologic conditions underlying dementia paralytica, were followed by countless endeavors in all parts of the world to establish the relation between diseased conditions of the brain and the disordered manifestations of cerebral activity. The results, as regards cerebral localization and mapping out of areas of the cortex, with their distinct functions, both sensorial and motor, have, as you know, been crowned with brilliant success. It, however, had a tendency to narrowing our conception of nervous diseases, by referring, or trying to refer, all disordered conditions of the nervous system, psychical, sensorial and motor, primarily to the brain or spinal cord. We know today that the anatomic investigation of the majority of insanities and of all the so-called functional diseases of the nervous system has met only with signal failure.

It may be held that this failure is due to the inefficiency of our physical and chemic means of investigation, but the probabilities are that we will have to go back to the old clinical methods of investigation: to study the human body in life as well as the organs after death. To look upon sickness, no matter what its character, may be a manifestation of a disordered condition of the whole body rather than a disorder of the function of one particular organ. In the light of recent investigations, especially of the functional diseases, we know that the brain and spinal cord are often more sinned against than sinning, and that a remedy for the relief of the functional affections is more often to be sought in correcting a disordered condition of blood, stomach, intestines, thyroid gland and kidneys, rather than of the brain. Bouchard, in his well-known publications on auto-intoxication, started the pendulum to swing away from anatomic investigations of the brain in functional diseases to the study of diseased manifestations of the more distant organs of the body. More beautiful and more classic disturbances of the functions of the central nervous system can not be found than the investigation of the nervous arrangements found in uremic intoxications.

I have purposely waited a very long time before publishing these cases, because I wished to be sure of the diagnosis. We know only too well how our most interesting and rare cases are cleared up and relegated to the classes of common diseases by time, which as a rule, shows us that that which we looked upon as a rare or extraordinary occurrence or disease was only an ordinary symptom out of time and place.

Case 1.—Uremia in which the first symptom was aphasia, which continued half an hour, followed by a convulsion and hemiplegia of the right side, with complete recovery in twenty-four hours. Mrs. C. K.—, aged 62, married, mother of five children. No miscarriages, no history of syphilis. Patient, who is very heavy, had never been sick before. At noon, Nov. 12, 1895, while at dinner, in the best of spirits, and with a good appetite, she suddenly quit eating in the midst of her meal, attempted to speak, then muttered some unintelligible syllables, grew very much excited on account of her inability to speak, arose from the table and walked about the room. On account of her unintelligible jargon the family thought that she was delirious and had suddenly become insane. She could

not make herself understood, was unable to answer the excited questions put to her, and on my arrival, about half an hour later, was much excited, tried to speak, but could not form her words. She stated afterward that she knew perfectly well what she wanted to say, but could not express her ideas in words. Her face was flushed and congested, the pulse full, hard and rapid. No vertigo, no vomiting, no difficulty of breathing. I obtained a specimen of her urine, and after ordering a purgative by mouth and enema, put patient to bed. On returning after about an hour, after making an urinalysis, I found her unconscious and in a violent convulsion. The convulsive movement, however, was limited to the right side. Respiration stertorous, face deeply congested, the left side flabby and relaxed, the right in a state of clonic convulsions, the left arm and leg were perfectly flaccid and relaxed. After cessation of consciousness the right arm and leg were completely paralyzed. A hot bath was given; profuse perspiration ensued on use of pilocarpin, diuretics and saline purgative. In about an hour consciousness returned, but patient was unable to speak. The paralysis of arm, face and leg was complete on awakening. About 6 P.M. patient began to talk, but not perfectly, still enough to make herself understood. In the course of the next few hours slight movement returned, and in twenty-four hours speech, as well as use of arms and legs, was again perfect. In two or three days the patient was able to be about. Was placed on a strict diet and treated with alkalines, baths and cathartics. The quantity of albumin gradually diminished, but never disappeared entirely. After eighteen months patient is still well; no recurrence of uremia or paralysis. Albuminuria still present; quantity very slight. One interesting point about this case is, the double focal lesion—the left third frontal convolution and the right psychomotor area.

Case 2.—June 15, 1896, Mr. H., aged 61, married. Parents both died at an advanced age, presumably from bronchopneumonia. Has not been sick from childhood. Has been a hard worker all his life, doing the work of a molder. While eating dinner, just at the commencement of the meal, patient suddenly lost consciousness, his head sank on his breast and he fell forward, slipping out of his chair on to the floor. He was immediately picked up, and regaining consciousness in a few moments, was supported to the next room and laid on a bed. The family noticed that his face was drawn to the left side, and that he made no use of the right arm or leg. There were no twitchings or convulsions of any kind. I arrived on the scene a half hour after the onset of the attack. Examination: Patient is still in a semi-unconscious condition; can be aroused, however, and is able to assist at the examination. Pupils are normal in size and respond to light, both sides are equal, ext. muscles of both eyes are normal in their function. The right side of face has lost its expression, the naso-labial fold is absent on opening the mouth. The left angle is drawn upward. The tongue is protruded to the left. No loss of sensation in region supplied by fifth nerve. The right arm can scarcely be moved; there is not a complete loss of muscular power, but the grasp of the hand makes scarcely a perceptible impression. The movement of the right leg is good, but there is scarcely any motor power. Reflex of right leg very much diminished. No loss of sensation of either arm or leg. The examination of urine revealed the presence of albumin. Purgation and sweating. On my return, in three hours, I was surprised to see the muscular power of the leg normal; that of the arm greatly increased. On the following morning the paralysis had entirely disappeared.

Albumin remained in urine for about four months. Patient was treated by alkalines and a non-nitrogenous diet. For past eighteen months the urine has contained no albumin at all, and the patient considers himself entirely well.

Case 3.—This case shows one of those variations of focal lesions in which the paralytic onset occurred independent of an attack of uremic convulsions. It is true that the diagnosis is somewhat in doubt. The apoplectic attack may have been caused by an embolism or a hemorrhage from a small artery in the neighborhood of the internal capsule. In my opinion, however, the paralysis was, at first, so complete and the recovery from it so rapid and complete that I have no doubt but it belongs to the category of uremic hemiplegias. J. P., Grand Rapids, Mich., aged 55, unmarried. No history of nervous trouble in the family. Has never been sick, nor nervous. Dec. 5, 1897, at 9:30 P.M., patient, while playing cards, suddenly noticed that he could not use the right arm. A few moments later, while standing on the floor, he sank and would have fallen had he not clutched the table with left hand. The face became distorted at the same time, by the loss of use of right side. No diminution or loss of consciousness; no loss of speech. Patient was taken home at once in a cab. Physician

called and diagnosed right-sided hemiplegia, with involvement of face. Urinalysis showed presence of large quantity of albumin. The attack was sudden, preceded only by violent frontal headaches of some weeks' duration. Regained use of arm in a few hours, and in twelve hours the recovery was complete. Use of leg regained in twenty-four hours. Six weeks after attack patient came to Cincinnati. The examination of nervous system was entirely negative. The examination of the urine showed the presence of 2 per cent. of albumin, large numbers of hyaline and granular casts. The case has been under treatment for Bright's disease for four months at the Good Samaritan Hospital. No uremic symptoms have shown themselves in this time. Quantity of albumin and number of casts remain the same.

Case 4.—The following case seems to bear out the theory that some local predisposition is necessary for the development of the paralytic attacks, some cause present in the center or area of the cortex which renders this center more liable to be attacked by the poisons circulating in the blood during an uremic attack rather than the other parts of the surface.

Mrs. F., aged 64, married, mother of seven children, all healthy, slightly nervous in temperament. Has never been sick before the development of Bright's disease some years ago. Was treated for arteriosclerosis and fatty degeneration of the heart for past three or four years. I saw patient first in January, 1896. Will not burden you with the text of examination except to tell you that she presented a typical fatty heart with well-marked arteriosclerosis. The urine on examination, dark in color, s.g. 1014, 2 per cent. of albumin, no sugar. In August, 1897, patient had a sudden attack of apoplexy, involving external recti of both eyes, especially of the right eye. The right side of face and tongue were involved. Complete loss of power in right arm and right leg, partial paresis of left leg.

Diagnosis: Acute bulbar paralysis, due to thrombosis of a branch of basilar artery in the upper part of pons. Patient gradually improved, and after three or four weeks was able to walk about, although with much difficulty. The arm recovered completely, the right side of face and tongue, as well as the right external rectus, have not recovered after a lapse of two years. On Feb. 12, 1898, was called to see patient. The attendant nurse stated that she had not been as well as usual for a few days; that she had had attacks of vertigo and felt sick at stomach; that she had been somnolent all morning and had not been able to move the right arm and leg at all.

Examination: Patient is in a stupor and can not be aroused sufficiently to assist in the examination. The pupils are contracted and equal in size. Unable to examine external muscles of eyes on account of coma. Constant twitching in the left side of face, loss of sensation of left side of face. The left arm and left leg are in a constant state of clonic spasm, the spasms are small in excursion and momentary in duration, following each other very rapidly. These spasmodic attacks have recurred three or four times in the course of two hours, an interval of fifteen minutes probably between each attack. During the interval the left arm and leg are seen to be completely paralyzed. The muscles are flaccid, relaxed, the limbs fall by their own weight without any resistance on part of the muscles; patellar reflex absent; sensation can not be examined into on account of presence of coma; right arm and leg normal.

Treatment: Purgatives; profuse perspiration induced by heat, applied by using hot water in rubber bags. Consciousness returned in about an hour. The diagnosis of left-sided hemiplegia confirmed by patient's inability to use the arm or leg. The paralysis of the arm and leg disappeared in four or five hours. On the next day the patient was in her usual condition. April 16, 1898, patient again had an attack of uremic coma, with left-sided hemiplegia, and again recovered as in previous attack.

A CLINICAL STUDY OF UREMIC PARALYSIS.

The four above-mentioned cases represent, with but slight exceptions, the clinical aspect of uremic paralysis, and we may draw up in a concise manner the clinical picture of this trouble.

The prodromal symptoms.—While the uremic focal disturbances seldom occur without premonition, we can hardly claim that it has any premonitory symptoms of its own. The onset of the attack is sudden, being either the ordinary uremic convulsion or a sudden attack of aphasia, Jacksonian epilepsy or paralysis, with or without convulsions or loss of consciousness. The premonitory symptoms are rather those of Bright's disease, uremic manifestations of a disor-

dered function of other organs of the body, or there may be general cerebral disturbances, such as headache, vomiting and vertigo, or localized edemata, which are manifestations of a general uremic condition. The uremic attack as such may be typical in its character or it may not be.

During an attack of uremic convulsions with loss of consciousness, heavy stertorous respiration, a full, bounding pulse and contracted pupils, the convulsions may suddenly cease on one side and continue on the other, the limbs in which the convulsions have ceased become flaccid, the muscles become relaxed and when lifted up and allowed to fall, drop in the direction that the force of gravity dictates, muscles offering no resistance whatever to the falling of the limb. The tendon reflexes are usually absent; there is very often a complete loss of sensation in the affected limbs. The face often becomes distorted by the convulsions of the muscles opposite to the paralyzed side, the mouth becomes drawn over to the side free from paralysis, while the cheek on the paralytic side becomes distended by each expiration. Boinet lays down the rule that the uremic hemiplegia coincides always with the uremic coma, and that it disappears when the coma disappears. While this is the rule, we know from the study of reported cases that there are very frequent deviations from it. In the typical cases, it is suddenly noticed during the progress of uremic convulsions that one side of the body is stricken with paralysis in the manner above described. This hemiplegic attack continues during the entire uremic convulsion, and if the outcome is fatal the paralysis is present up to the time of death. If the uremic attack yields to treatment, the hemiplegia sometimes disappears as soon as a patient regains consciousness. In other cases the paralysis rapidly diminishes and in a few hours the muscular power is completely restored. Dr. Rose of Berlin states that these attacks can occur like ordinary apoplectic attacks; after a good night's rest the individual awakens in the morning to find himself paralyzed. Only in very rare instances does the hemiplegia last longer than twenty-four or forty-eight hours.

In rare instances, as Case 3 illustrates, the hemiplegia may begin and become complete while the patient is in full possession of his senses, and the attack is not followed by coma. Case 2 illustrates a mode of onset also rare, in that the coma and hemiplegia occurred simultaneously without being followed or preceded by uremic convulsions. The paralysis has all the characteristics of being cortical in origin. In the beginning, perhaps but for a few minutes, it is complete, but rapidly gives way to paresis. The chief characteristics of the hemiplegic attack are the flaccidity of the muscles and the rapidity with which the paralysis disappears. Boinet states that another of its characteristics is its variability. It is complete in the morning, diminishes at night and increases again in intensity a few hours later. Lancereaux reports a case in which the hemiplegia occurred first on one side of the body, disappeared, and then occurred on the other side. Case 1 is similar to this, aphasia occurring first in a right-handed woman, and during the uremic convulsion which follows there is developed a left-sided hemiplegia.

It is probable that cases where the paralysis was of long duration and those that were followed by contractures were due to localized hemorrhages in the cortex.

The paralysis of the facial muscles may or may not be present in these uremic hemiplegias, and when it is affected it follows the laws of cortical facial paralysis, namely, the lower half of the face alone is involved. Raymond reports a case attended with ptosis. Passing amaurosis is common. Bernard, Chauffard, Raymond, Chantemesse and Level report cases in which the hemiplegia was associated with conjugate deviation of the head and eyes. Aphasia occurring with right-sided hemiplegia is quite common, but occurring alone, or as in Case 1, occurring in a right-handed person with left-sided hemiplegia is very rare. In all four cases which I have recorded, at the time of the examination there was an absence of the loss of sensation. Boinet, however, states that there is habitual association and hemianesthesia with the hemiplegia, and that this hemianesthesia has all the characteristics of being cortical in origin. It is always incomplete and exceedingly transitory, often disappearing before the disappearance of the paralysis. Raymond reports one case in which the anesthesia affected the entire body, another in which the conjunctiva and the mucous membranes were anesthetic.

It is possible that at times also the anesthesia may affect the special senses, as Boinet's case, where the sense of hearing was lost for three weeks.

The prognosis as a rule is bad; the majority of the cases reported were fatal.

The hemiplegic attack itself does not offer a bad prognosis. The prognosis depends upon the severity of the uremic attack, and in my opinion the presence or absence of hemiplegia does not have any influence on the prognosis of the attack, the hemiplegia disappears simultaneously with it or it rarely lasts longer than twenty-four hours. The second of the cases reported in this paper is completely cured, the hemiplegia and the kidney trouble having completely disappeared. I do not believe, however, that there is any relation between the intensity of the renal disturbance and the cerebral symptoms, in this sense, that the most severe forms of uremic convulsions are apt to be followed by cerebral symptoms due to some inherent localized weakness in the cortex of the brain. It seems that the cases that are associated with high temperatures are apt to be fatal. Conjugate deviations of the head and eyes are also unfavorable symptoms.

The diagnosis is very easy, if the examination is carefully made. While it is true that ordinary apoplectic attacks caused by hemorrhage in the brain or by embolism in the cerebral arteries may be accompanied with epileptiform convulsions, an examination of the urine will clear up the diagnosis by demonstrating the presence of large quantities of albumin in the urine. We must not forget, however, that patients with Bright's disease, especially those with interstitial nephritis have as a rule arterial sclerosis and may have an attack of hemorrhage into the brain during the uremic convulsions, which may give rise to hemiplegia during the comatose state. The variability and the flaccidity of the paralysis, the normal temperature and the rapidity with which the paralysis disappears, will enable us to make the differential diagnosis. It will not be so easy, however, to differentiate the purely functional cases of hemiplegia from those due to minute hemorrhages into the cortex (Senator).

A very interesting feature of uremic focal manifestation is the occurrence of localized Jacksonian epi-

leptic attacks. When we take into consideration, as we shall see later, the entire absence of any organic change in the cortex, this feature becomes all the more interesting, because of the almost universal association of an organic focal cortical lesion with the occurrence of circumscribed epileptic attack. Chantemesse and Tenneson, Chauffard and others have reported cases of this kind with autopsies, Dunin five clinical cases, Boinet one. In some of the reported cases the epileptic convulsions occurred during an uremic attack and in others preceded such an event or occurred entirely independent of a generalized intoxication. In Boinet's case four distinct and separate attacks occurred without the attack being followed by paresis. After the fifth attack, however, the hemiplegia set in. The above cited cases, and there are still more in literature, are pure cases of Jacksonian epilepsy. Case 3 in the present report presents a localized epileptiform seizure, but convulsions were present in the paretic extremities.

CAUSATION.

The real cause of these attacks is, as a matter of course, the uremic condition. As to the proximate cause, the authorities are divided. Most individuals afflicted with Bright's disease are subject to arteriosclerosis, and Senator holds that very often these focal disturbances are due to capillary hemorrhages. This opinion is shared by Wagner and Landois. That this may be true for some cases, especially those in which the paralysis does not disappear rapidly, can not be denied. But the autopsies which are on record seem to disprove this theory. Another theory advanced by Landois is that the paretic manifestations are the results of an exhausted condition of the cortical centers resulting from over-irritation resulting from the uremic excitation and subsequent convulsions. This theory, however, fails to explain the causation of those cases in which the Jacksonian epileptic attacks, the paresis or aphasia precede the uremic attacks or occur in uremia in the simple condition of coma.

There are but two possible explanations remaining: one is the toxic theory, and the other the theory of localized cortical edema of Traube. Traube's theory is the theory which ought to explain the occurrence of uremic convulsions in a mechanic way: as a result of edema of the brain there is produced a pressure upon the veins and capillaries which results in an anemia, consequently in a diminished supply of oxygen, and as a result we have the convulsions. Rosa (*Berlin Klin. Woch.*, Feb. 28, 1898) seems to think that it may possibly explain the focal symptoms of uremic attacks. Chantemesse and Tenneson seem to hold the same opinion, and Senator says that, "localized and circumscribed edema may explain the origin of uremic amaurosis and hemiplegia." The majority of authorities are agreed, however, that the uremic condition is the result of an auto-intoxication, and that the real cause of the manifestation is a chemic poison—urea, acetone or whatever it may be—which irritates the brain and produces the symptoms. But how does this bring us any closer to the answer as to how the focal symptoms are produced? We are once more thrown back on theory, for pathology reveals nothing to us. We know that certain parts of the cortex are more subject to the action of the chemic poisons than others, but if we wished to apply this to uremic conditions then the focal lesions ought to be

constant, occurring in every case of uremic convulsions. We can fall back on the old theory of individual idiosyncrasy, but it does not aid us much. Pierret and Senator hold that there may be circumscribed disturbances of circulation, a spasm of a vessel in a localized area of the cortex, resulting from some diseased condition of that vessel, such as arteriosclerosis. As a result of this spasm produced by the toxic condition of the blood, there is a localized ischemia with a subsequent paretic dilatation, stasis of blood, edema and intoxication of the cortex. Assenting to this theory, those areas of the cortex would be most likely to suffer which are supplied by vessels in a state of degeneration.

Baynaud and Pierret hold, and our Case 3 seems to bear out the statement, that focal manifestations due to old brain lesions, which, however, have disappeared, are apt to return as a result of all kinds of systemic infections and intoxications and especially in uremic conditions, which would seem to prove that areas of the brain weakened by some former insult and apparently recovered again, are more susceptible to intoxication than other areas which are normal.

PATHOLOGY.

The anatomic investigations of the fatal cases fail to throw much light upon the etiology of these attacks. They all agree in one respect, namely, that there are no gross anatomic lesions to be found in the brain or spinal cord. Prior to 1881 it was supposed that the hemiplegia occurring during uremic convulsions was always the result of a cerebral hemorrhage. Paetsch, about this time, reported three cases in which the hemiplegia continued up to the time of death, and in which the autopsy was practically negative. In one case he reports the presence of an excessive amount of cerebro-spinal fluid and edema of the brain; in another the brain was perfectly normal. The former autopsy was made by Israel, the latter by Jurgens. Paetsch, who holds the theory of localized edema and serous infiltration, lays much stress upon the occurrence of the edema, but we see edema of the brain quite frequently without any paralytic symptoms during life. Dunin publishes another case which presented during life nystagmus diplopia and hemianopia, in which the anatomic examinations, both gross and histologic, were entirely negative. It is interesting here as a collateral proof of the intoxication theory of these focal lesions to refer to two published by Oppenheim (*Charité Annalen*, 1888).

The first case was one of hemiplegia with aphasia, the other a typical case of Jacksonian epilepsy, both being associated with carcinoma, one of the breast the other of the stomach. It was supposed during life that the occurrence of the manifestations on the part of the nervous system were due to the occurrence of metastatic growths in the brain. The autopsies were entirely negative; no changes were found in the brain. The author believes that the attacks were due to intoxication, and that poisons may affect circumscribed areas of the cortex of the brain.

BIBLIOGRAPHY.

- Carpentier: *Presse Méd.*, Belge, 1880.
 Paetsch: *Zeitschr. für Klin. Med.*, 1881, vol. iii, p. 209.
 Jaekel and Raymond: *Rev. de Méd.*, 1885.
 Chantemesse and Tenneson: *Rev. de Méd.*, Nov., 1885.
 Sancieraux: *Union Medical*, 1887, p. 703.
 Seral: *Contribution à l'Étude de paralys. Uremique*, Paris, 1888.
 Chafford: *Arch. Médecin.*, July, 1887.
 Dreyfus: *Brissac. Gaz. Hebdom. Méd.*, July 20, 1888.
 Boinet: *Rev. Médecine*, 1892.

- Dercum: *Trans. Phila. Neurolog. Soc.*, Oct. 24, 1888.
 Dunin: *Berlin. Klin. Wochenschrift*.
 Rose: *Berlin. Klin. Woch.*, Feb. 28, 1898.
 Oppenheim: *Charité Annalen*, 1888.

DISCUSSION.

Dr. SEXTON—I think the considerations which were brought forward in the latter part of Dr. Hoppe's paper lead us really toward the truth in regard to the theory of most of these cases which he is inclined to classify as of uremic origin when hemiplegia is present. It is not quite clear to me how we can have a condition which must necessarily depend upon a local lesion in the brain, without that local lesion ever being demonstrable, and I think that the history of these cases as mentioned is somewhat negative in that direction. What I mean to say is that we can scarcely see the mechanism of the poison which is supposed to produce the hemiplegia. My own opinion would be that the great majority of these cases really depend upon a local hemiplegia, a rupture of minute vessels, in which the absorption of the clot is so perfect that after death the autopsies do not give any evidence of its existence. I do not think there is any doubt that we have many cases of hemiplegia which are due to small hemorrhages. The fact that a quantity of uric acid is found in the urine I do not regard as conclusive, because, as Dr. Hoppe mentioned, the condition of the kidneys favor these ruptures, and it is more plausible to explain the existence of the hemiplegia upon that hypothesis than upon the hypothesis that a limited area of the brain should be poisoned by the uremia, while the remainder of the brain escapes. I have seen a number of cases in which there was a transient hemiplegia, and a certain portion of these cases have albumin in the urine, and yet I was never able to satisfy myself that the presence of Bright's disease was the cause of the hemiplegia.

Dr. LOPOR—It has been my good fortune or misfortune to see a great many cases of hemiplegia that I believe are uremic in their origin. I cannot believe with Dr. Sexton that there is an organic base for the origin of them. Uremic hemiplegia following childbirth is more common than most neurologists know. Following in the course of Bright's disease I have noticed those cases that have a rise of temperature—and a rise of temperature is almost entirely secondary—do not get well rapidly or get well at all. In cases that have no rise, or a small rise in temperature, the hemiplegia disappears quickly after the disappearance of the albumin. I cannot see how a hemorrhage could take place from a ruptured blood-vessel and disappear almost as if by magic, as these cases do.

Dr. TOMLINSON—I would like to add the evidence of my experience to confirm the conclusions of Dr. Hoppe's paper, and to call attention to three cases published by Dr. Burr of Philadelphia, in which the history was the same, and which appeared to be apparently true hemiplegia, and yet in which the autopsy showed nothing. In the last year I have had four cases which came to postmortem, and all these cases were cases of interstitial nephritis, with the same symptoms manifested as those described by Dr. Hoppe. One thing I noticed particularly was that the paralysis was not complete; that there were some groups of muscles that were not involved, and that it appeared and disappeared in different parts, coming and going from time to time. The face and the limbs would be paralyzed; the paralysis would disappear from the arm and become more grave in the face, and vice versa. I can also confirm the prognostic effect of the rise of temperature; it has been the same in my own experience. Another thing I have noticed in these cases postmortem, has been the intense edema of the floor of the fourth ventricle. I have concluded that the cause of death in these cases has been the result of this edema, from the uremic poisoning. In my own cases I was able to know of the existence of the nephritis for a long time before the paralysis appeared, and by careful examination to find all the indications of chronic nephritis. Since I have become familiar with them I have been able to almost anticipate these cases and to know what the result will be according to the extent of the nephritis. The recovery of the cases will depend entirely upon the stage of the nephritis. On postmortem, in spite of the most careful study, I have never been able to find any local lesion or disturbance to account for it. I have not made up my mind entirely as to vasomotor edema or vasomotor spasm. I am disposed to think there is a spasm which temporarily deprives that portion of the brain of blood.

Dr. ESKRIDGE—I rise mainly to call attention to the differential diagnosis of Jacksonian epilepsy due to organic lesion from these spasms, due to presumed functional disturbance of the cortex. When we have an organic spasm in the cortex, the muscles involved in the spasm are the last to be affected. The temperature is always high on the affected side, and the patient loses consciousness. I have not met a case of spasm, due to uremic poison, in which there has been much difficulty

in making the diagnosis between that and a limited cortical lesion. I have met with cases in which I was able to make an absolute diagnosis as to whether the paralysis was due to organic lesion or some functional disturbance. As we all know, a few hours' waiting clears the diagnosis in these cases. As a rule, in organic lesions of the cortex, only those muscles affected in the initial spasm will be paralyzed after the convulsion is over. In paralysis due to the effect of a poison, in nephritis and hemiplegia, paralysis of one arm and one leg is common.

Dr. C. EUGENE RIGGS of St. Paul—I think that Dr. Hoppe is to be congratulated on the exceedingly happy way in which he has put this very practical subject before the Section. I am inclined to believe, however, that there is a golden mean that Horace tells about, in which we can probably take a safer position than by saying dogmatically that these are simply due to toxemia, or simply due to hemorrhage. I confess to great sympathy with Dr. Sexton's view. We know that one-third of the cases of organic hemiplegia are probably due to Bright's disease; we know that Bright's disease produces hyaline degenerative changes, and slight pressure will cause this disaster. But we also know that the resulting phenomena depend entirely on the location of the hemorrhage. We all have seen these limited aneurysms in the brain, and if one of these aneurysms happens to break, the first effect is insult to the brain, in which you will find certain phenomena described by Dr. Hoppe, while in a very short time that will all pass away. I think that death may not only be due to the influence of uremia—you have all seen poisons so extremely toxic as to cause death in a short time—but it may also account for the temporary phenomena of paralysis. That seems to me a thoroughly clear and legitimate conclusion. As soon as the kidneys act thoroughly and properly the paralysis will clear up. While the uremic toxemia may account for a number of these cases, still there are a number of other cases due to hyaline fibroid degeneration of the brain.

Dr. C. C. HERSMAN of Pittsburg—I can scarcely conceive the idea that these paralyses are caused by hemorrhage. In one case that I saw, some time ago, the paralysis was so profound that I was almost persuaded to believe it was a genuine attack of apoplexy, but the clearing up was very rapid, only lasting a few hours. My surprise was, not that it cleared up so soon, but that it was so complete. It seems to me that if a lesion were present, we could scarcely expect such rapid recovery. It is possible, however, that a miliary aneurysm occurring, the profound shock to the part might be the cause of paralysis, and on recovery from the shock, the lesion not being extensive enough to cause a destruction of tissue would answer for the cause and the rapid disappearance. The theory as to vasomotor disturbance has been disputed by many; in fact, all the theories that I have seen suggested have been disputed by others. Consequently we can not say that we have any definite conclusion that we can hold to; but I am inclined to think that the cause is not a lesion or a hemorrhage, any more than it is in the case of epilepsy, or any other convulsion which we might have that is not caused by hemorrhage.

Dr. HEINS—I have noticed that the gentleman in speaking of these cases spoke of them as rather milder cases, and therefore allowed the possibility of recovery in that way. They speak of the percentage of albumin. A case of nephritis is not measured entirely by the amount of albumin that you have. If the gentleman says that 2 per cent. of albumin shows a small kidney lesion, then I would like to ask him what would be a larger percentage? I do not wish to be critical, but it strikes me it is a matter of some importance. It seems to me, too, that it is a bad thing for practitioners of medicine to cast away their anchors, and when you have, as Dr. Sexton has mentioned, an actual lesion that is the probable cause, to throw it aside. It is better to reason it out that way, than to say that there was no hemorrhage, because then you simply have no cause at all.

Dr. HOPPE—In answer to Dr. Hein's question about the percentage of albumin; as a rule the prognosis of kidney trouble is inversely to the amount of albumin present, the most fatal cases of interstitial nephritis being those that have the smallest percentage of albumin. Sometimes men will die without the urine showing any albumin. So the amount of albumin present does not have any bearing on the severity of the attack or the prognosis. The only difference about the paper seems to be as to the immediate cause of the condition in the cortex. We have actual facts at our disposal, and autopsies which have been performed by the best men in the world. I would call your attention particularly to the autopsies made by Jurgens and Israel in Berlin; not only did they make gross pathologic examinations, but they made histologic examinations, and not

any trace was found of any change in the cortex. This is the history, postmortem, of all the cases in which the hemiplegia continued up to the time of death. When the Traube theory of localized edema was still in vogue, some of the autopsies showed edema of the brain. But we know that we find edema even in the brains of drunkards without having any symptoms of paralysis during life.

OCULAR HEADACHES.

Presented to the Section on Neurology and Medical Jurisprudence at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June 7-10, 1898.

BY WILLIAM H. WILDER, M.D.

Professor of Ophthalmology, Chicago Polyclinic; Instructor in Ophthalmology Rush Medical College; Surgeon and Pathologist Illinois Charitable Eye and Ear Infirmary; Ophthalmic Surgeon, Cook County Hospital.

CHICAGO.

This subject is such a large one, and the views and experiences of ophthalmologists as to the rôle played by the eyes in the causation of this important symptom are so diverse, that it is difficult to give it the careful consideration that it merits in the short time allowed. When we reflect upon the mysterious character of headache and the unsatisfactory explanations advanced to account for it, we can understand the difficulty the ophthalmologist encounters when he attempts to give a reason for the faith that is in him, and to correctly interpret his clinical data. Even the exact seat of the pain is not known. Who can say whether it is in the brain itself or in the membranes covering it? In acute inflammation of these membranes it is quite probable that the severe headache is caused by the irritation of the meningeal nerves, but that is entirely different from the functional affair we are considering.

Just what influence vascular dilatation exerts in the production of this neurosis is equally uncertain. If it should be true, as claimed by some observers, that sensory nerve-fibers exist in the vessels of the brain, we could understand how distention of these vessels would give rise to pain, even without the increase of intracranial pressure. It is not improbable that the congestion of the mucous membrane lining some of the accessory cavities of the nose is accountable for many of the frontal headaches coming on after excessive use of the eyes, and this view is strengthened by the clinical observation that such cases are relieved only when, with the correction of the ocular defect, there is combined suitable treatment for the diseased nasal mucous membrane. That there are headaches induced by eye-strain is undeniable, and the discovery that the relief of the strain is followed by alleviation of the pain marks a brilliant advance in our therapeutics.

Notwithstanding that other general conditions may underlie the development of this neurosis, and the ocular condition may be only one factor, still its presence may for the time be a dominant one whose correction will strongly influence the future history of the case. And should the headaches recur at some time subsequent to the relief of the eye-strain, such recurrence would not disprove the value of the treatment, but would prove that some of the other underlying conditions had become dominant factors. Such complex cases are not infrequent, and show the importance of careful examination of the eyes in all cases of headache in which there is any doubt as to the etiology.

In most cases of so-called ocular headache, there is a history of the pain coming on after either mod-