case, lace it up and suspend it; we can then take our leave feeling that our patient will rest in comfort and safety. As you will observe, these prepared cases are light, sufficiently firm, exactly fit every part of the leg, preserve the normal alignment of the bones, and by their gentle embrace invite repose of all soft parts that may incline to distortion by spasmodic muscular contraction. The malleoli and condyles are embraced; the calf and heel rest in depressions exactly fitting their normal contour and thus maintain extension on the principle of the double inclined plane, the heel, calf and lower third of the thigh lying in depressions. When tired of lying on his back the patient can, after the first nine days, lower the suspension apparatus and turn on his side, letting the leg rest on the bed, and I doubt if there would be disturbance of the fragments even should he throw the other leg over it. Should the fracture be compound, an opening can be made in the case for observation and dressing, but as the case spreads apart so easily, I prefer lifting it out by its cheese-cloth envelope for more thorough inspection and cleansing.

The anterior splint, composed of the same material as the case and exactly fitting all the normal curves of the leg, is two inches wide and fills the gap in front. The lacing passes back and forth over this, and, however tightly drawn, makes no impression on the circulation beneath. You will observe that the case is open clear down to the heel. This is to enable the leg to be lowered into its bed without letting up on extension until the calf and heel are imbedded and the case coapted to the leg. The foot-rest, of like material, is also molded to the normal curves of the foot, and can be fastened on or removed at pleasure.

In conclusion, let me say that the comfort and welfare of a patient with a broken leg should always receive our kindest consideration. Our thought, like an X-ray, should penetrate and unmask all abnormal conditions, and our effort should be to promptly adopt such remedial measures as the case may require. The dressing I have presented here is not perfect; but if it should be regarded as an improvement on existing methods, it shall be my endeavor to still further advance the lines toward my conception of a perfect "Normal Contour Fracture-Box."

Of the suspension apparatus, it is needless to speak. Each surgeon contrives his own. The necessity for suspension everyone must realize when we consider that a lateral motion of the hips can not be made without strain at the seat of fracture, and immobility of the malleoli and condyles which project does not insure against mobility of the fragments, enveloped as they are by the soft parts. Fixation at the knee and ankle by a fracture-box lying on the bed implies at least lateral fixation of the hips; otherwise there will be pain and mischief. Suspension, allowing, as it does, the leg to float freely with any movement of the body, is a safeguard against pain and injury from this cause. The old-time method of boxing and soaking the leg forcibly suggests Hamlet's idea of death: "To lie in cold obstruction and to rot." Then amputation in compound fractures was the rule rather than the exception. Now, with our antiseptic methods and abundant supply of scientific appliances, we approach such cases with a consciousness of mastery.

Let us have a Department of Public Health!

MALARIAL HEMATURIA.

BY F. G. DU BOSE, M.D.

SELMA, ALA.

The hemorrhagic type of malarial fever, while fortunately infrequent, is nevertheless a most serious disease. The overwhelming production of, and intoxication by, the malarial organism, causes extreme hyperemia of the viscera, and partial disintegration of the blood. Hemorrhage most frequently comes from the kidneys alone, although it may appear from the several mucous membranes simultaneously. These are occasioned by blood, nutritional and pressure changes in the congested area. I doubt if anyone has lived long in a malarial district and countenances Dr. Koch's theory that quinin is the causative agent. I have seen four cases where no quinin was taken prior to the hemorrhage; one of these, a child, died the following day from suppression of urine.

The previous history of these cases presents the mild remittent or intermittent fever in most instances. We have no premonition of the occurrence of hematuria. Its appearance on urination after a chill, is the first index to the real condition. Jaundice rapidly develops, varying from a yellow tinge to a bronze color. Temperature ranges from 101 to 105 F; pulse rapid, 100 to 120 per minute, small, with a thrill and often dicrotic; tongue thick, heavily coated, with imprint of teeth on margins. Vomiting and nausea are constantly present. Black vomit occasionally occurs. Bowels are constipated as a rule. Headache, pain in the back, and aching in the bones and extremities constitute the usual subjective symptoms. The mind is clear until uremic intoxication begins. The advent of delirium or coma is most unfavorable.

On account of the diversity of medical opinion regarding the treatment of this disease, I shall give in detail the method employed by me in ten successive cases without a death. The fatal case mentioned above was the first one in my practice, and received only the hypodermics of quinin and symptomatic treatment. As soon as the diagnosis is made, a hypodermic of morphia and atropia is given, followed immediately by a ten-grain dose of calomel, swept down with a glass of water. Five grains are repeated every two hours, till four or five more doses are given, or until the urine begins to be secreted in larger quantities. It has been my observation that calomel in such enormous doses, fortified by the hypodermic injection of morphin and atropia, is a rapid and most powerful diuretic. When so administered, I have never seen it act as a drastic purgative, nor have I witnessed evidence of mercurial poisoning. I remember after having given one patient thirty-five grains in divided doses, to have administered a goblet of Rubinat water eight hours after to secure the evacuation of the bowels. Besides being one of the best diuretics, it is also an efficient intestinal antiseptic, and perhaps a cholagogue. In cases where the patient can be seen two or three times a day, quinin should be given hypodermically, combined with morphin and atropia.

The following is the formula used by me:

Quinin bisulph.	5 gr.	32
Morphin sulph.	$\frac{1}{8}$ gr.	008
Atropin sulph.	$\frac{1}{200}$ gr.	00033
Aqua.	30 m.	195

Misce. Sig. For one injection. To be repeated every eight hours.

The skin over the calf of the leg, where the needle is to be introduced, should be scrubbed with a little

alcohol, and the medicine injected deeply into the muscles. I have never had an abscess to follow. I think these result either from lack of surgical cleanliness or from using a supersaturated solution of quinin injected at a higher temperature than that of the body. On cooling, recrystallization occurs; these crystals, acting as foreign bodies, are the beginning of abscesses. If patient can not be seen frequently, I give the following:

Quinin sulph.	5 gr.	32
Camphor monobromate.	2 gr.	13
Morphin sulph.	$\frac{1}{2}$ to $\frac{1}{8}$ gr.	.005 to .008
Atropin sulph.	$\frac{1}{200}$ to $\frac{1}{600}$ gr.	.00033 to .00011
Capsicum, pulv.	$\frac{1}{2}$ gr.	.03

Misce. For one capsule.

Sig. One every four or five hours.

If the temperature is above 102.5 F., I give the following:

Phenacetin	3 to 5 gr.	.20 to .32
Caffein.	$\frac{1}{2}$ to 1 gr.	.03 to .065
Sodium bicarb.	1 gr.	.065

Misce. For one powder. To be repeated every two hours until temperature is reduced.

Copious draughts of skimmed sweet milk and water are insisted on. One to two large saline enemata are given daily to increase diuresis. These are introduced high in the bowels and retained as long as possible. Sinapisms are applied across the back, over the kidneys, and to the epigastrium; hot-water bottles are placed upon the patient, especially during the time the paroxysm is likely to occur.

I have seen cases recover where death seemed almost inevitable. I have such implicit faith in the above treatment, that I no longer dread having charge of such cases.

Since writing the above, I have had occasion to treat a case of this disease, and herewith append the history:

Thomas J., white, age 40 years, a farmer, slept two miles from Blue Girth Swamp all the summer. He had chills every seven, fourteen, or twenty-one days from July, 1898. After each paroxysm, he would take three to five grains of quinin three times a day, and sometimes a small dose of calomel. He had taken no quinin for three weeks prior to this attack, which was initiated at 4 P. M., January 23, by a chill. Severe vomiting and purging followed, and at 8:30 P. M., temperature had risen to 104, pulse 80.

January 24, 9 A. M., temperature was 98, pulse 72. I gave fifteen grains of quinin. I was sent for in great haste at 5 P. M. and found patient had passed a few ounces of bloody urine. At this time temperature was 97.5, pulse 90. I immediately gave ten grains of calomel and a hypodermic of morphin, $\frac{1}{4}$ grain and atropia $\frac{1}{50}$ grain. At 9 P. M., his temperature was 104, pulse 120, he was constantly nauseated and vomited everything taken into the stomach, and complained of pains in the head, back, and extremities. I gave hypodermically, quinin bisulphate 5 grains, morphin sulphate $\frac{1}{8}$ grain, atropia sulphate $\frac{1}{100}$ grain, and prescribed phenacetin, camphor and caffeine to reduce high temperature. The bowels acted twice during the night.

January 25, at 8 A. M., he received another ten-grain dose of calomel and hypodermic of the quinin mixture. He had passed about 16 ounces of urine during the night, which was as black as ink. During this day his temperature and pulse remained the same as the night before. At 3 P. M., and again at 9 P. M., he received another hypodermic of quinin mixture. His bowels had not moved, although he was given a saline enema. His temperature was 104.5, and pulse 120; mind wandering, very restless, vomiting milk in large curds; gave another ten-grain dose of calomel and a saline enema at 9 P. M. The conjunctiva became yellow, and the skin slightly tinged during this day.

On the morning of January 26, he was very much jaundiced, his temperature was normal, pulse 110, urine passed in large quantities and much clearer. At 4 P. M., temperature was 101, pulse 115; he received two hypodermics daily of the quinin mixture, till January 29. The urine cleared up, temperature became normal, and the patient made complete recovery.

THE ETIOLOGY OF SEASICKNESS.

BY WM. EDGAR DARNALL, A.B., M.D.

Physician to the Atlantic City Hospital; Fellow of the American Academy of Medicine; Member of the Medical Society of New Jersey, and Vice-president of the Academy of Medicine of Atlantic City.

ATLANTIC CITY, N. J.

Seasickness is a thing that in these days of so much water travel is quite common, and to which it seems the medical profession has given but scant attention. There is little literature on the subject. It is one of those unsettled questions about which each man may pursue his own theory as to etiology unmolested, provided only that he hit upon some method for the relief of its symptoms. There is no pathologic evidence connected with it, so far as I know, and the clinical manifestations of the malady are all we have to guide us in seeking a rational cause for them. The assertion may safely be ventured, I think, that the seat of the trouble does not lie in the stomach itself primarily, but that the vomiting is a reflex phenomenon. It is not a disease, but the symptom of a condition.

Seasickness is no respecter of persons. It attacks rich and poor alike. Neither youth nor age permit escape. Robust health and constitution are no guarantee against it, and the fearful nausea and depression it produces come alike to all.

Without entering into a description of the familiar and vivid symptoms, which are indelibly stamped on the mind of everyone who has suffered from this malady, we may at once pass to the consideration of a cause which seems rational, and therefore worthy of study. This cause is the disturbance of the sense of equilibration, the organ of which is situated in the petrous portion of the temporal bone. In order to make the subject clear it will be necessary to review the anatomy and physiology of the internal ear.

The internal ear is composed of the cochlea, a vestibule, the semi-circular canals and the membranous labyrinth.

The vestibule is the common central cavity of communication between the parts of the internal ear. It is situated on the inner side of the tympanum, behind the cochlea, and in front of the semicircular canals. It is somewhat ovoidal in shape from before backward, flattened from within outward, and measures about one-fifth of an inch from before backward.¹ It is not necessary here to describe all the features seen on the walls of the vestibule, but on its inner wall should be mentioned a depression—the fovea hemispherica—perforated by several minute holes for the entrance of the auditory nerve. This is known as the “macula cribrosa.” On the upper wall or the roof is a depression—the fovea semi-elliptica—and to this is attached the utricle. Behind the vestibule are the three bony semicircular canals opening into it by five openings. They are about one-twentieth of an inch in diameter. Each presents at one end a dilation about twice the diameter of the canal, called the “ampulla.” The canals are superior, posterior and horizontal, and are arranged so that each is at right angles to the other two, thus forming three distinct planes for the body. The superior and posterior canals unite and form a common opening at the back of the vestibule.

The membranous labyrinth of the canal consists of a membranous tube loosely filling each canal, but dilated at the ampulla to correspond with the bone, and closely adherent to the bone at the ampulla. At each end these membranous canals are joined to a

¹ Gray's Anatomy.