appearance; the rest of the mucous surface was quite healthy and uninjured. The stone removed is marked *b*. In the left kidney there was a long spiked calculus, (*a* and *c* represent front and back views,) occupying two calices, so it could not be dislodged without laying them into one. The lining membrane in contact with the stone was so dark-coloured as to be almost gangrenous. This was, I suppose, the real cause of death. It is true that there was no extension of the pyelitis into the bladder, but I cannot believe that this operation would have set up the peritonitis. There was no circumstance attending the operation to excite peritonitis: there was no difficulty or delay in the extraction of the stone; no dragging of the soft parts from the opening being too small, and no incision into the post-vesical cellular tissue from the opening being too large; no injury or ulceration of the mucous membrane of the bladder giving rise to extravasation of urine.

I dare say that many of you will remember that, previous to his death, I suggested the probability of a stone incarcerated in the kidney being the cause of his sufferings after the operation. This opinion was derived partly from the fact that the stone I removed from the bladder had the shape and general characters of a renal calculus, and partly from the history of the case given by his mother. I suppose, therefore, that he died from sympathetic fever induced by the operation and incarceration of the renal calculus.

The following is the official report of the post-mortem examination by Dr. Montgomery:—

“Robert H—, aged twelve years, admitted into Abraham ward, Sept. 18th, 1861, under the care of Mr. Solly; died Oct. 10th, 1861. The body was emaciated. There was a wound in the perineum about an inch long. The boy had been operated upon for stone. The mucous membrane of the larynx and trachea was healthy. The lungs appeared healthy, and their substance was rather dry. The heart and its valves were normal. The liver was somewhat fatty; spleen very soft. In the left kidney there was a large branched calculus, and the substance of the kidney was rarefied; the ureter was not distended. The right kidney was healthy. There was a considerable quantity of yellow, coagulated lymph adherent to the abdominal walls near the pelvis. The bladder and rectum were glued together by recent coagulated lymph. The areolar tissue of the pelvis was extensively inflamed; it was suppurating. The walls of the bladder were considerably thickened.”

You will perhaps like to ask me whether, if I could have diagnosed the probability of a calculus in the kidney as well as in the bladder, would I still have operated? I answer most decidedly, Yes; and for this reason, that without the operation the poor child must have died in great agony, and that the operation gave him a chance of recovery. No surgeon can be certain that his patient has an incarcerated stone in the kidney, though he might suspect it, and trust to its passing, as the original one did, into the bladder. This has occurred before now, and a surgeon has been unjustly blamed for leaving a calculus in the bladder, when he really only left it in the kidney, where, of course, he could not reach it.

No. 53 was a healthy boy, eight years old. He went on from first to last without a bad symptom.

No. 54 was the boy on whom I had previously operated in the Hemel Hempstead Infirmary. He has again done well.

ON THE

OSTEOLOGY AND DENTITION OF THE ANDAMAN ISLANDERS.*

BY PROFESSOR OWEN, F.R.S.

FOR a long-coveted opportunity of becoming acquainted with this race, I am indebted to an energetic officer of the Indian Government—Dr. Mouat. On that gentleman's return to England in 1861, he showed me the skeleton of an adult Andaman male, which has since been presented by Dr. Mouat to the British Museum. The islands are divided into Great and Little, and the former into North, Middle, and South; so closely contiguous as to form a group 140 miles long, and not more than 20 miles across. They are inhabited by a race of undersized or dwarfed blacks, of great audacity and implacability to strangers. Their skin is jet black, their hair woolly, and their lips are thick, but less prominent than those of a Guinea

negro. They are in the habit of shaving off all their hair except the eyelashes. They generally inhabit the jungle, but they are migratory, and seldom stop in one spot. All have similar habits, and the same language and customs. They seem to have no notion of a Supreme Being, or of a future existence. They are naked, and not ashamed. They take fish left by the ebb tide in small nets, which are chiefly made by the women. They tattoo by incising the skin by bits of glass, without, however, inserting coloured matter, the cicatrix being whiter than the sound skin. The boys play with bow-and-arrow, and take good aim; at an early age they go into the jungle with their fathers. The encampments usually enclose a central place, in which there is one hut more carefully roofed-in than the others, and this is the residence of the local chief, who gives all orders with regard to fighting and retreating, as may be found necessary. Whenever a death occurs, the body is buried a few hours afterwards. It is covered with leaves, the carrier conveying it with his back turned to the corpse; and it is attended by two or three relatives to the place of interment, which is usually about a mile from the sea shore. The body is lifted into the grave by means of a sling, and then a small mound of earth is erected over it. About two or three months after the burial the bones are disinterred, carried to the encampment, and wept over by the relatives. The principal relative generally takes the skull, while others take various bones, which are nailed to the posts of their huts.

Upon the table is an Andaman skeleton. It is that of a person about thirty years of age. The cranium is well formed, and might be classed with those of the full oval type. The frontal region is narrow, but not low; the nasals are not flat, but moderately developed; the chin a little protruded, but not deep. The base of the skull offers all the usual human appearances. The teeth are not larger than those of Europeans, but seem large in proportion to the size of the jaws; all the lower molars have decided anterior and posterior roots. All the bones present the characteristic structure of those of the *homo sapiens* of Linnæus. The dimensions of the principal limb bones show that these Andamans are about four feet ten inches in height. They appear to have no idea of their own origin; and it has been inferred that they might be the descendants of African negroes sent by the Portuguese and wrecked at Pegu; but these are mere theories, and are not to be relied upon, as the Andamans do not present the peculiar characteristics of African negroes. I presume that the Portuguese would import from the Guinea islands persons of the usual stature; and it is hardly possible that the Andamans who are so bountifully supplied with provisions should have dwarfed to their present stature. I conclude, therefore, that they are aborigines, and merely represent the Africans in blackness, which they might have got from exposure for generations. They are more distinct from the Australian or Papuan negro than from the African, especially with regard to their molar teeth. There is no better evidence of their Malayan or Mongolian origin; but I offer no opinion that they originated in any locality of the opposite continent. Andaman might have been part of some extent of the dry land which had been seen in the range of the Himalayan mountains within the tertiary period. This has been the opinion of geologists for some time, and is alluded to by Professor Ansted in his “Ancient World,” and by Sir J. E. Tennent, who has offered valuable evidence in support of such an opinion from the peculiar zoology of Ceylon. The whole of the shores of the Andamans are skirted with coral reefs, and are the seat of active volcanoes; and we should, doubtless, err in opinion as to the origin of the population if we concluded that the islands are now as they were when they received their first population.

The Andamans are a most primitive people, at the lowest step of civilization, and have carried the inventive faculties no further than is necessary for the supply of their own wants. Their animal appetites are gratified in the simplest way; and when a male comes to years of puberty, a woman is assigned to him as his wife and servant. They are not cannibals; indeed, no traces of cannibalism have been found amongst them. Implacably hostile to strangers, the Andamans have made no advances to higher races which may have approached them. Perhaps the only change they have made is in the weapons which they use for warfare. So they may have existed for thousands of years; and there is really nothing on which the ethnologist can found an opinion as to the number of generations through which the Andamans have continued to exist. They may have existed through all the geological changes which their islands have experienced. The cranium of these people evidences an aptitude for civilization. They are dwarfish, but they have a much better formed cranium and limbs than

* Report of a paper read before the Ethnological Society, Jan. 14th, 1862.

the negroes either of Australia or Africa. (These points were illustrated by a well-executed diagram.) They have better developments of the nasal bones and of the upper jaw than the other tribes, but more particularly excel them in the oval shape of the skull, and its thinness.

OBSERVATIONS ON CERTAIN TUMOURS OF THE NECK,

WITH NOTES RELATING TO ITS SURGICAL ANATOMY.

By ANDREW MELVILLE M'WHINNIE, Esq., F.R.C.S.

(Continued from p. 34.)

BRONCHOCELE differs from all other swellings in the circumstance of its varying bulk dependent upon the general condition of the system. We find it affected even by external hygrometrical influences alone, increasing in volume to an inconvenient extent in damp weather, and becoming restored to its ordinary tranquil state under the opposite and more favourable condition of the atmosphere.

Enlargement of the thyroid may, under particular circumstances, simulate and be mistaken for other swellings. The following are the main features of such a one, which bore a resemblance, in some respects, to the tumour represented in Fig. 1.* The case derives additional interest from the fact of its having been considered by a distinguished practitioner to be an aneurism of one or other of the large bloodvessels of the chest.

An elderly man was lately admitted into one of the French hospitals on account of retention of urine from an enlarged prostate gland, from the effects of which he died. Whilst under treatment, the surgeon's attention was attracted by a marked peculiarity of the patient's voice (*une altération notable de la voix*), "characteristic of compression of the larynx, or of one of the inferior laryngeal nerves." On the right side of the neck, and immediately above the clavicle, was a tumour placed behind the sterno-mastoid muscle, ascending as high as the inferior border of the thyroid cartilage, and covered by a network of varicose veins. Externally, the common carotid, pushed aside by the tumour, could be distinctly traced over it. The swelling, which was firm, pulsated forcibly like an aneurism of a large artery, and which it was considered to be. This view seemed confirmed from auscultation, by which a bruit de soufflet was readily detected. M. Nélaton pronounced it to be an aneurism of the arch of the aorta, involving, perhaps, the brachio-cephalic artery. During deglutition the tumour remained perfectly fixed and immovable.

The post-mortem inspection showed how complete had been the error of diagnosis. The tumour was found to consist of an enlargement of the right lobe of the thyroid, extending downwards, as in my case, behind the corresponding sterno-mastoid and clavicle into the chest, where, by its pressure, it had considerably flattened the trachea. Its encroachment on the lower part of this tube explained the immobility of the tumour during the movements of the larynx. The right primitive carotid passed over its external surface. The left lobe was perfectly natural, and, from the absence of isthmus, was totally unconnected with the hypertrophied right one, as in the early foetal state. The left ventricle was found considerably hypertrophied, and the arch of the aorta was dilated to twice its natural calibre. Its pulsations had been conveyed through the thyroid tumour, which it had lifted up; whilst the bellows sound, referable to thickened and ossified semilunar valves, had been conducted in the same way, thus leading to the mistake in diagnosis.

In the museum of the Sussex County Hospital is a preparation which displays the following relations of a bronchocele to the large neighbouring vessels, the pulsation of which must have been strongly communicated to it during life:—The enlarged right lobe rests inferiorly upon the arteria innominata, and has the common carotid artery ascending over its front and outer surface. It is seen to have insinuated itself between that vessel and the subclavian; this, together with an enormous inferior thyroid artery, are placed behind the lower part of the tumour. Here the enlargement, to a smaller degree also,

of the left lobe would, however, if doubt existed, at once have indicated the nature of the affection. In vol. xxvi., second series, of the "Medico Chirurgical Transactions," just published, is a communication from my friend, Dr. Joseph Bullar, of Southampton, on the subject of pulsating bronchocele. A gentleman consulted him on account of two pulsating tumours, one on each side of the trachea, attended with so much purring sound, as well as vibration communicated to the touch, that the complaint had been mistaken for double carotid aneurism. The other patient was a female, aged fifty-two, having a bronchocele which pulsated. In these patients the eyeballs were particularly prominent. "Both were benefited by iodide of potassium internally and iodine externally, and by iron; but in both cases the chief point in removing the pulsation was by improving the general health: in one case by relieving great general debility by tonics, fresh air, nutritious food, and rest; and in the other, by diuretics, removing a dropsical condition and venous plethora. The most important lesson taught by them is in their prognosis. The appearance of both was formidable, and might easily excite the suspicion of aneurism or malignant disease; but in both the pulsation proved to be a harmless and remediable complication."

There are instances of tumours in the neck at first sight resembling goitre. Amongst those which have come under my notice may be selected the one represented in Fig. 4, here introduced as combining several points of interest worthy of notice.

Mrs. M. R.—, about forty years of age, of healthy constitution, was sent to me by a gentleman in the early part of this year, with a large pendulous tumour of the neck, which he regarded as a bronchocele of the left side; and, indeed, suspended

FIG. 4.

apparently from the thyroid region was a tumour of considerable size, with no discoloration of the skin, extending several inches below the clavicle, and realizing the description which has been given of a pendulous bronchocele—viz., "that it is not unlike the dew-lap of a turkey-cock, the bottom being the largest part."

Before placing my hand upon the tumour, or interrogating the patient, I requested her, in order to see the effect upon the throat in deglutition, to drink some wine. The swelling failed to accompany the movements of the larynx in swallowing. The absence, too, of sufficient corresponding fulness above, in the region of the thyroid body, coupled with the circumstance—to me an unusual one—of the swelling being confined to the left side, at once led to the belief that this disfiguring tumour was not of the nature of a bronchocele. Another surgeon happened at the time to come in, and pronounced it to be a hydrocele of the neck; a cicatrix at the lower part indicating a former puncture seemed to justify him in such view. Here, as in most cases, caution was necessary in forming a diagnosis. On inquiring into the history of the swelling, which was tolerably firm and unyielding to the touch, and not attended with pain, or subject to periodical change in volume, she stated that

* Vide THE LANCET of July 13th, 1861, p. 30.