accompanying table) refer to the effect of the combined serum and chrysarobin method on individual attacks of psoriasis. For this purpose it seems to me that the new method is of real value. As to its possible effect in preventing recurrences of the disease, the time is too short for conclusions to be drawn. To hope for such a result would be too much to expect in a disease so peculiarly prone to recurrence. Even if future attacks cannot be prevented, in the present state of our knowledge, it is at least a satisfaction to have an improvement in the method of treating the individual attacks of such an obstinate and chronic disease as psoriasis.

CONCLUSIONS

As a result of my own observations in a limited number of cases I would draw the following conclusions:

1. Autogenous serum, when used alone, does not appear to be of value in the treatment of psoriasis.

2. It is of decided value in many cases when used in conjunction with chrysarobin. Obstinate cases of psoriasis that have long resisted vigorous treatment with chrysarobin ointment will often yield to this remedy when injections of autogenous serum are given.

3. Intravenous injections of autogenous serum, if properly given, are entirely devoid of danger.

4. The technic of preparing the serum and giving the injections is comparatively simple.

I wish to express my thanks to Drs. Edward Pisko and Benjamin F. Ochs for their kindness in supplying some of the material.

[NOTE.—Since this article was submitted for publication a communication has appeared in the *Journal of Cutaneous Diseases* (1914, xxxii, 780) by Hilario entitled "A Contribution to the Auto-serotherapy of Certain Diseases of the Skin." Hilario reports excellent results in the treatment of five cases of psoriasis by the combined method of injections of autogenous serum and inunctions of a weak (2 per cent.) chrysarobin ointment. A comparatively large number of injections was given (as high as ten in one case) the amount of serum varying from 15 to 40 c.c. Hilario concludes that "The auto-serum plays a very active influence in the treatment of psoriasis in the way of reducing the resistance of the psoriatic tissue, thereby rendering it more apt to be acted on by the chrysarobin, in weak percentages."]

616 Madison Avenue.

Scientific and Literary Imagination.—The imagination of Darwin, of Pasteur, for example, is as high and productive a form of imagination as that of Dante, or Goethe or even Shakespeare, if we regard the human uses which result from the exercise of imaginative powers; and mean by human uses not merely meat and drink, clothes and shelter, but also the satisfaction of mental and spiritual needs.—Charles W. Eliot.

OBSERVATIONS ON MILITARY SURGERY IN THE EARLY WEEKS OF THE WAR

J. P. HOGUET, M.D.

Adjunct Attending Surgeon to Bellevue Hospital; Assistant Attending Surgeon to the General Memorial Hospital and Hospital for the Relief of the Ruptured and Crippled

NEW YORK

It has been said by a number of capable military surgeons within the last decade that the treatment of modern gunshot wounds would be a much simpler proposition now than in former years. It was shown that in the Spanish-American and the Russo-Japanese wars the effect of the modern high-speed pointed rifle bullet was a humane one. The wounds were small, non-lacerated, and above all, clean, with comparatively few exceptions.

This might have been true formerly, when the number of men engaged was comparatively few, when

transportation was rapid and when there was little use of the modern high-powered shrapnel shells, but from personal observation of the wounded in France during the early weeks of the war I can state without reservation that this is not the case.

Many surgeons have made the observation that in the month of August a large number of those injured by rifle bullets had absolutely clean wounds, especially when there was no involvement of bony structures. Since then the daily life of the soldier in the field has been an extremely arduous one. Facilities for personal cleanliness have been practically nil, and the average soldier's clothing in the field is in a frightfully

Fig. 1.—Ricochet or deformed bullet. This bullet had gone through the skin of the side of the thorax and then lodged in the elbow between the skin and the olecranon. The bullet had enough force left to fracture the humerus but not to shatter it.

dirty condition. This is well illustrated by the remark of one of the wounded who was being undressed on his arrival at a base hospital in Paris in November: he said that the trousers that were then being removed were the ones he started to the front in in the early days of August, and that he had not had them off once since then. The hospital bed was the first one he had been in since he had been called from his own home by the mobilization. It is hard to conceive how a bullet traversing such clothes as these can stay aseptic.

Shrapnel wounds—those caused by the lead shrapnel balls or parts of the casing of the shell—are, of course, attended by a greater amount of infection; on account of the greater size of the ball or shell fragment and their slower velocity, more clothes or other infectious material are brought into a wound, which is very much bruised and lacerated. Probably all these infections occur at the time that the man is wounded. Of course, there are a great number of wounded who

cannot be immediately removed from the firing line and who lie for hours exposed to the rifle and shell fire of both their own men and the enemy, but luckily these are the exception. Most of the wounded are first dressed at the front by the field surgeons, the

Fig. 2.—Typical butterfly fracture from bullet traveling at high speed.

great majority of whom do their work with judgment and ability notwithstanding the surroundings, with little equipment and amid danger.

Transportation of the wounded, considering all the difficulties and the great number to be transported, is being creditably done in France. The more seriously injured are being taken care of in the so-called "flying" ambulances which always stay within a few miles of the firing line. Those less seriously injured are removed to the trains, which now are fairly well equipped with the necessary supplies, and well heated and lighted. In charge of each train is an army surgeon, sometimes with assistants, and practically always with orderlies who attend to the comfort and minor wants of the men. Dressings are done *en route*, and these are practically always in good condition on the patient's arrival at the base hospital.

The military authorities up to the present have shown a desire to send as many wounded as possible to the south of France, and to keep the hospitals in the North empty, but at practically all the more important cities along the railroad there are temporary hospitals at the station into which those who are too sick to stand further traveling are taken, and then transferred to a permanent hospital in that city. At almost all stations on the railroads leading from the war zone kitchens have been established by the Red Cross, and the "secours aux blessés militaires," where hot food is supplied to the wounded.

Then, too, from the Pas de Calais district, where the fighting of late has been so heavy, a great number of wounded can be evacuated by the hospital ships to

such ports as Cherbourg and Havre, further south, or directly to England. These ships are well equipped, are large and comfortable, and each has a complete staff of surgeons who do whatever emergency operating there is to be done on board. All the patients are dressed as often as necessary, and on some of the ships there is a Roentgen-ray plant where every wounded man is roentgenographed, the plate developed and sent with him to the base hospital when he disembarks.

Practically all the wounds seen are from rifle bullets and shrapnel. One saber wound was observed at the American Ambulance at Neuilly¹ since the beginning of the war; there were no bayonet wounds. Accidental injuries, such as simple fractures of the leg or foot from being thrown from a horse, were sometimes seen, but were very rare.

RIFLE BULLET WOUNDS

The wound of entrance of the pointed .30 caliber cupronickelled steel rifle bullet now used by the Germans is generally a small one—smaller than the diameter of the bullet—while that of exit is generally larger. The course of the bullet is generally straight, but of course one that is nearly spent may be deflected or even stopped by a strong bone such as the femur. Large lacerated wounds are sometimes made by the so-called "ricochet" bullet (Fig. 1), that is, one that has struck a hard substance and has been deformed by bursting its jacket and then rebounding, striking the person. These are the wounds that give rise to the report that the enemy is using dumdum or explosive bullets. In order to prove the use of such it would be

Fig. 3.—Fracture of humerus from shrapnel.

Fig. 4.—Shrapnel wounds of the foot.

necessary to actually remove such a bullet from a wound.

The effect of the regulation German rifle bullet on each part will later be taken up, but its effect on bone

1. "Ambulance" is the name given to the military hospital; Neuilly is a suburb of Paris.

must be discussed here. When a strong bone, such as the humerus or femur, is struck by a bullet within its zone of maximum speed (5,465 yards), one of two things may happen: Either the bone is cleanly perforated—and this is very rare—or else it is completely shattered. In the latter case we have the typical butterfly fracture, as is shown in Figure 2.

In these cases there is such a transmission of energy from the bullet traveling at high speed that extensive comminution results. Near the center of the fracture the fragments are apt to be small, but long fissures extend up and down the shaft of the bone—in some cases into the epiphyses. These fissures make some of the fragments of a large size. On the other hand, when a bullet at the end of its trajectory strikes a bone, a simple fracture results, which very much resembles those caused by indirect violence in occupational injuries. In one instance the bullet, after striking the bone, had its force so nearly expended that it remained in

Fig. 5.—Fracture of femur from shrapnel.

the limb. Short flat bones, such as the clavicle or ribs, are generally shattered, especially when the blow is a glancing one. Small bones like the tarsus or carpus are most often extensively comminuted.

SHRAPNEL WOUNDS

Wounds from shrapnel are, in general, more severe than those from rifle bullets. A shrapnel wound is one made by a shrapnel ball, or part of the casing of the shell. The former is a solid lead ball of about .50 caliber, forty balls to the pound. When the shell explodes these balls are scattered in all directions in the so-called cone of dispersion, but not at a very great velocity. The shell itself is shattered into innumerable pieces of varying sizes. Hence it can be readily seen that shell wounds result from large missiles traveling at a comparatively low rate of speed, and are therefore characterized by a large amount of laceration to the parts. Perforating shrapnel wounds are rarely seen, in most cases the missiles lodging in the tissues and forcing in with them a large amount of clothing or parts of the soldier's equipment. These wounds, of course, vary greatly in size and are prac-

tically always multiple—seventy-five having been counted in one soldier. The destructive effect of shrapnel on bones is very great. Here the butterfly fracture is hardly ever seen, except when the bone is struck by one piece. In most cases there is an extensive comminution amounting really to a grinding of fragments, being extremely small, about the size of pebbles (Figs. 3, 4 and 5).

VARIOUS INJURIES

The exact statistics as to the relative frequency of injuries to various parts will not be brought out till after the close of the war, but it might be well to describe briefly here the more characteristic injuries of each part.

One saber cut of the scalp, which was a simple wound without fracture of the skull, was noted. One case in which a rifle bullet had completely traversed the skull from side to side was seen at the Beaujon Hospital on the service of Dr. Tuffier, and was the only one of its kind observed. Cases of furrow wounds of the skull were not uncommon; in most of these the furrow was anteroposterior and generally on the side of the head; they were most often accompanied by an extensive fracture of both inner and outer tables, but very little comminution was present. In one case of this class in which I operated at the American Ambulance, there was an extensive right-side extradural and subdural clot. There was a complete motor paralysis of the opposite arm and leg, which cleared up two weeks after the removal of the clot. Compound fractures of all parts of the skull were very frequent from shrapnel wounds, in which a portion of the overlying tissues had been torn away. Many cases were seen in which a portion of the soft

Fig. 6.—Abscess of frontal lobe of the brain with drain in position.

parts of the forehead, together with an eye, had been completely destroyed by shrapnel. Here there was generally a fracture of the frontal bone through which brain tissue exuded. In one patient, received after he had been in a hospital at Arras for a week subsequent to being wounded, I discovered at operation an abscess

in the frontal lobe on the right side. There were also some small particles of shrapnel behind the right eye (Fig. 6).

Wounds of the face and eyes by rifle bullets were not uncommon, especially since a great deal of the fighting has been done in trenches. Two cases in which the bullet had entered one cheek, completely

Fig. 7.—Shrapnel ball in pericardial sac. The picture is not very distinct as the ball was moved by every heart beat.

traversing the head and coming out at the opposite mastoid process, were seen. Both of these were uncomplicated, except for a complete facial paralysis on the side of the wound of exit. Several cases of the destruction of one or both eyes from bullets coming from the side were also observed. Shrapnel wounds of the face were one of the commonest of injuries. Some of the patients simply had very small particles of shell-casing embedded in the skin. Many others had large portions of the face destroyed, and of these there were numerous cases of injuries to the eyes and nose.

Fractures of the inferior maxilla by both bullets and shrapnel were numerous. They were all very complicated cases, as they were compound either externally or toward the mouth, or both, and the fractures were practically always multiple. In the shrapnel injuries part of the chin or side of the face was often torn away, and in many cases some of the underlying bone was gone. In most of the Paris hospitals these cases were turned over to the dental surgeons, whose work was excellent, and whose results were often brilliant. A number of very complicated fractures with loss of bone and overlying tissues were seen at the Beaujon hospital. Here the plan of treatment was to apply an interdental wire splint immediately on the patient's entrance. The wound of the soft parts was dressed until infection ceased, and any operation in the way of removal of dead bone, or plastic work, was done only when the wound was perfectly clean. This method apparently gave better results than when the interdental splint was applied only when the wound was clean.

Wounds in the neck are comparatively rarely seen. One artillery-man was seen with a large lacerated wound in the middle of the neck from a piece of shell. When taken off the field this patient seemed to be breathing his last. As the piece of the shrapnel was visible through the external wound, it was removed by the field surgeon. The patient immediately began to breathe normally, and it was then discovered that the phrenic nerve had been pressed on. In this case there was surprisingly little loss of blood. Another patient admitted to the American Ambulance with a bullet wound in the neck was almost exsanguinated, and it was found on necropsy that the vertebral artery had been completely severed. Large hematomas of the neck were sometimes seen, and it was found best in all these patients to cut down and ligate the injured vessels immediately. One or two isolated cases of injury to the cervical and brachial plexuses were noted, but were very rare.

Wounds of the chest were not common. Penetrating rifle wounds were sometimes seen. In some the bullet had completely traversed the thorax, and in a few it had remained in it. In all of these cases there was usually an extensive hemothorax, which practically always disappeared slowly of itself, or after a few aspirations. Only one case of empyema was observed, and in this case there was a perforation of the chest wall by a piece of shrapnel 1 by 2 inches in size. Practically all the chest cases showed a complicating fracture of one or more ribs with a considerable amount of comminution. A few had signs of pulmonary consolidation, which lasted a short while and cleared up.

Two very interesting heart cases were seen. In one a shrapnel ball had entered the left lumbar region,

Fig. 8.—Bullet from a machine gun lodged in the abdominal wall, between the peritoneum and the internal oblique muscle.

completely traversing the left side of the abdomen, and was found by roentgenoscopic examination lying free within the pericardial sac. The ball moved freely up and down at every movement of the heart and apparently caused the patient no inconvenience (Fig. 7). The other case of this kind was seen at necropsy

in a patient who lived only a few days after entering the hospital. Here a large piece of shrapnel had completely traversed the abdomen and lodged in the heart muscle.

A surprisingly large number of patients had non-penetrating wounds of the abdominal wall. Many of these simply had very small particles of shrapnel

in the iliac, or lumbar regions, but several unusual ones were seen higher up in the abdomen, the highest being in the sixth intercostal space in the anterior axillary line.

Bullet wounds in the lower part of the abdomen anteriorly or through the buttocks posteriorly have been found to be especially interesting, and particularly fatal. The reason for this is that here there is often a perforation of the large intestine with a resulting abscess. Apparently a retro-peritoneal cellulitis is set up in the neighborhood, so that a large blood-vessel may be eroded and a fatal hemorrhage ensue.

Wounds of the perineum are comparatively rare, though several cases were seen in which shrapnel balls or shell fragments were found lodged there (Fig. 9). The greatest amount of destruction of tissue from shrapnel observed was seen in a case of this kind. Practically all of the perineum had been destroyed here and a large part of the tissues on the inner aspect of both thighs.

By far the most frequent and most interesting cases from a surgical point of view that the war has brought out so far are those of injuries to the extremities. Here again the very superficial injuries from rifle bullets, and shrapnel must be emphasized. Some of the very superficial injuries from rifle bullets, or some of the more superficial wounds *en sèton* may sometimes remain clean, but practically every other wound of an extremity from either bullet or shrapnel is infected at

Fig. 9.—Shrapnel in perineum.

this stage of the war. It is for this reason that the compound fractures of the humerus and femur are so very serious, and the latter much more so than the former. When a fracture occurs in either of these bones very long fissures are most often present. There is infection of the soft parts and bone from the very beginning. In the former the infection seems to travel along the fascial planes between the muscles, so that abscesses may subsequently be found at a considerable distance both from the fracture and the wound of entrance. In the bone an acute osteomyelitis starts in at the main point of fracture and slowly spreads along each of the fissures. Hence we see the small fragments extruded first and later the large ones.

Several instances have been noted in which the fissures have extended into the epiphyses, and on account of this osteomyelitis we find an infected joint in a rather short period of time. Where the diaphysis of a long bone is struck there is extensive comminution, but fissuring into the shaft rarely occurs. In these cases, of course, infection of the joint is immediate. Com-

That the modern rifle bullet is not a harmless one when it penetrates the abdomen now seems to be an indubitable fact. The typical penetrating wounds are not seen at the base hospitals, or if they are they are seen only in the stage in which a localized abscess has been formed. From accounts of a number of men who have been doing active service at the front, few laparotomies are done there on account of inadequate facilities. The opinion of most of these men is that there is little effort on the part of the mucous membrane to close up the perforation. Most patients with abdominal wounds are now being transported to the nearest base hospital in a sitting position, and there treated expectantly. Luckily localized abscess formation is the rule, and in no case of this kind is operation performed at the base hospital unless there is a definite abscess. One case was seen at the American Ambulance in which the bullet had entered the left iliac region and had made its exit through the middle of the sacrum. This patient had no abdominal symptoms for almost one month, when an abscess began to develop on the left side. A number of cases of fecal fistulas were seen which had been evacuated to the base hospitals from those at the front after the acute symptoms had subsided. The majority were low down

Fig. 10.—Metal shoe-lace clasp driven into the leg by a shell explosion.

pound infected fractures of the humerus, even when accompanied by an extensive loss of bone and soft parts, yield much more readily to treatment than do similar injuries to the femur. Whether this is due to a better blood-supply to the upper extremity, or whether the foyer of fracture is easier of access here will not now be discussed, but the fact remains, and will probably become more and more evident during the course of the war. Amputation of the arm has rarely to be resorted to unless there is injury to the brachial artery high up, or unless some complication, such as gas gangrene, sets in. Amputation through the thigh, however, must be done in a fairly large proportion of cases, as it must be borne in mind that these patients become rapidly septic so that the first indication is to save the patient's life, the second to save the limb and the third to preserve as much function as possible. Conservatism in the treatment of these fractures is the rule in France and England. I have it on very good authority that such is not the case in the German army.

Conservatism is, of course, to be encouraged, but the treatment of these fractures is one that requires absolutely the best of surgical judgment. Good and efficient drainage must be established early and be maintained. Loose fragments of bone or pieces of shrapnel must be removed. The postoperative treatment is more important than the operation. The limb must be immobilized, and in the case of the femur, extension must be employed. A plaster hip spica, although carrying out both immobilization and extension, has been found to be impracticable. It soon becomes loose as the swelling of the limb decreases, and is dirtied in a very short time, as the amount of discharge from these wounds is enormous. The simplest method and the most practicable consists in extension with weights and immobilization by means of wooden splints on a fracture pillow. An idea of the gravity of these fractures, and the difficulties in their treatment, may be somewhat appreciated when it is remembered that these patients have for several months been living under a terrible strain, and often with insufficient food. The reason that conservatism can be used at all is that all these patients are young men.

Of the two dreaded complications of gunshot injuries — tetanus and gas gangrene — it is much too early in the war to speak. The French government had a supply of antitetanic serum on hand, so that it was a rule in many base hospitals that every soldier should be given an immunizing dose on admission. An idea of the comparative rarity of tetanus can be gathered when it is stated that in four hospitals in Paris — each with a capacity of about 400 beds — not one has had more than six cases.

Gas gangrene, however, is much more common and is especially prone to follow shrapnel wounds of the thigh. The methods of treatment differ in various hospitals. Most of the French surgeons do not believe in extremely large incisions for drainage, but make comparatively small ones, which are frequently irrigated with peroxid. Many are in favor of draining and forcing oxygen gas under pressure into the affected tissues for a considerable length of time — from twelve to seventy hours. This can be repeated at intervals if the process does not resolve. Both these methods of treatment have given good results in some

cases, but in many others amputation has to be done, and very often this is of little avail.

For the privilege of having been able to operate in and treat a number of the cases mentioned here I wish to thank Dr. Du Bouchet, Surgeon-in-Chief of the American Ambulance at Neuilly. To the many other surgeons in Paris, by whose courtesy I was able to see the cases under their care, my thanks are also due. The roentgenograms reproduced here were kindly made for me by Dr. Jaugeas and Mr. King of the roentgenographic department of the American Ambulance at Neuilly.

40 East Forty-First Street.

THE RELATION OF POSTURE AND INTESTINAL DERANGEMENTS TO COXITIS*

ROBERT B. OSGOOD, M.D.

BOSTON

The title of this communication is misleading. One has a right to infer that I shall at least attempt to explain the relation of posture and intestinal derangements to coxitis. I shall offer no explanation, however, and there should be an interrogation mark after the last word.

The paper is simply a report and a consideration of three cases in which there was an apparent relationship between faulty posture or intestinal derangements and coxitis, and in which the discovery of this relation made unnecessary any prolonged local treatment of the coxitis. The excuse for its presentation is the hope that not all of those present have noted such an association. If this is so and your experience should tally with mine it is possible that you may occasionally be spared a modicum of anxiety, a few families may be spared the distress which the diagnosis of tuberculosis always brings, and certain patients may be spared a prolonged and irksome treatment which may not be demanded.

You are all familiar with the belief held by Sir Arbuthnot Lane¹ that unless the capacity of the several tissues of the body to resist the entry of certain organisms is inhibited by an auto-intoxication resulting from intestinal stasis, it is impossible for these diseases to develop. Therefore, to meet these diseases he adopts measures to improve the drainage scheme and cites cases in which rapid disappearance of large masses of tuberculous glands in the abdomen has occurred, following disconnection of the large bowel.

Fraser,² writing on the types of organism in bone and joint tuberculosis in children, is quite convinced that the more common bovine form enters usually by the mouth in milk. It is probably true that an intestinal stasis favors extension to the bones and joints.

Barrington-Ward,³ working at Great Ormond Street, reports a case of double tuberculous disease of the hip with sinuses, in which orthopedic treatment had been unavailing, but in which rapid recovery followed a short circuiting operation after the manner of Mr. Lane.

It is not, however, to suggest that faulty posture and intestinal derangements bear a causative relation

* Read before the Section on Orthopedic Surgery at the Sixty-Fifth Annual Session of the American Medical Association, Atlantic City, N. J., June, 1914.

1. Lane, Arbuthnot: *Lancet*, London, Dec. 14, 1912.

2. Fraser, S.: *Brit. Med. Jour.*, April 12, 1913.

3. Barrington-Ward, L. E.: *Surg., Gynec. and Obst.*, July, 1912.