Thrombosis, carrying with it the possibility of embolism, may develop after any type of operation, but the element of risk seems greater in the case of operations in the upper abdomen than in those involving the pelvic viscera. Post-operative pulmonary embolism usually occurs between the eighth and twentieth days of convalescence. In the majority of cases, fortunately, the embolism is not fatal. When it involves a small vessel or when the embolism is lodged in an organ where vascular anastomosis is free, embolism is often unrecognized. In an analysis of 16 fatal cases, Busch found that death was instantaneous in one-fourth. In the others, ten minutes or more elapsed before death took place. Novak discusses the various factors to which an etiological role has been attributed.

A Statistical Study in Serology. By Robt. S. Preston, Richmond, Va. *The Old Dominion Journal of Medicine and Surgery*, April, 1915, pp. 202-205.

The author in his study states that he has found it essential to the correct interpretation of incomplete complement fixation tests that the clinical and laboratory finding be correlated. Incomplete Wassermann reactions and indefinite luetin tests are as apt to be overestimated as are negative Wassermans. The latter does not exclude the possibility of a luetic infection. To do this to the best of our ability several tests should be performed at intervals, and if necessary after a provocative injection of salvarsan has been administered. Salvarsan is more efficient in producing the Herxheimer phenomenon than mercury, though mercury may be utilized previously for this purpose on account of the facility of its administration.

Of 11 cases that had received antispecific treatment 41 per cent gave a positive Wassermann reaction. Those cases (368) diagnosed and grouped as being syphilitic in nature gave 79 per cent positive reactions. Two hundred and forty-four cases treated or untreated diagnosed as primary, secondary and tertiary lues gave 90 per cent positive tests.

In the routine cases for diagnosis one or two cases of paroxysmal hemo-globinuria gave a positive Wassermann, as also did a case of hematuria, one case that had previously had a splenectomy performed, one case clinically a gastric carcinoma, and one with gastric crisis operated on for perforating ulcer. Of seventeen cases of aortic diseases, six gave positive tests. Of three cases of Menier's diseases partial fixation was obtained in two, but no response could be obtained by treatment. Of eight cases of syphilis of lungs and pleura, two were diagnosed pleurisy and four were known to have been considered phthisis previously.

The Significance of Bacillus Coli in Pasteurized Milk. By L. P. Shippin, Baltimore, Md. *Journal of the American Medical Association*, April 17th, 1915, pp. 1289-1291.

The presence of *B. coli* in pasteurized milk is commonly taken to indicate either improper pasteurization or subsequent contamination of the

milk. This organism was supposed to have a thermal death point below the temperature used in pasteurization, and in consequence it was believed that it could not survive this process. However, De Jong and De Graef claim to have isolated strains which are capable of withstanding a temperature of 149 to 152.6 degrees F. for thirty minutes. These strains would not be killed by the degree of heat usually applied in the process of pasteurization, and their presence would not mean that the milk had been improperly heated.

The writer succeeded in isolating from pasteurized milk seven strains of *B. coli* which survived temperatures ranging from 143.6 degrees F. to 147.2 degrees F., applied for thirty minutes. This confirms the work of De Jong and De Graef, and indicates that the presence of *B. coli* does not mean necessarily improper pasteurization or subsequent contamination of milk.

Diuretics in Renal Diseases. By Leslie Byron Wiggs, Richmond, Va. *The Virginia Medical Semi-Monthly*, April 9th, 1915, pp. 7-10.

The author protests against the promiscuous use of diuretics in the presence of renal damage. He supports his views by the results of experiments on rabbits in which he had produced fatal acute nephritis. The use of diuretics in these animals in each instance shortened their lives. He believes that if this is true in the acute conditions that it is equally true in the chronic types, for in both instances the cause of the nephritis is irritation, and he argues that it is illogical to further irritate them by the use of diuretics, for "cells that fail to functionate as a result of stimulation or irritation will probably not be permanently benefited by further stimulation or irritation." What they need under such circumstances is rest, and his clinical experience justifies such an hypothesis.

He summarizes as follows:

1. In chronic cases of nephritis with rising blood pressure and a normal chloride and watery content in urine associated with a low urinary urea, diuretics of the vascular type, may be used with caution.

2. In severe acute cases diuretics are contraindicated.

3. In cases definitely known to be mild or subacute in type, less irritant diuretics may be used guardedly. . . . Since, clinically, these types are difficult to differentiate from the early manifestations of severe forms, it seems a good rule to interdict the use of diuretics in doubtful cases.

4. Theoretically, there seem to be few logical reasons for the use of renal stimulants or irritants; therefore their indications in any case are clothed in extreme doubt.

Bell's Palsy and Neuritis of the Eighth and Twelfth Nerves Following Salvarsan Injection. By Virginius Dabney, Washington, D. C. *Interstate Medical Journal*, May, 1915, pp. 478-480.

Foreign observers have reported deaths and cases of neuritis but such reports are not common with us, whatever be the motives. This has served to lead many into the use of salvarsan through lack of warning as to the possibility of

serious results. The wide publicity given the remedy of Ehrlich, and the erroneous interpretation of his claims, together with the lack of knowledge of the users, seem responsible. Salvarsan is dangerous, and should not be used without mercury as a preliminary. It is an arsenical preparation and shares all the poisonous properties of that mineral. Case: Healthy locomotive fireman, age, 23; syphilitic infection two years before; hemiatrophy of tongue, facial palsy, labyrinthine irritation. Two weeks after injection paralytic symptoms appeared; second injection in 30 days. Apparently pressure on 7th nerve in fallopian caused atrophy. That the Bell's palsy and neuritis of the 8th and 12th nerves was due to the salvarsan and was neither a syphilitic neuritis nor a "nervous relapse" is the writer's firm belief, based on these facts: He had had no mercury to mitigate the tremendous reaction of the arsenic on the syphilitic deposits in the nerves; nervous relapses occur only some time after the cessation of treatment (this occurred immediately after treatment had begun); the Herxheimer reaction occurs within 36 hours (this case in ten days).

American Thermal Springs: Their Therapeutic Uses; Radioactivity of Springs in Germany, Austria, and Virginia, with a Regime for Patients With Gout, Rheumatism, and Arthritis. By Guy Hinsdale, Hot Springs, Va. Medical Record. May 1st, 1915, pp. 722-725.

The origin and source of heat of thermal springs are discussed with a list of some of them east of the Rocky Mountains. A list of radioactive springs in Austria, Germany and Virginia shows that the latter rank very high, the Magnesia Spring at Hot Springs, Virginia, containing 274 Mache units per liter. The Virginia springs were tested by Professors Hemminger and Zueblin about a year ago. Any spring containing a radioactivity of over 100 Mache units per liter should be classed as strongly radio-active. Dr. Hinsdale refers to the febrile reaction commonly noted when gouty and rheumatic subjects bathe in thermal springs and he appends a regime for those patients: (1) A tub bath at 102, 103 or 104 degrees F. Vigorous or gentle rubbing in the tub, especially of affected joints or other areas. Duration of bath 8 to 12 minutes. (2) A hot, dry pack, on leaving the tub. Duration of pack 10 to 20 minutes. (3) Cloths wet in ice water while patient lies in tub and in pack. (4) A douche of cold water all over the body after the pack. (5) A more or less vigorous rubbing all over the body in pure alcohol, or cocoanut oil may be used. After three successive days of this treatment a day of rest. Rest in bed after each bath. Twenty or twenty-one baths is the proper number for average cases.

A Treatment for Puritus Ani. By Harvey B. Stone, Baltimore, Md. Read before the Medical and Chirurgical Faculty of Maryland, April, 1915.

The success of alcohol injections in producing local anaesthesia in facial neuralgia cases, suggested that the same anaesthesia might be produced in the perianal skin by destroying the superficial cutaneous nerve filaments with alcohol

injections. The danger of producing motor paralysis of the sphincter muscle was appreciated, as was also the possibility of causing superficial skin sloughs. In view of these disadvantages the method was first tried on dogs. It was found that deep injections that brought the alcohol into contact with the nerves of the sphincter muscle produced incontinence. Also, injections into the cutis produced skin sloughs. A proper placing of the injection, however, under the skin, but not in it, on the one hand, and on the other hand not deeply carried into the perianal structures, succeeded in producing anaesthesia without either sloughs or incontinence.

In view of the success of these experiments the method has been applied to clinical cases. So far five cases have been subjected to this treatment. Two small superficial sloughs resulted from imperfect technique in the injection. There has been no interference with the control or function of the muscle. All of the cases have been relieved entirely of the itching in the area injected, although some of them had suffered persistently in spite of all sorts of treatment for many years. How long the relief will continue is not yet known. The case of longest duration has shown no recurrence after five months.

The injection is made with 50 per cent alcohol preceded by novocain. The whole area in which the itching occurs is underlaid with the injection.

The Medical Treatment of Peptic Ulcer With Especial Reference to the Lenhartz Treatment. By George Blumer, New Haven, Conn. Johns Hopkins Hospital Bulletin, May, 1915, pp. 127-132.

The article reviews the different regimens of rest and diet which have been used in the medical treatment of peptic ulcer and classifies them into three groups: (1) The Leube group, in which the underlying principle is as complete rest to the stomach as possible; (2) the Lenhartz group, in which the principle is the frequent administration of small quantities of acid-binding food to neutralize the excess of free hydrochloric acid and keep up the general nutrition; (3) The fat group, in which advantage is taken of the non-stimulating and non-irritating effect of fats.

The article discusses the advantages and drawbacks of the different diets and the conclusion is reached that each has advantages and disadvantages, and that the wise physician will use them as frame-works and try to individualize the diet to suit each patient.

Larva Migrans. By Henry C. Dozier, Ocala, Fla. Journal of the Florida Medical Association, May, 1915, pp. 326-329.

The following conclusions drawn from a series of cases of Larva Migrans, have not been observed or reported so far as the author knows. See original article for argument and details:

First.—Infection from water (sea-shore or lake), the larva being probably that of the gad-fly, horse-fly, or deer-fly, who attach their eggs to water or wet earth, and are carnivorous.

Second.—It seems probable that a point of entrance is necessary for the larva to enter the skin. Some scratch or denuded area, as is so