VI. *On the Cuticles of some Recent and Fossil Cycadean Fronds.* By H. HAMSHAW
THOMAS, M.A., F.G.S., Curator of the Botanical Museum, Cambridge, and NELLIE
BANCROFT, B.Sc., F.L.S., 1851 Research Scholar, Newnham College, Cambridge.

(Plates 17–20. and 32 Text-figures.)

Read 6th November, 1913.

CONTENTS.

	Page
I. Introduction	155
II. Cuticles of some Recent Cycads	157
(a.) Material and Methods	157
(b.) Previous references	157
(c.) Description of the Cuticles	158
i. Introduction	158
ii. Detailed description	158
iii. General summary of the structure of the recent Cuticles	174
III. Cuticles of Fossil Cycadean Fronds	177
(a.) Previous observations on cuticular structure	177
(b.) Detailed description	179
(c.) The Cuticles of some other types of Fossil Cycadean fronds	194
(d.) Summary of the structure of the Fossil Cuticles	195
IV. Comparison of the Cuticular Structure of Recent and Fossil Cycadean Fronds, with notes on that of other Gymnosperms	197

I. INTRODUCTION.

IN an investigation of the cuticles of recent and fossil Cycads our attention is principally concentrated on the form and structure of the stomata, a study of which has attracted many observers. The physiological functions of the guard-cells are so important, and their shape and action are so uniform in plants of widely separated classes, that they may be regarded as highly adapted to their special work, and as structures which were evolved at a comparatively early period in phylogenetic development. But while the morphology of the stomata is uniform, many differences of structure, especially of thickening, do exist, rendering the work of classifying stomata by no means easy. One of the few general attempts to make a survey of the whole range of these structures, to classify them in groups, and to find broad general tendencies in their variation, is that of Porsch (1905), who has introduced a conception of a phylogenetic ground-plan modified by adaptations to external conditions. The idea that stomatal structures

may be valuable from the phylogenetic standpoint has not yet been worked out in any detail, nor has any attempt been made to compare modern plants with their fossil ancestors with regard to this character. One of us (Thomas, 1911) has shown, however, that some of the Carboniferous Calamites possessed the transverse bands of thickening which are so characteristic of the guard-cells of the modern Equisetums, and it seemed important to make further investigations of this kind. An especially favourable opportunity is presented in the Cycadophyta; the modern representatives of this group possess considerable variety of frond form, while the fossil fronds are also varied, and are abundant in a fairly good state of preservation in Mesozoic rocks. From a purely palaeobotanical standpoint the work offers some attractions, because we have so often to deal with isolated fossil fronds which possess few points of distinction apart from external form; the possibility of classifying them and obtaining information as to their affinities by means of a study of their epidermal structure seems important.

So long ago as 1856 Bornemann made a brief comparative study of the epidermal characters of recent Cycads for the purpose of identifying fragments of fossil Cycadean fronds from the Lettenkohle beds of Thuringia, but his studies of the fossil forms were not very critical. He thought that the Mesozoic fronds were nearly related to if not generically identical with, some of the modern forms, and therefore expected that the study of the latter would at once facilitate the identification of the fossils. We are now tending to regard the fossil Cycads as more and more widely separated from their recent representatives, and we may at the outset expect the similarities in their structure to be few.

Recent work on the fossil and living members of the Cycadophyta has raised them to a very important position among the Gymnosperms. The investigation of the Carboniferous Pteridosperms, and the recent discoveries with regard to the Bennettitales have caused many botanists to seek for the origin of the Angiosperms amongst the ancestral plexus of Cycadean forms, much attention having been focussed on the Mesozoic plants of this affinity. It is, therefore, important to utilise every form of evidence as to the relationships of these plants.

The method of enquiry adopted in the present paper is, first, the detailed investigation of the epidermal structures found in the modern Cycads, and the attempt to find a type of stomatal structure characteristic of the group, the departures in detail from this type being noted. Subsequently, the epidermal structures of all the available Mesozoic fronds are described, and an attempt is made to estimate their relations in this respect to one another and to the modern fronds. In studying the fossil fronds, although we have many beautifully preserved examples of their cuticles, yielding much information in surface view, we have as yet been unable to get satisfactory preparations showing the guard-cells of the stomata in section; we are thus placed in an unfavourable position for comparing them with the recent structures.

II. CUTICLES OF SOME RECENT CYCADS.

(a.) MATERIAL AND METHODS.—Most of the material used in the investigation of recent cuticles was specially obtained from the Glasnevin Botanical Gardens, Dublin, and from Kew, and was fixed in medium chromacetic acid. In some cases ordinary laboratory material was used.

The types examined included *Cycas* (5 species), *Dioon* (2), *Ceratozamia* (3), *Macrozamia* (6), *Zamia* (8), *Encephalartos* (8), *Bowenia* (1 form), *Stangeria* (2 forms), *Microcycas* being the only genus unrepresented.

The upper and lower cuticles of the leaves were examined in surface view and in section. In many cases the epidermises were stripped off easily, but in others it was necessary to use Schulze's macerating fluid. Sometimes pinnæ were treated with the fluid and then washed in ammonia, as in the case of the fossil cuticles (*vide infra*); after this treatment only the cutin layer remains, with little or no trace of underlying cells.

Permanent preparations of cuticles removed in either of the above ways were made. Bismarck brown, Congo red, and diamant fuchsin were used for staining. Lloyd (1908), in his researches on the Physiology of Stomata, seems to consider Congo red more satisfactory as an epidermal stain than Bismarck brown; in the present case diamant fuchsin was most used, as it is very rapid. The stain fades a little, however, on exposure of the preparations. Unstained preparations of cuticles were also largely used. In the case of the sections, one of the most satisfactory stains was found to be Scharlach R (also used for macerated cuticles) prepared as directed by Hill (1912). By it the cutin layer and cuticularised parts were clearly defined, staining a bright brownish red, all the other tissues remaining unaffected. Phloroglucin with hydrochloric acid was used to show the lignified lamellæ which occur in every case on the dorsal and ventral walls of the guard-cells. Serial sections stained with cotton red and aniline blue, and with diamant fuchsin were employed in the correlation of surface views and sections of stomata. Other tests were used, but an account of them need not be repeated here, as descriptions are to be found in the literature quoted.

(b.) PREVIOUS REFERENCES.—The most important previous accounts of the epidermises of Cycads include that given by Bornemann (1856), whose description is in connection with some fossil types and includes remarks on the chemistry of the epidermal walls and on the preparation of "artificial cuticles"; Kraus (1866) and Nestler (1895) also give good descriptions of some Cycadean cuticles. These researches show that each genus is fairly well characterised by a peculiar epidermis; sometimes the genera may be divided into sub-groups according to the structure of the epidermis (see Bornemann, 1856, p. 42, and Kraus, 1866, p. 13). Figures and descriptions of the stomata of some Cycads are given by Schwendener (1881), Strasburger (1866–67), Mahlert (1885), and Porsch (1905). More detailed references will be given in the description of each genus. The last-named author mentions the fact that though there is a general type of stoma characteristic of the Gymnosperms as a whole, there is one special type for almost each genus when the finer structure is taken into consideration.

(c.) DESCRIPTION OF THE CUTICLES.

(i.) Introduction.

In a consideration of the cuticles* of recent Cycads with a view to comparing them with those of fossil fronds, the features which seem to be of greatest value are those which may be determined from an examination of surface views. The actual form, size, and arrangement of the guard-cells, and the thickening of their walls as seen in section, are very constant throughout the group, and the variations mostly depend upon the appearance in surface view. The chief features to be taken into consideration are as follows :—

(1) The form and outline of the epidermal cells of upper and lower surfaces of the leaves, both between and over the nerve-courses ; the pitting and thickness, straightness or irregularity of the cell-walls.

(2) The distribution and position of the stomata, whether scattered or in definite restricted areas ; whether irregularly placed or having their axes in a fixed direction with regard to the axis of the pinna.

(3) The appearance of the stomata in surface view, depending upon the presence or absence of an intercalated series of cells between the guard-cells and the general epidermal cells, and the consequent degree of sinking of the guard-cells below the epidermis ; the shape of the stoma-opening.

(4) The number and arrangement of the subsidiary cells† (the epidermal cells surrounding the stoma, or in the case of much sunk guard-cells, those around the stoma-opening at the surface of the leaf).

(5) The presence or absence of crystals, either in the epidermal cells or in inter-cellular spaces.

(6) The presence or absence of hairs or hair-scars.

(7) The thickness of the cutin layer, and the presence or absence of foldings or striations upon it.

(ii.) Detailed Description.

STANGERIA, T. Moore. Pl. 17. figs. 1-6 ; text-fig. 1.

REFERENCES :—Bornemann (1856), pp. 41, 45 ; Taf. 12. figs. 19, 20. Kraus (1866), p. 34 ; Taf. 23. figs. 28, 29. Nestler (1895), p. 363.

The typical *Stangeria paradoxa*, T. Moore, and its variety *schizodon*, were examined and found to be exactly similar in epidermal and cuticular characters. The genus is distinct from all other Cycads, in that the epidermal cells of both upper and lower surfaces of the leaf-lamina have undulate lateral walls, like those of the epidermal cells of ferns. These cells are irregular in form and arrangement, and are usually three- or four-angular. Over the midrib of the pinnæ the undulations are absent, and the cells

* The term "cuticles" is used to include the layer of epidermal cells and the overlying layer of cutin.

† "Wall-Zellen" or "Nebenzellen" of the German authors.

are more regularly rectangular (text-fig. 1). Over the lateral veins also the cells are somewhat rectangular, narrow, and elongated, but the walls are undulate. At the pinna-margins the cells gradually become smaller and straight-walled. According to Kraus the cell-walls are unequally thickened, more secondary material being laid down on the outer side of a bend than the inner. Very little difference was noted in the present case, perhaps owing to the comparative thinness of the walls; and after careful preparation and staining, it seemed as if the slight inequality might be due to optical effect rather than to any real difference in the amount of thickening.

Fig. 1.

Stangeria paradoxa: surface view of the cuticle from the midrib of the upper side of the leaf, showing a stoma with characteristic thickening due to the dorsal (*d*) and ventral (*v*) lignified lamellæ of the guard-cells; *g*, the raised polar ends of the guard-cells. Note the absence of folds of the cutin in the neighbourhood of the stoma. $\times 450$.

Stomata occur scattered and irregularly-placed with regard to the pinna-axis, on the lower side of the lamina; they are generally absent in the very narrow areas over the lateral veins. They occur on both upper and lower mid-rib areas, their long axis being parallel with that of the pinna (text-fig. 1). The upper, as well as the lower stomata appear to be functional, as there is a well-developed air-space underlying each.

The stomata lie practically at the level of the epidermis (Pl. 17. figs. 1 & 5); the guard-cells are not sunk as in the case of *Zamia* (cf. Pl. 17. fig. 5 with Pl. 18. fig. 13). Each stoma is surrounded by from four to six epidermal cells having slightly thinner, straighter walls than the rest of the epidermis. These "subsidiary cells" are particularly marked in the case of the lower stomata, and may usually be distinguished into polar and lateral cells; Pl. 17. fig. 1 shows an example with two polar and two lateral subsidiary cells. Division of the lateral cells often takes place, as text-fig. 1 shows.

The cavities of the subsidiary cells in many cases contain crystals of calcium oxalate*. The stomata are elliptical in shape, and in surface view the lignified lamellæ† of the ventral and dorsal walls of the guard-cells give a very characteristic thickened appearance (Pl. 17. fig. 1; text-fig. 1) to the area overlying the guard-cells. This thickened area surrounding the pore of the stoma shows two thin patches at each pole, owing to the raising up of the polar ends of the guard-cells, bringing them nearer to the surface at these points, as described by Porsch for *Bowenia*. The projections of thickening substance between each pair of thin areas strengthening the ends of the stoma-opening are due to the extension of the ventral lignified lamellæ of the guard-cells (see text-fig. 26, p. 176). Laterally the cutin overlying the lamellæ shows fine striations (Pl. 17. fig. 1; text-fig. 1).

Hair-scars occur on the lower surface of the pinnæ, particularly on and near the midrib and at the margins. They are only to be seen here and there on the upper midrib. The scars are somewhat like a truncated cone in shape and are strongly cuticularised (Pl. 17. figs. 2 & 6).

The cutin layer overlying the epidermal cells is fairly thick, particularly on the upper surface of the pinnæ. It has closely-lying striations which are seen from sections to be due to folds (Pl. 17. fig. 3). These folds are particularly marked on the upper surface, and are irregular on the lamina, but are regular and run in a longitudinal direction on the midrib (text-fig. 1). They radiate from the poles of the stomata and from the bases of the hair-scars (Pl. 17. figs. 1 & 2; see also references quoted). Folding of the cutin is not to be found in any other genus of Cycads, and constitutes, with the undulate epidermal walls, a distinguishing feature of *Stangeria*.

CYCAS, Linn. Pl. 17. figs. 10-15; text-figs. 2 & 3.

REFERENCES:—Bornemann (1856), pp. 40, 42; Taf. 11. figs. 3-6. Kraus (1866), p. 31; Taf. 19. figs. 1-6; Taf. 20. figs. 10, 11. Strasburger (1866), Taf. 41. fig. 143. Mahlert (1885), Taf. 1. fig. 20. Nestler (1895), p. 343.

SPECIES EXAMINED:—*C. revoluta*, Thunb.; *C. circinalis*, Linn.; *C. undulata*, Desf.; *C. Micholitzii*, Dyer; *C. media*, R. Br.

The epidermal cells of *Cycas* are similar on both sides of the pinnæ, and are more or less irregular in shape and arrangement. They are usually somewhat elongated, and are three- to six-angular. Over the midrib the cells are more regular and rectangular in shape and run parallel with the pinna-axis. In these points there is general agreement with *Stangeria*. In *Cycas*, however, the lateral epidermal walls are not undulate, being at the most somewhat curved. A distinguishing feature of all species of *Cycas* examined‡ is the pitting of the outer walls of the epidermal cells (Pl. 17. figs. 10-13, fig. 15; text-fig. 2). In surface view and sections, these walls are seen to be perforated

* Cf. Kraus (1866), Pl. 23. fig. 29; the example figured in Pl. 17. fig. 1, does not show these crystals, owing to previous maceration.

† Porsch (1905); see also Karzel (1907).

‡ See also the lists given by Bornemann and Kraus.

by more or less regularly-disposed pores, which occur principally towards the periphery of the cells and which are covered by the cutin layer. This pitting occurs in both the upper and lower epidermis, and is not known to be present in any other genus of the Cycadaceæ*. The pores vary considerably in size in the different species of *Cycas*: the largest were observed in *C. circinalis*, the smallest in *C. Micholitzii*. Pitting of the lateral walls also occurs, but in a varying degree (Pl. 17. figs. 10 & 13). It is well marked in *C. revoluta*; in *C. undulata*, however, pits occur only here and there.

Fig. 2.

Cycas Micholitzii.—Lower cuticle in surface view: s, a stoma-opening, surrounded by six subsidiary cells. The thickened appearance round the stoma-opening is due to the showing-through of the cutin layer lining the air-space and overlying the guard-cells. $\times 450$.

Fig. 3.

Cycas revoluta.—A stoma in surface view, focussed below the level shown in Pl. 17. fig. 12, so that the dorsal or upper surfaces (*d*) of the guard-cells are seen surrounded by fifteen small cells, the intercalated series shown in section in Pl. 17. fig. 13. $\times 450$.

Stomata occur only on the underside of the pinnæ, and are scattered and irregular in position. As both Bornemann and Kraus have previously mentioned, the species of *Cycas* fall into two groups, according to their epidermal and stomatal characters, which, moreover, correspond with the external characters of the leaves. The first group contains species such as *C. revoluta*, with narrow pinnæ, the margins of which are inrolled †; the second group, of which *C. circinalis* is an example, contains species with broader, flatter pinnæ, less xerophytic in texture than those of the first group. All the types examined, with the exception of *C. revoluta*, belong to this group.

In the first type the epidermal cells are irregular in shape and disposition, particularly on the under side of the pinnæ, where, however, they are more isodiametric; their greatest diameter often lies transversely to the pinna-axis. The most distinctive feature of the cuticle consists in the complication of the stomatal apparatus. The guard-cells are well sunk below the surface, and between them and the subsidiary cells, which in this case overarch the guard-cells and form an "outer air-space," is an intercalated series consisting of from twelve to fifteen small cells (Pl. 17. fig. 13; text-fig. 3‡). These

* *Microcycas* has not yet been examined.

† Bornemann and Kraus have also examined *C. inermis* in this group.

‡ Cf. Kraus (1866), Taf. 19. figs. 4 & 5.

intercalated cells appear to be derived from the same mother-cells as the subsidiary cells, which are also from twelve to fifteen in number. The subsidiary cells are narrowly and obliquely elongated (Pl. 17. fig. 13), so that they arch over the depression containing the guard-cells in such a way as to give the appearance, in surface view, of a somewhat hemispherical elevation, perforated by a small rounded opening (Pl. 17. fig. 12)*. The intercalated cells and subsidiary cells are pitted in the same way as the general epidermal cells (Pl. 17. fig. 13).

In the second group, the epidermal cells are irregular in form and position, as in the first type; the stomata, however, are more simple in organisation (Pl. 17. fig. 14). They are a little sunk below the general level of the epidermis, but there is no intercalated series and the subsidiary cells do not overarch the depression; hence the very different appearance of the stomata as seen in surface view (text-fig. 2). The subsidiary cells are fewer in number, though very variable, usually between five and nine (text-fig. 2); in shape they are from isodiametric to elliptical.

In *C. circinalis* and *C. revoluta*, fairly young stages were examined; these showed very distinctly the presence of hair-scars on the under surface of the leaves.

The cutin layer is in all species fairly thick; it is not characterised by any special foldings or markings, as in *Stangeria*.

ENCEPHALARTOS, Lehm. Pl. 18. figs. 10 & 11; text-figs. 4-8.

REFERENCES:—Bornemann (1856), pp. 40, 44; Taf. 13. figs. 1-6. Kraus (1866), pp. 33, 34; Taf. 20. figs. 12-15; Taf. 21. fig. 16. Mahlert (1885), Taf. 1. fig. 23. Nestler (1895), pp. 346-348; Taf. 11. figs. 5-8; Taf. 12. fig. 9.

SPECIES EXAMINED:—*E. villosus*, Lemaire; *E. Woodii* (? hort.); *E. horridus*, Lehm.; *E. caffer*, Miq.; *E. lanuginosus*, Lehm.; *E. Lehmanni*, Eckl.; *E. Ghellinckii*, Lem.; *E. Laurentianus*, Wilden. (See also the lists of Bornemann and Kraus.)

Reviewing the species examined as a whole, there seems to be a considerable amount of variation in the form of the epidermal cells, even in the cells of the same pinna (text-fig. 5). Generally speaking, the cells are somewhat elongated and the longitudinal axes run in the direction of the longitudinal axis of the pinna (Pl. 18. fig. 11; text-fig. 4). The transverse walls are usually obliquely placed with regard to the pinna-axis; the cells are three- or four-angular and rhombic in shape.

In *E. Lehmanni* (text-fig. 5), *E. caffer*, and *E. lanuginosus*, where the cells are small and more nearly isodiametric, their greater diameter is usually transversely or obliquely placed with regard to the axis of the pinna; in *E. horridus* the cells are irregular in form and direction, particularly in the region of the lobe-bases.

The epidermal cells of the lower surface are of two kinds—those of the nerve-courses and those of the stomatal areas. The cells of the nerve-courses are similar in form to those of the upper surface. They are usually somewhat narrow and elongated and are arranged in longitudinal lines. The cells of the stomatal areas are typically shorter, broader, and more isodiametric.

* Cf. Bornemann (1856), Taf. 11. fig. 4.

The walls of all the epidermal cells are without undulations, but are often slightly curved. In most of the cases examined, the lateral walls (*never* the outer walls) of both upper and lower epidermal cells are pitted in the same way as the lateral walls of *Cycas*. This is very marked in *E. Lehmanni* (Pl. 18. fig. 10; see also fig. 11). In *E. Woodii* and on the lower surface of *E. villosus* pittings are infrequent. In the examined species of *Encephalartos* the inner and lateral walls of the epidermal cells are often slightly

Fig. 4.

Encephalartos Woodii.—Upper cuticle in surface view, showing form and arrangement of epidermal cells.
× 450.

Fig. 5.

E. Lehmanni.—Upper cuticle in surface view.
× 450.

thicker than the outer walls (Pl. 18. fig. 10) Towards the pinna-margins the cells become smaller and thicker-walled. In some cases, as in *E. horridus* and *E. villosus* (Pl. 18. fig. 11), the elongated epidermal cells are septate.

Stomata are confined to the lower surface of the pinnæ, and more or less definitely to the areas between the nerve-courses. Their longitudinal axis is typically coincident with that of the pinna. They are in most cases considerably sunk below the general level of the epidermis without overarchng of the subsidiary cells; so that the outer air-space is fairly long and narrow and tubular in shape (text-fig. 6). There is a somewhat irregular system of intercalated cells separating the guard-cells from the general epidermal cells. In median sections of the stomata, typically two intercalated cell-series appear (text-fig. 6); the sections may not, however, pass through a cell of both series. Nestler (1895, p. 348) mentions that one or two intercalated series are present; while according to Kraus (1866, p. 34) the guard-cells lie immediately under the epidermis except at the pinna-bases. The stoma-opening is somewhat elongated and elliptical to quadrangular in shape (text-

figs. 7 & 8). It is surrounded by from four to six subsidiary cells, and is strongly cuticularised.

Simple hairs and hair-scars occur in *E. Ghellinckii*. In *E. caffer* crystals of calcium

Fig. 6.

Encephalartos Lehmanni.—Section of the lower cuticle, showing a stoma in median section. Note the presence of intercalated cells between the guard-cells and subsidiary cells, and the heavy cuticularisation. $\times 450$.

Fig. 7.

E. Lehmanni.—Lower cuticle (after maceration) in surface view, showing a stoma-opening (*s*) surrounded by six subsidiary cells (*p*, polar subsidiary cells; *l*, lateral subsidiary cells). $\times 450$.

Fig. 8.

E. villosus.—Lower cuticle in surface view; the stoma-opening is surrounded by five subsidiary cells, one of the polar cells being divided. $\times 450$. This cuticle was not removed by maceration, so that the thickening of the guard-cells and the lining of the outer air-space show through in the surface view, giving a much thickened appearance to the stoma-opening.

oxalate, such as figured by Nestler (1895, Taf. 11. figs. 6–8), were not observed; very little material of this species was, however, available.

The cutin layer is of a considerable thickness in the case of both upper and lower epidermises (Pl. 18. fig. 10; text-fig. 6).

DIOON, Lindl. (Pl. 18. figs. 5-9; text-figs. 9 & 10.)

REFERENCES :—Bornemann (1856), pp. 40, 45; Taf. 12. figs. 11-14. Kraus (1866), pp. 39, 40; Taf. 22. figs. 21-23. Mahlert (1885), Taf. 1. fig. 22. Porsch (1905), p. 11; Taf. 1. fig. 7.

SPECIES EXAMINED :—*D. edule*, Lindl.; *D. spinulosum*, Dyer.

The two species were found to be of a similar type, characteristic for the genus. In *D. edule* the upper epidermal cells and those of the nerve-courses of the lower side of the pinnæ are similar, and are of two kinds—thick-walled cells elongated fairly regularly in the direction of the longitudinal pinna-axis, and more or less isodiametric thin-walled cells. These two kinds of cells may be distinguished in transverse section (Pl. 18. fig. 7). The thin-walled cells appear to be the more numerous. The cells of the stomatal regions between the nerve-courses are usually smaller than elsewhere. They are almost isodiametric and are uniformly thick-walled. In *D. spinulosum* the epidermal characters are similar to those in *D. edule*, though all the cell-walls seem to be slightly less thickened. In both cases the lateral walls of the cells are often slightly pitted (Pl. 18. fig. 5) (Kraus, 1866, fig. 21), and all the walls are more or less heavily layered

Fig. 9.

Dioon edule.—Lower cuticle in surface view, showing six subsidiary cells surrounding a stoma-opening, *s*. Cf. the epidermal cells of the nerve-course (*n*) with those of the upper surface shown in Pl. 18. fig. 5. $\times 450$.

Fig. 10.

D. spinulosum.—A stoma in median section, showing the dorsal (*d*) and ventral (*v*) lignified lamellæ of the guard-cells, and three intercalated series of cells (*i¹-i³*) between the guard-cells and subsidiary cells. (Cf. Pl. 18. fig. 6.) $\times 450$.

(Pl. 18. figs. 5 & 9). In the case of the thick-walled cells of *D. edule*, the lumen is often almost filled up.

Stomata occur very definitely limited to the areas between the nerve-courses on the under surface of the leaves. They are much sunk in flask-shaped cavities, which are slightly deeper in *D. edule* than in *D. spinulosum*. In both species there are typically three intercalated series of cells between the guard and subsidiary cells (Pl. 18. fig. 6;

text-fig. 10) (Porsch, 1905, Taf. 1. fig. 7). The cavities are lined with a layer of cutin, and the epidermal cells surrounding the stoma-openings (subsidiary cells) arch over slightly. They are also a little raised beyond the surface of the pinna, and are rounded, as seen in surface view. They are from six to eight in number and are much thickened, especially in *D. edule* (text-fig. 9).

Crystals of calcium oxalate occur in *D. edule* on the lower side of the pinnæ in the nerve-courses. They were not observed on the upper side. Kraus figures a crystal from the lower epidermis, but does not specially note their occurrence there. He mentions that "here and there a thin-walled cell contains a crystal." So far as could be determined, however, from the appearance in surface view and in section, the crystals occur in cavities or spaces between the cells rather than in the cells themselves (Pl. 18. figs. 8 & 9); for in surface view the rounded ends of the neighbouring cells seem to project into a space, and in section Scharlach R shows the presence of a layer of cutin entirely surrounding or lining the crystal-containing space.* *D. spinulosum* has no crystals.

The cutin layer in both species is very thick, especially in *D. edule* (Pl. 18. figs. 7 & 8).

CERATOZAMIA, Brongn. Pl. 18. figs. 1-4; text-figs. 11 & 12.

REFERENCES:—Bornemann (1856), pp. 40, 45; Taf. 12. figs. 7-10. Kraus (1886), pp. 38, 39; Taf. 21. figs. 17-20. Mahlert (1885), Taf. 1. fig. 21.

SPECIES EXAMINED:—*C. mexicana*, Brongn.; *C. fusco-viridis*, D. Moore; *C. latifolia*, Miq.

These three species show the same general characters, all possessing two kinds of epidermal cells, thick- and thin-walled. The thick-walled cells apparently predominate in *Ceratozamia* (cf. *Dioon*), though they are somewhat variable in number and in distribution in the three species examined. There is considerable variation, moreover, in the relative thickness of the thick-walled cells; *C. mexicana* shows a very marked contrast between the two kinds of cells (Pl. 18. figs. 1 & 3); *C. latifolia* shows comparatively little difference between them (text-fig. 11); while *C. fusco-viridis* is intermediate in this respect. The cells—particularly the thicker-walled ones—are much elongated in the direction of the longitudinal axis of the pinna. The cells of the nerve-courses, in the lower epidermis, are similar to those of the upper side; while between the nerve-courses, as is usual, the cells are shorter and broader. Thin-walled cells occur in these areas also, especially in *C. mexicana* (cf. *Dioon*).

In *C. latifolia*, the elongated thick-walled cells of both upper and lower epidermises are often septate (text-fig. 11); Kraus notes the same fact in *C. Küsteriana*.

C. mexicana shows pitting on the lateral walls of the thickened cells (Pl. 18. figs. 1 & 2). This is more marked on the lower surface than on the upper, especially in the case of the subsidiary cells of the stomatal apertures. In *C. fusco-viridis* also, the lower epidermal cells are much pitted; in *C. latifolia*, however, pitting was not noted, except

* A similar occurrence of crystal-containing cavities is quoted by Nestler in the case of *Encephalartos caffer*. In *D. edule* the cavities are apparently larger than in that species.

in the cells of the nerve-courses. The cell-walls, particularly in *C. mexicana*, are very strongly layered (Pl. 18. figs. 1-4). *Ceratozamia* has wide stomatal regions and narrow nerve-courses, the demarcation of the areas being most distinct in *C. latifolia*. The stomata lie parallel to the pinna-axis, or somewhat obliquely to it, and are sunk below the level of the epidermis, having one intercalated series of cells (Pl. 18. fig. 4). The air-space is lined with a layer of cutin, extending as is usual to the ventral walls of the guard-cells, and its somewhat oval opening is surrounded by from four to six subsidiary cells; five were observed in *C. latifolia* (text-fig. 12), six in *C. mexicana* (Pl. 18. fig. 2),

Fig. 11.

Ceratozamia latifolia.—Upper cuticle in surface view, showing thin-walled and thick-walled cells. At *x*, an elongated cell is divided by a septum. $\times 450$.

Fig. 12.

C. latifolia.—Lower cuticle (after maceration) in surface view, showing a stoma-opening (*s*) surrounded by five subsidiary cells (*p* and *l*). $\times 450$.

and usually four in *C. fusco-viridis*. Kraus notes four in *C. Küsteriana*. The subsidiary cells do not overarch the air-space to any extent.

The cutin layer is throughout fairly thick, particularly on the upper surface of the leaves (Pl. 18. figs. 3 & 4).

MACROZAMIA, Miq. Pl. 18. figs. 14 & 15; text-figs. 13-18.

REFERENCES :—Bornemann (1856), pp. 40, 44; Taf. 11. figs. 16, 17. Schwendener (1881), fig. 13. Nestler (1895), pp. 352, 354; Taf. 12. figs. 10, 11; Taf. 13. fig. 16.

SPECIES EXAMINED :—*M. spiralis* (= *Fraseri*, Miq. ?); *M. corallipes*, Hook. f.; *M. Denisonii*, F. Muell.; *M. Hopei*, T. Hill; *M. cylindrica*, C. Moore; *M. Macleayi*, Miq. (= *spiralis*, Miq. ?).

The six species examined show agreement in the occurrence of thick-walled and thin-walled epidermal cells. They fall into two groups, however, according to the arrangement of the cells. *M. Macleayi* (= *spiralis* ?), *M. Denisonii* (text-fig. 13), and *M. Hopei* (text-fig. 14) differ from the other three species in the irregularly-shaped three- or four-angular cells placed transversely or obliquely to the long axis of the pinna. In this first group, moreover, the thick-walled cells are very marked, their cavities being often almost filled up; there is also a heavy primary wall, particularly in *M. Denisonii* (text-fig. 13). In *M. corallipes*, *M. cylindrica* (Pl. 18. fig. 15), and *M. spiralis* (= *Fraseri* ?) the cells are three- or more often four-angular; they are generally elongated in the direction of the longitudinal axis of the pinna, though the thinner-walled cells are often squarish. In this second group, the walls are often much pitted, particularly in the case of the thin-walled cells (Pl. 18. fig. 15). In the other type pitting is less

Fig. 14.

Fig. 13.

Macrozamia Denisonii.—Upper cuticle in surface view, showing thin-walled and thick-walled cells, and the heavy primary cell-walls. Note the irregular arrangement of the cells. $\times 450$.

M. Hopei.—Upper cuticle in surface view, showing thin-walled and thick-walled cells, and pitting of the thin-walled cells. $\times 450$.

evident. Layering of the cellulose is frequently marked, as in *M. cylindrica* (Pl. 18. fig. 15). The cells of the upper surface and of the nerve-courses of the lower surface are similar in both groups, but the cells between the nerve-courses are usually irregular in shape and size, and are much more uniformly thickened (text-figs. 15 & 16), although occasionally thin-walled cells occur. Sometimes, as in *M. corallipes*, the elongated thickened cells are septate.

Stomata occur only on the lower surface between the nerve-courses; their long axis is parallel with or slightly oblique to the long axis of the pinna. A regular longitudinal arrangement, almost in lines, is seen in *M. spiralis* (= *Fraseri* ?). In all cases the stomata are sunk below the epidermis, an apparently rather irregular system of cells

intervening between the guard-cells and the subsidiary cells. There seem to be typically two series of these intercalated cells (Pl. 18. fig. 14), but the number of series as seen in section through a stoma shows a little variation, apparently owing to irregularity in position of the cells; this is seen in text-fig. 17. In *M. cylindrica* the outer air-space is long, and narrower towards the opening, owing to the slight overarching of the subsidiary cells, which are slightly raised above the level of the epidermis (Pl. 18. fig. 14). The opening as seen in surface view is oval or elliptical in shape. In *M. Denisonii* the outer air-space is shallower and wider (text-fig. 17). In all cases the air-space is fairly heavily lined with cutin. The subsidiary cells vary from five or six to eight in number, but are typically six, as in *M. Denisonii* (text-fig. 15) and *M. Hopei* (text-fig. 16).

Fig. 15.

Fig. 16.

Fig. 17.

Macrozamia Denisonii.—Lower cuticle in surface view (after maceration), showing stoma-opening surrounded by six subsidiary cells. $\times 450$.

M. Hopei.—Lower cuticle in surface view, showing a stoma-opening with six subsidiary cells (*p* and *l*). The dorsal surfaces of the guard-cells are indicated at *d*. $\times 450$. (This cuticle was not removed by maceration, so that all the thickening of the air-space and guard-cells shows through.)

M. Denisonii.—Lower cuticle, showing a section through a stoma. Note the irregular intercalated cells, between the guard and subsidiary cells. $\times 450$.

In *M. cylindrica*, in the nerve-courses of the lower surface, and also here and there on the upper surface of the pinnæ, occur crystals of calcium oxalate (text-fig. 18); they appear to be contained in small isodiametric cells, and not in cavities, as in *Dioon edule*, and in *M. spiralis* (= *Fraseri* ?), where the crystal-containing spaces are very small and not at all frequent.

Fig. 18.

Macrozamia cylindrica.—Lower cuticle (from a nerve-course) in surface view, showing a crystal-containing cell. (Cf. the form and arrangement of the cells with those shown in Pl. 18. fig. 15, from the upper epidermis of *M. cylindrica*.) $\times 450$.

The cutin layer is moderately thick (Pl. 18. fig. 14; text-fig. 17), but, as usual, it is thinner on the lower than on the upper surface of the leaves.

ZAMIA, Linn. Pl. 18. figs. 12 & 13; text-figs. 19–23.

REFERENCES :—Bornemann (1856), pp. 40, 43, 44; Taf. 11. figs. 7–15. Kraus (1866), pp. 41, 42; Taf. 22. figs. 24–27.

SPECIES EXAMINED :—*Z. Skinneri*, Warsz.; *Z. muricata*, Willd.; *Z. Loddigesii*, Miq.; *Z. integrifolia*, Ait.; *Z. Fischeri*, Miq.; *Z. Ottonis*, Miq.; *Z. Lindeni*, Regel, = *Z. cylindrica*, Liebm. (= *Leiboldii*, Miq.?).

These eight species agree in essential characters of the epidermis. The cells in all are elongated in the direction of the longitudinal axis of the pinnæ, and are three- or four-angular in shape. In *Z. Ottonis* the cells are somewhat irregular, with groups of smaller cells here and there. In some species, for example *Z. Skinneri* (text-figs. 20 & 23) and *Z. integrifolia* (Pl. 18. fig. 12), there is a marked distinction into two kinds of cells, thin-walled and thick-walled. In *Z. Skinneri* and *Z. Loddigesii*, the thin-walled cells are only occasional, while the other cells often show a considerable amount of thickening. In *Z. Fischeri* (text-fig. 19) and *Z. Ottonis* the walls throughout are fairly thin. *Z. muricata* has walls of intermediate thickness. The cells of the nerve-courses of the lower epidermis are similar to those of the upper surface. In *Z. cylindrica* and *Z. Lindeni* the cells of the nerve-courses are fairly thick-walled, elongated

and narrow, running quite straightly, and having somewhat pointed ends. The cells between the nerve-courses are shorter and broader and more irregular, but they are still slightly elongated, except in the immediate neighbourhood of the stomata. In the stomatal areas the cell-walls are sometimes thinner than they are on the upper surface; this is best seen in *Z. Skinneri* and *Z. Lindeni*. In *Z. Fischeri* the upper epidermal cells and those of the nerve-courses and stomatal areas are of about the same thickness.

Pittings are sometimes plainly seen (Pl. 18. fig. 12), particularly in the case of the thin-walled cells of nerve-courses. Occasionally the cellulose of the epidermal cells is layered; and in some cases very much elongated cells are septate as in *Z. Ottonis* and *Z. Skinneri*.

Fig. 19.

Zamia Fischeri.—Upper cuticle in surface view.
× 450.

Fig. 20.

Z. Skinneri.—Upper cuticle in surface view. Cf. the amount of variation in the thickness of the cell-walls with that shown in text-fig. 19. × 450.

Stomata occur on the lower sides of the leaves in definite areas between the nerve-courses; they are sunk only slightly below the level of the epidermis, there being no intercalated series of cells between the guard-cells and subsidiary cells (Pl. 18. fig. 13). The lignified lamellæ of the guard-cells may be seen, as in *Stangeria* (and *Bowenia*), on focussing slightly below the level of the subsidiary cells. The subsidiary cells are characteristic in number and arrangement; there are from four to six, consisting of an elongated cell at each pole of the stoma, and two (sometimes only one) lateral cells at each side (text-fig. 22). Sometimes a second series of lateral cells is present (text-fig. 21).

Fig. 21.

Zamia Lindenii.—Lower cuticle in surface view. The thickened stoma-opening shows a double series of lateral subsidiary cells at each side (*l*, *l*). $\times 450$. (This cuticle was not removed by maceration, and thus shows the thickening substance of the guard-cells and air-space, showing through to the surface.)

Fig. 22.

Z. muricata.—Lower cuticle in surface view. The stoma-opening in this case has five subsidiary cells (*p* and *l*) surrounding it. $\times 450$. (Not macerated; see text-fig. 21.)

Fig. 23.

Z. Skinneri.—Upper cuticle in section, showing the variation in the thickness of the cell-walls. (Cf. text-fig. 20.) $\times 450$.

In *Z. integrifolia* hair-scars are occasionally seen, particularly on the under surface in the nerve-courses. The scars seem to occur over a group of the thinner-walled cells. They occur also in *Z. Ottonis* and *Z. Skinneri*.

The cutin layer is similar in thickness to that of *Macrozamia* (cf. Pl. 18. figs. 13 & 14).

BOWENIA, *Hook.* Pl. 17. figs. 7-9; text-figs. 24-26.

REFERENCES:—Nestler (1895), pp. 356, 357; Taf. 13. figs. 19-21. Porsch (1905), p. 13; Taf. 1. figs. 1-4.

SPECIES EXAMINED:—*Bowenia spectabilis*, Hook., var. *serrulata*, André (= *Bowenia serrulata* (André), Chamberlain (1912, p. 419)).

Bowenia serrulata agrees with such *Zamias* as *Z. Fischeri* in the arrangement of the three- or four-angular epidermal cells and also in the thickness of their walls (cf. text-figs. 19 & 25). The cells are elongated, with straight walls, and are slightly narrower

over the nerve-courses, even on the upper surface of the leaf. As is usual, on the lower surface, the cells between the nerve-courses are shorter and broader. No pitting of the walls was observed.

Stomata occur in both the upper and the lower epidermis (Pl. 17. figs. 7 & 8). They are few and scattered on the upper side, occurring principally in the middle of the lower half of the pinnæ, between the veins; they decrease in number towards the upper half and margins. They are fairly abundant on the lower side. In all cases their long axis is approximately parallel with that of the pinnæ. The guard-cells are situated practically at the level of the epidermis (Pl. 17. fig. 9), there being no intercalated series of cells between them [and the subsidiary cells—they are less sunk than those of any other genus except *Stangeria* (cf. Pl. 17. fig. 9, *Bowenia*, with fig. 5, *Stangeria*;

Fig. 24.

Bowenia spectabilis var. *serrulata*.—Upper cuticle in section. $\times 450$.

Fig. 25.

B. spectabilis var. *serrulata*.—Upper cuticle in surface view. $\times 450$.

fig. 14, *Cycas circinalis*; and Pl. 18. fig. 13, *Zamia integrifolia*, for degrees of sinking of guard-cells without an intercalated series of cells). The stomata are surrounded by from four to six subsidiary cells, the walls of which are fairly thin (Pl. 17. figs. 7 & 8), particularly in the case of the stomata of the upper leaf-surface. On account of the position of the guard-cells, *i.e.* almost at the general level of the epidermal cells, the lignified lamellæ of dorsal and ventral walls are plainly seen, giving a characteristic appearance in surface view (Pl. 17. figs. 7 & 8; cf. *Stangeria*, fig. 1). As previously described by Porsch in the case of *B. spectabilis**, the raising-up of the polar ends of the guard-cells is indicated in surface view by the presence, at each pole of the stoma, of two

* Porsch (1905), p. 13; Taf. 1. figs. 1 & 3. See also Nestler (1895), Taf. 13. figs. 19 & 20, b.

thin pear-shaped areas in the thickened area overlying the guard-cells. The projections of thickening material formed by the extension of the ventral lignified lamellæ are seen at the poles of the stoma (see text-fig. 26, showing the correlation of serial sections through a stoma with the surface view). The cutin layer overlying the lamellæ is faintly striated as in the case of *Stangeria* (Pl. 17. figs. 7 & 8).

Neither hair-scars nor crystal cavities were observed. The cutin layer, compared with that of the other genera, is not very thick (Pl. 17. fig. 9; text-fig. 24).

(iii.) General Summary of the Structure of the Recent Cuticles.

The epidermal cells of Cycad leaves have usually slightly curved or straight walls. *Stangeria* has undulate walls somewhat like those of ferns*. The walls generally attain a considerable thickness, and often show layering as in *Ceratozamia*, *Dioon*, and some *Macrozamia*s. *Bowenia* has comparatively thin walls, while *Ceratozamia* and *Dioon* on the whole have the thickest, though these are characterised by the possession of two kinds of epidermal cells, some elongated and thick-walled, and others shorter and thin-walled. A variation in the thickness of the walls occurs also in *Macrozamia*, some *Zamia*s, and to a certain extent in *Encephalartos*. The epidermal cells are usually elongated, and in *Cycas* and *Stangeria* are alike on both upper and under sides of the leaf. In other types the cells of the under side following the course of the veins are similar, or nearly similar, to those of the upper side, while the general epidermal cells between the "nerve-courses" are more isodiametric. Usually, in the upper epidermis, the nerve-courses are not marked by any difference in the form of the epidermal cells; *Stangeria* shows a slight elongation and narrowing of the cells above the veins.

Stomata occur generally only on the under side of the leaves between the nerve-courses, their long axes coinciding or forming a slight angle with the long axis of the pinnæ. The occurrence of stomata in these definite areas seems to be correlated with the above mentioned difference in the form of the epidermal cells there, as compared with those of the nerve-courses (see *Zamia*, *Macrozamia*, *Ceratozamia*, *Dioon*, and *Encephalartos*). In *Cycas* and *Stangeria* the stomata are scattered and irregularly placed; there are no fixed stomatal areas.

Bornemann (1856, p. 42) has recorded the occasional occurrence of stomata on the upper side of the pinnæ in *Zamia muricata* var. *picta*. Kraus (1866, fig. 15) figures a stoma of the pinna-base in *Encephalartos Altensteinii*; it is apparently from the upper side, though he does not mention the occurrence of stomata other than on the under side of the pinnæ. Nestler (1895, p. 362) was unable to find such stomata in *Encephalartos*; in *Bowenia*, however, he records them from the basal parts of the upper side of the pinnules, gradually decreasing in number and finally disappearing towards the middle. The present investigations have revealed the presence of stomata on the upper side of the pinnæ in *Stangeria*. They are, however, confined to the midrib (text-fig. 1).

* Bornemann (1856), p. 42; Kraus (1866), p. 13.

Nestler (1895, pp. 360, 364) gives a table showing the numbers of stomata per square millimetre in a representative selection of species. The *Ceratozamia*s (with from 24 to 31 per sq. mm.) and the *Macrozamia*s (29 to 31) are seen to have the fewest, and the *Dioon*s (*D. edule*, 81, and *D. angustifolia*, 72) the most numerous stomata. The investigated species of *Cycas* (with from 49 to 56 per sq. mm.), *Encephalartos* (41 to 52), *Stangeria* (54), and *Zamia* (43 to 60) have very similar numerical distribution of stomata, and are intermediate between the *Ceratozamia*s and *Macrozamia*s on the one hand and the *Dioon*s on the other. *Bowenia spectabilis* has 35 stomata per sq. mm. on the under side and 14 on the upper side of the pinnae.

The stomata of the Cycads as a group are larger than those of any other group of Gymnosperms*. Their size seems to be fairly constant, being about 0.075×0.034 mm., while their structure conforms to a general plan—the “Gymnosperm type” of Porsch. The guard-cells in median and polar section show a constant form and position relative to the general epidermal level.

The ventral and dorsal walls of the guard-cells have in every case lignified lamellae†, though these are slightly variable in extent and in clearness of definition‡. They give to those stomata, the guard-cells of which are only slightly sunk, a very characteristic thickened appearance in surface view (Pl. 17. fig. 1, *Stangeria*; figs. 7 & 8, *Bowenia*; text-fig. 26, *Bowenia*). In *Bowenia*, for example, the surface view of a stoma shows the pore bounded on each side by a somewhat crescentic structure formed by the lignified lamellae on the upper or dorsal surface of the guard-cell. This is covered by a layer of cutin. The lamellae on the lower or ventral surface of the guard-cells are only seen at the poles of the stoma, where the guard-cells are raised and the lamellae are extended, appearing as thickened projections. Text-fig. 26 shows the correlation of serial sections through a stoma with the surface view. The gradual raising of the guard-cells towards the poles and extension of the ventral lamellae with accompanying diminution of the dorsal lamellae are plainly seen. In the case of guard-cells which are much sunk, the surrounding, and usually overarching, epidermal cells obscure the characteristic appearance caused by the lignified lamellae. In intermediate conditions such as exhibited by *Zamia* (Pl. 18. fig. 13), the lamellae of the guard-cells may be seen on focussing slightly below the epidermal cells. They are quite comparable with those of *Bowenia* and *Stangeria*.

The epidermal cells surrounding the guard-cells and stoma-opening—“subsidiary cells”—are usually fairly constant in number and arrangement in a genus. In no case was material available young enough to trace the entire development of the guard-cells and subsidiary cells from an epidermal mother-cell. Four seems to be the original number of subsidiary cells, the typical arrangement of these being one at each pole of the stoma and one at each side. It seems possible from a comparison with *Gnetum* and *Welwitschia*§, together with a consideration of fairly young examples in *Cycas revoluta*

* Malhert (1885), p. 57.

† These are not figured in every case; they are shown in text-figs. 10 & 26 and in Pl. 18. figs. 4 & 6.

‡ Karzel (1908). § Takeda (1913¹), p. 351; & (1913²), pp. 365, 366, text-fig. 8.

and *Ceratozamia mexicana*, that the polar subsidiary cells represent ordinary epidermal cells, while the lateral ones are the product of division of the epidermal mother-cell giving rise to the guard-cells also. Frequently the lateral cells are transversely divided into two, so that six subsidiary cells—two polars and four laterals—are seen. Occasionally only one of the lateral cells is divided; division of the polar

Fig. 26.

Bowenia spectabilis var. *serrulata*.—A, surface views of stomata (cf. Pl. 17. figs. 7 & 8);

B, 1-4, serial sections through a stoma, corresponding to the positions 1-4 in the surface view. These sections show the gradual raising of the guard-cells towards the polar ends of the stoma, and the change in size and position of the dorsal (*d*) and ventral (*v*) lignified lamellæ of the guard-cells, these changes giving the characteristic surface view of the stoma. B, 5, shows a polar section of a stoma in which the dorsal lamellæ do not appear; this section corresponds to position 5, in the surface view. (Diagrammatic.)

cells is comparatively infrequent. An increase in the number of subsidiary cells is often seen where guard-cells are much depressed below the general level of the epidermis, there being a mechanical necessity for this increase—the deeper the outer air-space, the greater the number of cells required to ensure its rigidity, by means of their lateral walls.

In *Bowenia* and *Stangeria*, where the guard-cells are not sunk, there are from four to six subsidiary cells, differentiated into polar and lateral (text-fig. 1; Pl. 17. figs. 1, 7, & 8). Occasionally there is some displacement of the cells, so that the differentiation is not always quite definite (text-fig. 1; Pl. 17. fig. 8). *Encephalartos* (text-figs. 7 & 8), *Ceratozamia* (text-fig. 12; Pl. 18. fig. 2), *Macrozamia* (text-figs. 15 & 16), and *Zamia* (text-figs. 21 & 22) typically show from four to six subsidiary cells, differentiated into polars and laterals. An interesting distinction in *Zamia* is the frequent presence of a second series of lateral subsidiary cells, parallel with the first, due perhaps to the longitudinal division of the original lateral cells*; this is well seen in *Zamia Lindenii* (text-fig. 21). The subsidiary cells in these four genera are at a different level from that of the guard-cells, the latter being sunk to a greater or less degree, forming an outer air-space. The opening of the air-space at the surface of the leaf is elliptical to quadrangular in shape. In some cases, particularly in some *Macrozamia*s and some species of *Encephalartos*, there is a certain amount of overarching of the subsidiary cells. In cases where the guard-cells are much sunk, the opening of the air-space on the leaf-surface is more rounded in shape, and the subsidiary cells, which tend to increase in number for the better strengthening of the large air-space, are not distinguished into polars and laterals. This is seen in *Dioon* and *Cycas revoluta*, where all the cells are of the same shape. Occasionally also the subsidiary cells are raised beyond the general level of the epidermis, forming small elevations perforated by the stoma-openings (Pl. 17. figs. 12 & 13, *Cycas revoluta*).

In some cases crystals of calcium oxalate occur in the epidermal layers; in *Stangeria* they are to be found in the cells round the stomata; in *Macrozamia cylindrica* crystals occur on the upper leaf-surface and on the lower surface in the cells of the nerve-courses. In other cases, e. g. *Dioon edule* and *Macrozamia spiralis* (= *Fraseri*?), they occur in spaces between the epidermal cells.

Hair-scars occur in several cases, such as *Encephalartos Ghellinckii*, *Cycas circinalis* and *C. revoluta*, *Zamia integrifolia* and *Stangeria*.

The cutin layer overlying the epidermal cells is fairly thick—usually thicker on the upper than on the under side of the leaf. *Dioon*, *Ceratozamia*, and some species of *Encephalartos* show the heaviest cutin layer, while that of *Bowenia* is comparatively thin. The cutin of *Stangeria* has very characteristic folds.

III. CUTICLES OF FOSSIL CYCADEAN FRONDS.

(a) Previous Observations on the Cuticular Structure of Fossil Cycadean Fronds.

Bornemann, whose work has been mentioned above, was the first to make observations upon the structure of the cuticles of fossil fronds, and, in fact, made a comparative study, based on the recent forms, of a type which has never since been attempted (1856). The

* Cf. Takeda (1913¹), p. 351; (1913²), pp. 365, 366; text-fig. 8.

fossil remains with which he had to deal were, however, very fragmentary; little or nothing could be made out as to the macroscopic characters of the fronds whose cuticular structure is figured, and from his figures one is led to suppose that the preservation of the fragments or the methods of treatment were not of the best. In the cases figured the walls of the epidermal cells are all represented as straight, and the stomata figured do not allow any close comparison with the examples now before us. It would appear that the forms figured on his Taf. 7. figs. 1-6 are allied to *Thinnfeldia*, but in no other cases are the stomata sufficiently well represented.

Schenk, in his Fossil Flora of Frankonia (1867), made a great advance in the study of cuticles; he described and gave accurate figures of the cuticles of a number of forms, including *Teniopteris stenoneura*, *Thinnfeldia* spp., *Nilssonina polymorpha*, *Otozamites* (*Otopteris*) *Bucklandi*, *Pterophyllum* spp., several conifers and ferns. He does not give many good figures of preparations showing stomatal structure, but he was the first to observe the sinuous walls of the cells in various species of *Pterophyllum* and in *Otozamites*. He mentions that he has made preparations from the English species *O. acuminatus*, *O. Beani*, and *O. graphicus*, also of the English species of *Teniopteris*, which was found to be comparable with the Otozamitean structures*. He gave some short descriptions of stomata, noticing that they occur usually on one side of the lamina only; but there are many points of importance on which we can obtain no information from his work. The same author subsequently figured the cuticles of *Dioonites Dunkerianus* (Goep.)†.

Zeiller (1882) described and figured the epidermal structure of a *Thinnfeldia* (*Cycadopteris Brauniana*) and *Frenelopsis Hoheneggeri*; he has also figured a small piece of a cuticle of a Palæozoic *Pterophyllum* (Zeiller, 1906).

In his 'Jurassic Flora of the Yorkshire Coast' Prof. Seward (1900, p. 234, fig. 43) has given a good figure of part of a lower cuticle of *Ctenis* sp. The form of the epidermal cells, often bearing papillæ, was well shown; the stomata were not clearly seen, but there were depressions representing the cavities in which the stomata were sunken.

Most of the recent observations on the epidermal structures of Mesozoic Cycadean fronds have been made by Prof. Nathorst. He has described and figured the cuticles of species of the following genera:—*Dictyozamites*, *Pseudocycas*, *Ptilozamites* (*Ctenozamites*), and *Nilssonina*‡. He has given many excellent photographs of these, but has not yet paid special attention to the structure of the stomata, which is one of the most important features from the point of view of the present inquiry. As will be seen later, many of the cuticles now described and figured can be very closely compared with the specimens from the Scandinavian plant-beds. This is especially the case in the instance of *Dictyozamites Johnstrupi* from Bornholm and *Dictyozamites Hawelli* from Marske. The close similarity between these forms, and also the similarity between specimens of other types occurring in both the Lower and Middle Estuarine beds of different parts of North

* Schenk (1867), p. 165.

† Schenk (1871), pl. xv. figs. 2-4.

‡ Nathorst, Paläobotanische Mitteilungen i.-iv., 1907-8; and 'Ueber die Gattung *Nilssonina*,' 1909.

Yorkshire, indicate that the cuticular structures are not essentially affected by purely local conditions, but show characters which are related to the more general conditions and also to questions of phylogeny.

More recently Prof. Nathorst (1912, p. 36) has given a short summary of the results of his cuticular work, and has arranged the genera which have been studied in five groups or orders characterised by the following forms:—(a) *Pterophyllum*, (b) *Otozamites*, (c) *Pseudocycas*, (d) *Nilssonia*, (e) *Ptilozamites*. He does not give a detailed description of the characters of the cuticles of these orders, and they appear to be somewhat unequal in value. He thinks that classes (a), (b), and possibly (c), belong to the Bennettitales, while (d) and (e) are more related to the modern Gymnosperms.

For the purposes of comparison with another group of Jurassic Gymnospermous plants, the excellent figures of stomata of some genera of Ginkgoales given by Prof. Seward * are very useful. The Ginkgoales possess a very definite and constant type of stomatal structure which is somewhat similar to that found in some Nilssonias.

(b) Detailed Descriptions of the Cuticular Structure of some Fossil Cycadean Fronds.

I. Bennettitales.

PTILOPHYLLUM (WILLIAMSONIA) PECTEN (Phill.), n. comb.

The large and somewhat variable group of fronds usually placed in the aggregate species *Williamsonia pecten*, can now be divided into several sections according to the structure of their epidermal cells. The smaller type of frond with regular straight linear or linear-lanceolate pinnæ, closely approximated and attached by almost the whole length of their bases, can be distinguished from the larger type with broader pinnæ, which are slightly falcate, more truncated at the apex and contracted at the base, and with a somewhat more spreading venation. So far as can be seen from external appearance, these two types seem to be connected by a series of gradations, as may be noticed in the figures given in Prof. Seward's 'Fossil Flora of the Yorkshire Coast,' where pl. 3. fig. 1 resembles the small type, and pl. 3. fig. 6 probably represents a frond of the larger type (the differences mentioned above are, however, much more noticeable in other examples from Marske and Whitby collected by one of us). When cuticular preparations are made, the two types of frond can be readily distinguished by the presence or absence of large numbers of thickened circular markings arranged in rows on the cuticles of the second form. These, as mentioned later, were probably of the nature of the thickened bases or scars of hairs of which no trace is seen externally without considerable magnification. The fronds possessing these curious markings, as well as other slight differences, must probably be placed in a new species. The separation of these two types probably involves the distinction of a third and larger type of frond, and the recognition, that the three forms from Marske, figured recently by one of us (Thomas, 1913), are probably three

* Seward (1911¹), pl. 4. figs. 56, 57; pl. 5. figs. 58–62 & 64.

distinct species. We cannot here enter into this question in detail, but it may be remarked that the existence of several species of fronds is rendered probable by the discovery of a number of different species of male and female flowers which probably belonged to some of the plants bearing these *Ptilophyllum* fronds. Three or four forms of male flowers and five or six species of female flowers of the *Williamsonia* type are now known. Besides these we know of two or three other types of flower which probably belonged to plants bearing fronds of some of the other forms placed in the Bennettitalean group in the present paper. In view of the uncertainty of connecting "flowers" with fronds, it appears to be better, at present, to use the generic term *Williamsonia* to apply only to reproductive structures, while referring to the fronds under the names of their form genera. Thus we shall refer to *Williamsonia pecten* (Phill.) fronds as *Ptilophyllum pecten* and to *Williamsonia gigas* fronds as *Zamia gigas*, Lindl. & Hutt.

We may now proceed to the description of the cuticles of fronds of *Ptilophyllum pecten* (Phill.) *. The cuticles of the upper and lower epidermis are very distinct (see Pl. 20, fig. 1). The upper epidermis is composed of uniform cells showing no traces of veins and having no stomata. The walls of the cells show an extremely sinuous outline, a characteristic in the Bennettitalean group; each wall is bent into a regular series of folds which sometimes almost touch each other. The appearance of this cuticle is seen in the photograph reproduced on Pl. 20, fig. 2.

The lower epidermis is composed of cells with a somewhat similar outline, usually rather thinner than those of the upper epidermis and with larger lumina. There is a marked differentiation between the areas above the veins which possess no stomata and the intermediate portions where the stomata occur in two or three rows (Pl. 20, fig. 1). The cells above the veins exhibit a somewhat more elongated form than the rest. A few circular hair-scars are sometimes seen. Near the margin of the leaf the cells become more elongated and their walls thicker.

The Stomata.—In each square millimetre of the lower surface between 50 and 60 stomata occur; this number seems fairly constant in the specimens examined. The individual stomata are roughly circular, with a diameter of about 0.039 mm. The guard-cells are very regularly orientated, so that the slit lies transversely, at right angles to the veins of the pinna. The regular arrangement of the stomata is a characteristic feature of the *Ptilophyllum-Otozamites* group of fronds; in other genera they are irregularly scattered. The guard-cells do not seem to be appreciably sunk. On either side they have a small subsidiary cell (*s*, text-fig. 27), on to which the sinuous walls of the epidermal cells abut. The actual structure of the guard-cells is somewhat difficult to make out from the surface view of the remains of their cuticularised portions (Pl. 20, fig. 3). From the analogy of the modern stomata (*cf.* text-fig. 26) it seems probable that the structures seen in our preparations do not represent the outlines of the cells round the stomatal pore but only parts of their walls which were differently thickened at

* For methods of obtaining cuticular preparations see Thomas (1912), also Nathorst (1912) and Bather (1908). In this section the term cell-wall is used for the ridges on the cuticular film which now represent the walls of the original cells.

two levels. The appearance observed when preparations of stomata are seen in surface view (text-fig. 27; Pl. 19. fig. 1) is as follows:—On either side of the central slit-like pore are two elliptical or hemispherical structures; they are somewhat flattened where they abut the pore, and have rounded ends, appearing, in fact, rather like an ordinary guard-cell appears in surface view. Between these and the subsidiary cells lie two other thickened patches, more or less hemispherical in shape, and apparently somewhat overlying the central structures. They possess a very characteristic outline, and have, like the central ones, a deep brown colour; they often show peg-like extensions towards the centre (see text-fig. 27). Very similar patches are seen in almost all the stomata of this group. They often have a slightly shorter diameter (measured in the polar plane) than the subsidiary cells, but are longer than the central structures; in some cases they appear to be attached to the latter, but in other preparations appear to be free from them at the ends, only lying over them in the centre (*cf.* Pl. 19. fig. 1).

Fig. 27.

Stoma of *Pitlophyllum pecten*, showing subsidiary cells (*s*) and thickenings of guard-cells.

The interpretation of these structures would be somewhat difficult but for our knowledge of the thickenings possessed by the guard-cells of modern forms. In the surface view of these we never see the outlines of the guard-cells, but only the form of their thickenings or woody lamellæ. In *Bowenia* the thickening on the upper face of each guard-cell forms a hemispherical structure at each side of the pore, while the lamellæ on the lower faces contribute the projecting pieces seen at the poles (*cf.* text-fig. 26). In the present case we seem to be dealing with a very similar structure. It is clear that the central patches bordering on the slit do not represent the actual guard-cells. They are often extended into sharp points at the poles, are much smaller than the subsidiary cells, and do not abut directly on to them; moreover, on this view the presence of the intermediate patches cannot be easily explained. It seems likely that we see in the fossil fronds two pairs of thickening patches or lamellæ belonging to the upper and lower sides of the original highly inclined guard-cells. These lamellæ were somewhat similar to the woody lamellæ of the modern forms, but were more strongly cutinised. The central lamellæ would be originally on the lower side of the cells and would be much shorter than in the modern structures. The outer and more thickened patches would represent the upper lamellæ of the guard-cells, which, however, overlapped the lower lamellæ to a much smaller extent than in *Stangeria* and *Bowenia*, but are almost identical in shape

and in their relations to the subsidiary cells, with the thickenings of *Stangeria* seen after maceration (see text-fig. 32). The figures below (text-fig. 28) show an attempt to reconstruct the sectional views of the stomata at various points in accordance with the explanation given above.

Fig. 28.

Diagrams representing surface view of stoma of *Stangeria* (I.), with hypothetical sections (II.) at the different points indicated in I. The shaded and dotted portions represent the probable positions of the thickened portions.

In some modern stomata we find that the upper lamellæ do not extend unchanged to the pore, but the latter is surrounded by a thinner oval area. A precisely similar arrangement is seen in some of the *Ptilophyllum* and *Zamia* cuticles (see Pl. 19. fig. 2).

Unfortunately, it has proved impossible to obtain sections through the cuticles showing the structure of these stomatal remains, but a comparison of the material before us with the stomata figured by Lignier provides some interesting features and gives considerable support to the interpretation just advanced. In his paper on *Bennettites Moriæi*, Lignier* observed that in some sections through the involucral bracts, the stomatal guard-cells, cut through transversely, were seen. In one of his figures, Pl. 1. fig. 9, which is reproduced below (text-fig. 29), two stomata are seen in

Fig. 29.

Section through the epidermis of a bract of *Bennettites Moriæi* showing two stomata cut through. The characteristic form of the guard-cells and subsidiary cells is well shown. (After Lignier.)

section cut through transversely somewhat near their poles. The guard-cells are seen to have the pear-shaped outline, so characteristic of the modern forms (*cf.* text-figs. 26, etc.), with a marked upward extension. The upper walls of the guard-cells are drawn as being somewhat thickened, the wall abutting on the slit being thin;

* Lignier (1894), p. 18, pl. 1. figs. 8, 9.

on either side of the guard-cells are two well-marked subsidiary cells. The figure further shows that the guard-cells of adjacent stomata were approximately parallel to each other, and that the inclined guard-cells became more vertical at the poles and their upper or outer thickening became less. In figure 8 of the same plate another stoma is shown cut through more or less longitudinally, but the section has not here passed through the centre. No stomata were seen in surface view, but in a later paper* surface views are given of stomata from the interseminal scales of the same specimen, which had been obtained by collodion casts. Though the details are not very clearly shown, it is evident that these stomata were of the same type as those which are described here, the guard-cells and subsidiary cells of some of them being quite characteristic. The structure of these stomata in Lignier's species of *Bennettites*, a genus closely allied to our Yorkshire *Williamsonia*, not only provides an interesting and very close comparison with the modern *Bowenia* type, but also goes some way towards demonstrating that the interpretation which we have given above, of the meaning of the surface views of stomata in this group, is correct.

We have dealt at some length with the structure of these *Ptilophyllum* stomata because the form of these organs is remarkably constant throughout the group of fronds which we have assigned to the Bennettitales. Moreover, their appearance is so complex that at first sight it is very difficult to understand them; no previous observers seem to have made any attempt to grapple with the problem, but have merely spoken of the thickening patches as the guard-cells.

PTILOPHYLLUM sp.†

Typical examples of *Ptilophyllum pecten*, from the Lower Estuarine beds of Whitby and Marske, show perfectly smooth cuticles with very few hair-scars. In the larger fronds, however, from the same beds, a large number of very characteristic structures are seen, which are also present on the fronds of the small *pecten*-form which occur plentifully in the Cloughton Wyke beds of Middle Estuarine age.

These structures are shown in Pl. 20, fig. 4. They may be described as small annulate projections from the surface of the cuticle .03 to .04 mm. in diameter, occurring in numerous regular rows running parallel to the length of the pinna. The rows are very close together, being often separated by a space less than their individual width. The annulate structures of the same row are frequently contiguous and are sometimes unevenly thickened, being broader and darker laterally and thinner at the points next to the adjacent rings, so that an appearance is produced of pairs of crescentic patches. They are so numerous as to obscure entirely the outlines of the epidermal cells and make the stomata very indistinct.

We may regard these structures as the bases of hairs, with which the lower surface of the frond was covered. Scattered hair-scars occur on the ordinary types of frond, which

* Lignier (1912), p. 426, fig. 1.

† This species will later be described as *Ptilophyllum hirsutum*; a figure of typical outline of a frond is given in the 'Fossil Flora of the Marske Quarry,' text-fig. 3 B. Thomas (1913).

do not differ markedly from those on some modern Cycad fronds, but they are not so massive as in the present example, nor are they arranged in such a regular sequence. When, however, the epidermal cells are arranged in regular rows, the outgrowths from them may be expected to possess the regular arrangement which we see here. The fronds just mentioned appear to possess stomata which are somewhat similar to those above described (see text-fig. 30).

An interesting comparison can be drawn between the hair-scars seen so abundantly on these fronds, and the hairs on the bracts and lower part of the strobilar-axis in the flower of *Williamsonia scotica*, recently described from the Kimmeridge of Sutherland by Prof. Seward *. The hairs in the latter formed a very striking feature; they were very numerous, and arose from the thickly cuticularised outer wall of an epidermal cell which formed a dark ring round the lower end of the hair. Higher up the walls of the hair were much thinner. If leaves bearing hairs of this type had been preserved as impressions in the same way as the material now before us, we could scarcely expect the substance of the hair to be preserved, though the cuticularised base would remain as

Fig. 30.

Stomata of *Ptilophyllum* sp., showing form of the guard-cell thickenings.

a clearly visible ring on the outside of the epidermis. It is possible, too, that hairs of this type might have been deciduous. The occurrence of these characteristic hair-scars forms another link connecting the *Ptilophyllum* fronds with the type of inflorescence now known under the generic name of *Williamsonia*.

ZAMITES (*WILLIAMSONIA*) GIGAS, Lindl. & Hutt.

There have yet been found few examples of this species from which good cuticular preparations can be made. One specimen, however, from Marske, which has large pinnæ and apparently belongs to this species, yields preparations which show the cells of the upper epidermis fairly clearly. The lower epidermis is somewhat crumpled, and the preparations made from it are always rather dirty; its cells are very indistinct, and the pits in which the stomata appear to have been sunk are usually filled with matrix. We can, however, make out some of its epidermal characters. The upper epidermis is thinly cuticularised, and is composed of cells of the usual type with very sinuous outlines. The lower cuticle is also very thin, and the outlines of the cells cannot be seen. The stomata are arranged in fairly regular rows with their axes at right angles to the rows; the guard-cells appear to be considerably sunken and are surrounded by a brown, rather highly cuticularised area, probably the walls of the pit. The guard-cells possess

* Seward (1912), p. 108, pl. 12. figs. 40-42 & 45.

the same characteristic thickenings as previously described; the crescentic patches are terminated in sharp points (*cf.* text-fig. 31) and partially cover the ventral patches, while the central space on either side of the slit is thinner. The small projection from the centre of the crescentic patch which is characteristically seen in *Ptilophyllum pecten*, *Dictyozamites*, etc., is here apparently absent.

A large frond from Marske with a form somewhat like that of *Zamites gigas* yielded, after some difficulty, a cuticular preparation of the upper epidermis, in which the cells are rounded in shape and quite distinct from the other examples in the present group.

ZAMITES (?) sp. Scale-leaves from stem.

In the Natural History Museum (Palæobotanical Department) of Stockholm, there is an interesting specimen, collected at Whitby by Prof. Nathorst in 1909. It appears to be the remains of a branching stem about 5 cm. wide; one part of it is fairly well preserved, and seems as if covered with numerous overlapping scales or leaves; it is

Fig. 31.

Stoma of *Zamites gigas*, Lindl. & Hutt., showing form of the thickening on the upper side of the guard-cells.
The circle surrounding the stoma represents the outline of the shallow pit in which the stoma was sunk.

somewhat similar to those of the James Yates Collection, figured by Wieland (1911, figs. 2, 3).

The scale-leaves appear as if lanceolate in shape, of rather thick texture, and striated; they are very indistinctly outlined. They yield, however, cuticular preparations which show a certain amount of structure, though considerably obscured by a mass of hypodermal or other tissue, which was evidently thick-walled and has resisted both decay and maceration. The actual epidermal cells are not clearly seen, but some of the stomata are quite visible; they appear to be similar in both form and arrangement to those previously described (see Pl. 19. fig. 2).

OTOZAMITES GRAPHICUS (Leckenby, ex Bean MS.) n. comb.

Fronds of this form from Marske yield good preparations, but the cuticles are very thin and fragile, and hence only small pieces are obtainable. The similarity of its cuticular structure to that of the foregoing types is very noticeable. The upper epidermis has small squarish cells with rather thick walls showing closely approximated folds. The lower cuticle is very thin, with the usual type of cells; the stomata are rather sparsely scattered, only 40-50 per sq. mm. occurring in the specimens examined. In this they contrast with *Otozamites Feistmanteli* from the same bed, in which the stomata are twice as numerous. The stomata are arranged more or less in lines between the veins which are marked by the greater elongation of the epidermal cells; they have a fairly regular orientation, the pore being at right angles to the veins. They are not deeply sunken.

The stomatal structure (see Pl. 19. fig. 4) is like that of *Zamites*. The thickening patches are hemispherical to spindle-shaped, and the central area is thin. The sides of the pore seem to be slightly thickened, and the two lateral subsidiary cells are also somewhat thicker than the other epidermal cells.

OTOZAMITES FEISTMANTELI, Zigno.

The dark shale bed at Marske has yielded good specimens of the smaller *Otozamites*, which gives good cuticular preparations. These show some interesting points; the stomata are clearly seen and are very numerous, being almost contiguous in lines between the nerves. About 100 occur in each square millimetre. In structure and form they are similar to the stomata of the larger species, but the lateral thickening patches thin out gradually towards the slit, showing no sharp margin, but more or less merging in towards the central area. The latter is very slightly cuticularised except at the edges of the pore, which stands out clearly as a dark central line (*cf.* Pl. 19. fig. 3). The subsidiary cells are again slightly thickened, and their relations to the other epidermal cells are often well seen.

DICTYOZAMITES HAWELLI, Seward.

The cuticular structure of *Dictyozamites Johnstrupi*, Nath. has already been described in some detail by Prof. Nathorst (1907, p. 12, Taf. 3. figs. 2-8). The structure of the present examples from Marske does not differ materially from that of the Bornholm examples, but the structure of the stomata can now be further elucidated. As in the former cases, the upper epidermis is devoid of stomata, and is composed of cells with very sinuous walls, the reticular course of the veins being marked by the elongation, narrowing, and thickening of the cells above them. The folds in the cell-walls of this species are not so deep as those previously described. The lower epidermis is less strongly cuticularised than the upper; the course of the veins is again marked by elongated cells with less folded walls. The stomata occur only between the veins; they are more or less irregularly scattered, but still show a distinct tendency to have the pore

placed at right angles to the adjacent veins. The stomata are numerous, about 120 per sq. mm. Small papillæ occur on many of the epidermal cells, but are not so conspicuous as in *D. Johnstrupi*. The structure of the stomata appears somewhat complicated owing to the presence of what seem like small thickened outgrowths or papillæ from the surface above the guard-cells, and which are shown in Pl. 19. fig. 8, where the guard-cells have been by some means pulled apart. In other cases they appear to overlap the central space, as shown in Pl. 19. fig. 6, and Pl. 20. fig. 7, or give the impression of the guard-cells interlocking with each other, as mentioned by Prof. Nathorst. These projections are probably of a nature very similar to those observed in the case of *Ptilophyllum pecten* (cf. Pl. 19. fig. 1). Sometimes they are not visible (cf. Pl. 19. fig. 7), being probably on the under side of the preparation; in this case the thickening patches are very comparable to those in *Otozamites*. On either side of each stoma are subsidiary cells, somewhat thickened and often much elongated (cf. Pl. 19. figs. 6-8). In preparations mounted with the lower side of the stomata uppermost, the latter seem to have the same form as in several other examples of this group, the ventral patches on the guard-cells being somewhat thinner.

ANOMOZAMITES (WIELANDIELLA) NILSSONI (Phill.), n. comb.

Leaves of an *Anomozamites* form have been found in Yorkshire near Gristhorpe, Marske, and Roseberry Topping. They are very similar in shape to those of *Wielandiella angustifolia*, Nath., and may well have belonged to plants bearing floral structures similar to those found by Nathorst in connection with his species; such structures are not, however, yet known with certainty from England, though stems have been found which are almost identical with those of the Rhætic plant. The cuticular preparations obtainable from the Yorkshire *Anomozamites* leaves are exceedingly beautiful, and show clearly the epidermal and stomatal characters. The leaves from the Lower Estuarine Series at Marske differ slightly from those of Gristhorpe in the possession of numerous spherical papillæ on their lower cuticles; the stomata are also slightly smaller.

In the specimens studied the upper epidermis is very uniform, and is composed of more or less rectangular cells; the courses of the veins are not clearly marked off, but occasionally they are indicated by lines of slightly more elongated cells. The cell-walls are again sinuous or zig-zag, but the folds have a different form from those in the other genera, each of the ridges and grooves being angular rather than nail-shaped or rounded; each point appears to possess a small pad of cutin. The upper epidermis is devoid of stomata. In some preparations the cuticle from the rachis of the frond is also seen; this has regular rectangular cells with straight walls, and occasional structures like rudimentary stomata. The lower epidermis has a structure similar to that of the upper, but the cells are less regular, and the stomata are irregularly scattered (cf. Pl. 20. fig. 8). The courses of the veins are seen, though not very clearly, and no stomata lie upon them. There are about 70 stomata per sq. mm., and their form is very clearly seen in most examples. It is shown in Pl. 19. fig. 5, and Pl. 20. fig. 8. The central slit is

bordered by two fine linear membranes, beyond which are the thickening patches somewhat hemispherical in shape, with a straight inner surface and rounded or pointed ends. The inner surface of each appears to be carried on as a wall, a fact which is of some importance in the interpretation of the structure. There are two crescentic subsidiary cells.

In their structure these stomata resemble the examples seen in *Otozamites*, especially *O. graphicus*, and also in *Teniopteris vittata* (cf. Pl. 19. fig. 12).

TENIOPTERIS VITTATA, Brongn.

The entire linear-lanceolate fronds with a strong petiole, central midrib, and dichotomising nervation which are so common in the Yorkshire plant-beds were designated *Teniopteris vittata* by Brongniart and, as the name implies, regarded as fronds or pinnæ of a fern. Phillips in 1829 (p. 147) regarded it as allied to the recent *Scolopendrium*, but subsequent authors* have for the most part considered it to be allied to *Marattiopsis* and have placed it among the Marattiales. Another view was that it had affinities with the modern genus *Oleandra*, since it possessed a petiole with a clean-cut base, which suggested that it had become detached from its original stem or rhizome by the formation of an absciss-layer†; thus the generic name of *Oleandridium* was used by Schimper (*loc. cit.*), Feistmantel, and others.

On the other hand, the view was put forward by some authors, especially Bornemann (1856, p. 59), that it was really the pinna of a pinnate Cycadean frond such as *Stangeria*, and the name *Stangerites* was instituted to replace *Teniopteris*. Several subsequent authors have kept in mind this possibility, but no definite evidence has been forthcoming in support of either view. *T. vittata* has never been found bearing synangia or sporangia; it has also never been found attached as a pinna to a larger frond. Recently, however, it has been found in association with Cycadean stems of the *Wielandiella* type, and one specimen appears to show organic connection between a frond and one of these stems. It is almost certain that we have to deal with entire fronds and not with pinnæ, and thus a very close comparison is offered with the fronds of *Anomozamites* (*Wielandiella*); the latter possess the forking nervation of *Teniopteris* and may be regarded as segmented fronds of the *Teniopteris* type. *T. vittata* itself is sometimes found with the lamina divided into irregular segments. When we come to examine the cuticular structure we obtain very strong confirmation of the Cycadean view.

In its cuticular structure *Teniopteris vittata* approaches very closely to *Anomozamites Nilssoni*. The upper epidermis is fairly strongly thickened and was composed of squarish to hexagonal cells, with somewhat sinuous walls (Pl. 20. fig. 5). The vein-courses are only slightly differentiated, the cells being more regularly arranged in rows. The outline of the cell-walls is interesting owing to the presence of a solid piece of thickening at the point of every fold (this is just visible in the photograph); in some cases it gives the appearance of an almost straight wall ornamented with projecting

* Schimper (1869), p. 607.

† Seward (1900), p. 158.

tooth-like masses of thickening, but it can always be referred to the same general type of wall-folding. We see the same tendency in *Anomozamites*. The lower epidermis is composed of cells similar to those of the upper epidermis, but of a somewhat larger size; their walls are thinner and may be sinuous or have the folds filled in, as just mentioned. A few rounded hair-scars are present. The stomata are absent from the nerve-courses and are irregularly scattered between (*cf.* Pl. 20. fig. 6); about 60–70 occur in each square millimetre. They have the same general form as previously described, but are apparently rather more complex. Some of the examples from Marske (Pl. 19. fig. 12) do not differ greatly from the *Anomozamites* stomata except in the proportions of thickening, the areas abutting immediately on the slit being the more strongly thickened. The stomata in the Gristhorpe examples appear to have been slightly sunken; in most cases the pore is clearly seen and is surrounded by a strongly thickened margin running to a point at either pole (Pl. 19. fig. 11). The outer (upper) thickening patches are often not so clearly defined as in the other genera, but have the same relations to the auxiliary cells as those of *Anomozamites*. These outer patches are often thinner than the inner ones and occasionally appear to project into the centre, almost above the slit (Pl. 19. fig. 10); they sometimes develop a small projecting hump, similar to that seen in *Ptilophyllum pecten* and *Dictyozamites*. Though it is very difficult now to understand what the structures, which we see in the surface view, must have meant in the solid, yet it is quite evident by comparison with the cuticles of other groups, that *Teniopteris vittata* possessed essentially the same epidermal and stomatal characters as the other fronds in the Bennettitalean group, being especially like *Anomozamites*.

When this is considered in addition to the facts quoted above, it seems that we have strong reasons for removing this species and its near allies from among the ferns and for including it in the Bennettitales.

TENIOPTERIS MAJOR, Lindl. & Hutt.

Some *Teniopteris* fronds occur at Gristhorpe which are somewhat larger than those of *T. vittata*, but are readily distinguishable by their distinct and widely separated dichotomising veins, which are one millimetre apart or more and are very characteristic; these fronds are probably distinct from *T. vittata*, and may be referable to the form described by Lindley and Hutton as *T. major*.

In its cuticular characters it does not differ much from the species just described; the cells of the upper epidermis are rather large and have well-developed angularly folded walls, slight additional thickenings being present on the tips. The lower epidermis has somewhat scattered stomata which resemble very closely those of *T. vittata* (Pl. 19. fig. 9).

In this case the epidermal characters do not, so far as we have seen, present any sharply marked features on which we can rely for the separation of the related forms *major* and *vittata*.

II. Nilssoniales.*

NILSSONIA, Brongn.

While the cuticles of the genera already described show a very considerable agreement, both in the structure of their epidermal cells, and also in the form and arrangement of the stomata, the cuticles of the remaining genera *Nilssonia*, *Ctenis*, and *Ptilozamites* are quite distinct from them. In these forms the epidermal cells are somewhat irregular in shape, being rounded, hexagonal, or rectangular; the walls are straight without any indication of folding. The stomata appear to be somewhat sunken, and when any trace of the guard-cells can be seen, they apparently lack special thickening and have a more or less rounded contour. The appearance of these cuticles therefore marks them at once as distinct from those of the Bennettitalean group.

The Nilssonias must have been important constituents of the Jurassic vegetation of the Yorkshire area. Fronds of *Nilssonia compta* occur abundantly in the Middle Estuarine bed in the neighbourhood of Gristhorpe Bay. In the same locality *N. mediana* occurs; this is almost identical in form with a plant which is very abundant in the Lower Estuarine beds of Marske and Roseberry Topping, but which possesses very much more delicate cuticles. The same form occurs at Whitby, while another type, closely allied to *N. compta*, but probably a species new to England, is found at Roseberry Topping. The cuticles of these three species or forms have been examined, and their characters are briefly described below. The cuticles of the Swedish species *brevis*, *polymorpha*, and *pterophylloides* have been described and figured by Prof. Nathorst. They apparently agree somewhat closely with the English species described here, though little or nothing could be made out about the form of their stomata.

NILSSONIA COMPTA, Phill.

The examples of this species from the Gristhorpe bed indicate that they originally had a somewhat delicate texture, and owing to this and to the thinness of the cuticles it is not easy to obtain good cuticular preparations from them. Even after portions of the carbonised lamina have been successfully removed from the matrix they frequently break up into numerous small fragments during the clearing process. It is believed that this affords a character of real value in describing and identifying fronds. In many descriptions of leaf-impressions, the lamina is described as leathery or coriaceous, from the appearance of the specimen when still attached to the rock or even from the appearance of its cast; in most of these cases little value can be assigned to such a statement. When, however, we are able to remove the carbonised or mummified plant from the matrix and investigate the structure of its cuticle, we have definite grounds for a statement as to the comparative texture of the lamina.

In the case of the plants preserved in the fine-grained mudstone of the Gristhorpe bed, we can say with some confidence that such fronds as *Tæniopteris vittata*, *T. major*, *Ctenis falcata*, and especially *Ptilozamites Leckenbyi*, had a firm and coriaceous lamina. Others, such as *Anomozamites Nilssoni* and *Nilssonia compta*, were of intermediate

* See page 196.

character, while *Nilssonia mediana* and *N. orientalis* had a distinctly delicate or thin lamina, approaching the type possessed by the ferns and seldom separating from the matrix in an entire condition.

The upper epidermis of *N. compta* is very distinct from the lower. It possesses small rather irregularly-shaped cells arranged, to some extent, in rows. The individual cells vary greatly in shape; they may be isodiametric with rounded or angular outlines, or they may be more elongated, and of rectangular to fusiform shapes. The latter kind probably occur above the nerves. All the cell-walls are straight and not sinuous. There are usually no stomata present, but in one or two cases, structures which may be stomata, are seen near the margin of the lamina. A few rounded papillæ occur, these being of the type common to most of the Nilssonias. Their form is better seen on the rachis and in other species.

The cells of the lower epidermis are much more clearly seen; the nerve-courses are distinctly marked by the occurrence of rows of elongated cells, often bearing papillæ, but devoid of stomata. The spaces between the nerves are occupied by rounded cells, with a good deal of thickening in the corners. The cells surrounding the stomatal apertures are much thickened. The stomata on these leaves are not numerous; estimates from the very fragmentary cuticles indicate between 40 and 50 in each sq. mm. It is almost impossible to make out the exact structure of the stomatal openings, but the following features are recognisable. Each aperture is surrounded by crushed thickened cells, which evidently formed, in their original condition, a chimney-like structure, being raised beyond the surface of the leaf, and enclosing a pit, at the base of which the guard-cells were placed. Amongst recent Cycadean cuticles, a similar condition is seen in *Dioon* and in *Cycas revoluta*. The fossil cuticles do not show the projection of the cells surrounding the stomatal apertures, owing to crushing in preservation. The cells are seven or eight in number, and often show a star-like arrangement. Their relations to the ordinary epidermal cells are not clear. When some of the preparations are mounted so that their inner side (viz., that abutting on the mesophyll) is uppermost, the remains of the guard-cells can often be distinguished (Pl. 19. figs. 14 & 18). The guard-cells themselves may be somewhat indefinite, but the pore is usually quite distinct. The thickening of the guard-cells seems to be of a simple type, and not complex as in the Bennettitalean fronds; this may perhaps be correlated with their sunken position.

The cuticle from the rachis below the frond presents some distinctive features. It is composed of elongated spindle-shaped cells, giving the appearance which sclerenchymatous cells show in longitudinal section. At frequent intervals occur oval cells, which are considerably more cuticularised than the rest. In the centre of each of these is a rounded area, probably representing a short papilla. Structures of this nature are also found sparingly on both surfaces of the lamina, as mentioned above.

NILSSONIA MEDIANA (Leckenby, ex Bean MS.), n. comb.

Examples of this species from Gristhorpe do not yield good preparations owing to the delicate nature of the cuticle; specimens from Marske, which are very similar in external form, have a well-thickened cuticle, and preparations from these show the outlines of

the cells very clearly. The cuticles of *N. mediana* differ from those of *N. compta* only in detail. The cells of the upper epidermis are more elongated and spindle-shaped, and those of the lower epidermis are very sharply differentiated according to their position above or between the nerve-courses (Pl. 20. fig. 10). Some papillæ are present, and each of the rounded cells from the areas between the nerves is thickened externally and forms in consequence a slight convex projection on the surface of the cuticle. The stomata can be more clearly seen than in *N. compta*. The cells surrounding the stomatal aperture are only slightly raised and thickened, and therefore they are less crushed. The raised portions are in this species seen to be merely small outgrowths from the ordinary epidermal cells, slightly thickened (*cf.* Pl. 19. fig. 15).

NILSSONIA sp.

A new form of *Nilssonia* has recently been discovered at Roseberry Topping, Yorkshire. It is closely allied to *N. compta* of the Middle Estuarine beds, but has more rounded and less regular leaf-segments, approaching the *N. polymorpha* type. It also has a very marked wrinkling of the lamina. It yields good cuticular preparations (Pl. 20. fig. 9), though these are somewhat brittle. The stomatal apertures in this species are surrounded by several strongly thickened conical cells which originally must have formed conspicuous projections, but which are now usually much crushed (see Pl. 19. fig. 13). The papillate hairs are also well marked in this new form.

NILSSONIA ORIENTALIS, Heer.

Of the three species of *Nilssonia* described in Prof. Seward's 'Jurassic Flora of Yorkshire,' *N. compta* and *N. mediana* were well defined and represented by many specimens in the then existing collections, while the third was only known from a few examples. The name given to this was *N. tenuinervis*, a title which had originally been proposed by Nathorst (1880, p. 35) for linear fronds, little divided and having numerous fine veins. It now seems impossible to separate this English form from the Siberian type described by Heer (1878, Pl. 4. figs. 4-9) as *N. orientalis* *. In his original MS. notes on the specimens amongst which *N. tenuinervis* was found, Prof. Nathorst † remarked on the great similarity in external form between the new species and some examples of *Tæniopteris vittata*; and, in fact, they can only be distinguished readily in well-preserved examples. In 1909, when working at the revision of the Swedish species of *Nilssonia* and studying their epidermal characters, Prof. Nathorst (1909, p. 27) made cuticular preparations of some fronds from Cloughton Wyke, Yorkshire, which were in his collection. These preparations showed a type of structure unlike that of the other *Nilssonias*, and identical with that described above for *Tæniopteris vittata*. Nathorst concluded that the Yorkshire frond was not a true *Nilssonia*, but was rather allied to the ferns; he accordingly instituted the new genus *Nilssoniopteris*, giving as

* With reference to the nomenclature of this form see Thomas (1913), p. 240.

† Our best thanks are due to Prof. Nathorst for the kind way in which he placed all his material at my disposal, for showing me many of his manuscript notes and adding materially to the elucidation of this difficult species.—H. H. T.

its characters the distinct cuticular structure and the dichotomising of some of the secondary veins. In 1911, while working in Prof. Nathorst's laboratory, one of us was able to re-examine his specimens and preparations, and found that one of Prof. Nathorst's fronds, that figured on Pl. 6. figs. 23 & 25, was really a specimen of *Tæniopteris vittata*, showing a forking venation. As this form yields good preparations it had been used in the description of the examples figured on his Pl. 7. fig. 21.

From the other specimen only very fragmentary preparations could be obtained, but they were sufficient to show that their structure was quite distinct, agreeing with that of true *Nilssonias*, in which the cuticles are extremely delicate. In this form the fragments of cuticle (one of them is shown in Pl. 20. fig. 11) show that the upper epidermis is composed of elongated rectangular or tapering cells, with straight walls similar to those of *N. mediana*. The lower epidermis has fine rectangular or squarish cells with straight walls and small dark rounded structures like the papillate hairs of *N. compta* and *N. mediana*. No stomata can be clearly seen.

Other examples of the same kind have been obtained from the Gristhorpe bed; they agree in the characters given above, but do not add any information. They show a well-marked minute wrinkling of the cuticular film which rather obscures detail.

From these facts it appears that the name *Nilssoniopteris* must be abandoned, for the cuticles of the Yorkshire specimens referable to *Nilssonia orientalis (tenuinervis)* are of the same type as those of other species of *Nilssonia*.

CTENIS FALCATA, Lindl. & Hutt.

Specimens of this species found in the Gristhorpe bed flake off very readily from the matrix and yield fairly good cuticular preparations.

The upper epidermis is very strongly cuticularised, and the thickening evidently originally extended down the radial walls to a considerable extent, for the cell outlines are now represented by thick badly defined lines of cuticle. The cells above the nerves are especially strongly cuticularised, often causing the nerve-courses to stand out as dark, slightly raised lines; only two or three rows of cells, however, are affected, and not a broad series. The anastomosing of the veins is clearly shown on both the upper and the lower cuticles. The cells of the upper cuticle are irregular in outline, but are more or less isodiametric, varying between squares and hexagons. Some of them seem slightly papillate, but are not markedly so. No stomata are present on the upper surface. The lower cuticle (Pl. 20. fig. 13) is somewhat similar to the upper in thickness, but the cell outlines are larger and more distinct. The vein-courses are marked by a more regular arrangement of cells, and the stomata occur somewhat scattered in the intervening areas. Their numerical distribution is from 60–70 per square mm.

As in *Nilssonia*, the guard-cells are sunk and are not clearly seen. The cells surrounding the stomata-openings are not much differentiated in form or thickening from the ordinary epidermal cells; where they adjoin the guard-cells, however, there is a solid ring of thickening round the stomatal cavity (Pl. 19. fig. 19). The latter is apparently not very deep, and the guard-cells are often seen at its base. They are

slightly thickened, and in many cases are not preserved, but the stomatal slit is frequently visible.

Ctenis falcata presents a type of stomatal structure comparable with that seen in *Nilssonia mediana*. In the former species, however, there is little or no tendency towards the papillate outgrowths from the subsidiary cells surrounding the stomatal apertures, which is so pronounced a feature in some Nilssonias, for example *N. compta*.

PTILOZAMITES (CTENOZAMITES) LECKENBYI (Leckenby, ex Bean MS.), n. comb.

The cuticles of this form have already been figured by Prof. Nathorst (1908, Taf. 1. figs. 3-5) as providing an example of a frond with stomata on one side of the pinnæ only. He did not enter into a detailed description of the structures seen.

In comparing the structure of the cuticle with that of other genera, the following points may be noted. The leaves have a very stout texture with thick cuticles so that the leaflets separate off whole from the matrix and can be easily handled. In most of the specimens examined the serration of the margin is not pronounced, but a few teeth are seen near the apex. The upper cuticle is very thick and shows very clearly the outlines of the epidermal cells which must originally have had their radial walls very considerably cuticularised. The cells are irregularly shaped, but are straight-walled and somewhat rectangular. The cells above the nerves are more elongated than the rest. The lower cuticle is also thick and shows cells similar in form to those of the upper epidermis. The cells are slightly elongated above the veins, but the courses of the latter are not clearly defined and are not always devoid of stomata. The stomata were numerous, their distribution being about 90 per square mm. (Pl. 20. fig. 12). In form they show considerable resemblance to those in *Ctenis*, being slightly sunk, and having the walls and opening of the stomatal cavity much thickened (Pl. 19. fig. 17); the cuticle round the opening, in fact, forms a little ring, slightly raised beyond the surface of the pinna. The seven or eight subsidiary cells are somewhat thickened and appear as a group darker than the rest. On mounting a lower cuticle so that the mesophyll side is uppermost, the outlines of the guard-cells are faintly seen; these appear to have some resemblance to those of the *Williamsonia*-*Tæniopteris* group. The slit lies at right angles to the longer diameter of the stomatal cavity, and on either side of it there are distinct though faint indications of guard-cells (?) with pointed ends (Pl. 19. fig. 16).

(c) The Cuticles of some other Types of Fossil Cycadean Fronds.

The cuticular structure of some other genera have been described by other authors, but in several cases the figures given are not sufficiently detailed or accurate to allow of their close comparison with the examples here described. Prof. Nathorst has figured and described the cuticles of two species of *Pseudocycas* from the Lower Cretaceous (1907, pp. 7, 9, Taf. 2. figs. 1-11). These cuticles show more or less rectangular cells arranged in regular rows and having undulating walls. The stomata are not clearly recognisable, but from Taf. 2. fig. 11, they appear to be somewhat similar in general

character to those of *Anomozamites* and the Bennettitalean group. Nathorst compares the structure of these forms with those of *Otozamites*, *Pterophyllum*, &c., and it does indeed seem probable that these *Pseudocycas* fronds should be included in the Bennettitalean group, though in the disposition of the stomata in a furrow they are distinct from any of the known Jurassic forms.

Berry, in his account of the Lower Cretaceous flora of Maryland (1911, p. 334), gives a diagrammatic figure of the cuticle of *Dioonites* (= *Zamites*) *Buchianus*, Ett. The cells as in *Pseudocycas* are arranged in regular rows. They are rectangular and had somewhat undulating walls. The stomata are all orientated in the same direction (cf. *Ptilophyllum*, &c.), presumably with their long axes parallel with the long axis of the pinna. If the analogy with *Ptilophyllum* held, however, the stomata should be orientated at right angles to the long axis of the pinna. The structure of the guard-cells is shown diagrammatically and cannot therefore be strictly compared with our types; there is considerable probability, however, that they are similar in form to those of *Otozamites* and *Tæniopteris*. Schenck has figured *Dioonites Dunkerianus* with sinuous walls and stomata irregularly scattered. We can probably, therefore, include *Zamites* (*Dioonites*) in the Bennettitalean group.

The cuticular structure of various species of *Thinnfeldia* has been described by Schenk, Seward, Gothan, Zeiller, and other authors. It is very doubtful, however, whether this genus should be included among the Cycadophyta. The cuticular structure seems to be distinct from that described above, and is apparently more closely related to that of the Ginkgoales, especially so far as the stomatal form is concerned. There is an approach to the structure shown by *Nilssonia mediana*, but on the whole *Thinnfeldia* seems to be quite distinct.

(d) Summary of the Structure of the Fossil Cuticles.

The comparison of the structures just described brings out very clearly one point, viz., that the Cycad-like fronds from the Jurassic rocks of Yorkshire fall into two sharply divided groups according to their epidermal structure. The fronds in the first group have epidermal cells with sinuous walls and comparatively thin cuticles; the cells are in general rectangular in shape, but in most cases their shape is not very apparent. The stomata occur only on the lower side of the leaf and are usually absent below the vein-courses; they exhibit a strong tendency towards arrangement with their axes at right angles to the veins. Though there is some variation in the details of stomatal structure, the general plan is as follows. The guard-cells were level with the surface or only slightly sunk; the outlines of the guard-cells themselves are never seen but only those of the thickenings which they possessed. The thickenings of the guard-cells usually appear as lamellæ of definite shape and give the stomata a characteristic appearance; on either side of the pore is a pair of smaller patches, probably representing the lamella from the lower side of the guard-cells, and outside there is another pair of lamellæ, usually somewhat hemispherical in shape, representing the thickenings from the upper side of the guard-cells. At the poles there are usually two clear spaces,

looking sometimes like cells, but probably representing the thin places above the ends of the guard-cells. More or less surrounding the stomata at the sides are two large subsidiary cells, often somewhat thickened. The above description applies both to (a) pinnate compound fronds, e. g. *Ptilophyllum*, *Otozamites*, *Dictyozamites*, and *Zamites*, some of which are known to have belonged to plants bearing flowers of the *Williamsonia* type, and to (b) simple fronds, e. g. *Teniopteris*, which is usually entire, and *Anomozamites*, which is usually pinnately lobed. Fronds of the latter have been found to belong to plants with a peculiar type of Bennettitalean flower (*Wielandiella*). There are then, strong grounds for regarding the type of cuticular structure described above as characteristic of the fronds of the Bennettitales.

The cuticles of the second series, which includes the genera *Nilssonias*, *Ctenis*, and *Ptilozamites*, are sharply marked off from those described above. In this section the cuticles may be thin (as in the *Nilssonias*) or very thick; they appear to have been formed from cells of elongated, rectangular, rounded, or fusiform shapes, with straight walls. The stomata are on the under side of the leaves and exhibit no regular arrangement, the guard-cells are somewhat deeply sunken, and the six or eight thickened subsidiary cells which surround them often form a somewhat overarching canopy. The guard-cells are hemispherical or spindle-shaped, they are only thinly cuticularised, and definite lamellar thickenings are seldom seen.

The cuticles in this group do not agree with one another so closely as do those in the Bennettitalean group, *Ctenis* and *Ptilozamites* are similar, but the *Nilssonias* are not so closely related. The fronds were variously shaped, those of *Ctenis* and of some species of *Nilssonias* were large and pinnate, *Ptilozamites* was bipinnate, while the fronds of *Nilssonias orientalis* were simple and often entire or only slightly lobed. It may be noticed that the thickness of the cuticle in these forms bears an apparently definite relation to the size of the frond. Of the bipinnate *Ptilozamites* no complete fronds have been obtained, but they were probably of considerable size, their cuticles were very thick; the large fronds of *Ctenis*, and also of the Lower Estuarine form of *Nilssonias mediana* had cuticles of considerable thickness. The smaller fronds of *Nilssonias compta* from Gristhorpe had thinner cuticles, while the much smaller entire fronds of *Nilssonias orientalis* had very delicate cuticles. We see no such gradation among the Bennettitalean group. We know very little about the affinities of the plants which bore the fronds placed in this second section. Prof. Nathorst has found fronds of *Nilssonias pterophylloides* associated with numerous seed-like bodies in such a way as to render their original connection probable, but the stems and reproductive organs of the other species and genera are as yet quite unknown. Though it is quite possible that some of these plants were closely related to the recent Cycadales, there is as yet no definite evidence in support of this view. It may be convenient if we institute the name of NILSSONIALES for the group of fronds which has just been described.

IV. COMPARISON OF RECENT AND FOSSIL CUTICLES.

The difficulty of comparing the epidermal structure of the recent and fossil fronds is considerable owing to the impossibility of obtaining good sections through the latter. It is, therefore, necessary in most cases to employ only the information obtained from surface view, and since in most cases the different types of stomata are chiefly to be distinguished by their appearance in transverse section, our work becomes much more difficult. It is possible, however, to notice a number of points of resemblance, and in fact many interesting comparisons can be made.

In the first place we may notice that the fossil fronds, like the modern ones, exhibit some xerophytic tendencies in the thickness of their cuticles, in the frequently sunken stomata, and in the very small number of stomata per square millimetre, the numerical distribution of stomata in both being remarkably similar. These facts may indicate that the xerophytic characters of the modern Cycadean fronds are ancestral.

Comparisons can be drawn between the shape and structure of the guard-cells. The sections through stomata of *Bennettites Morierei* figured by Lignier, indicate that the Bennettitalean stomata had a structure of the general Gymnosperm type, and that the guard-cells showed the characteristic Cycadean form. The appearance in surface view of the guard-cells of several genera, when they are not obscured by overlying cells, is somewhat characteristic. It is well seen in *Stangeria*, *Bowenia*, and sometimes in *Zamia*, *Macrozamia*, and *Encephalartos*. The guard-cells are thickened in a peculiar way with a small oval thin patch round the pore, on either side of which is a spindle-shaped or crescentic thickened mass. There is a thin patch above each end of the guard-cells, but a solid or Y-shaped thickening at the points of contact of the guard-cells. The resulting shape, which may be seen from the figures, seems to be fairly constant for all modern Cycads, and is very characteristic, being modified in the case of the deeply sunken forms. The same type of stoma is seen in most of the fossil fronds examined.

The Bennettitalean stomata possessed guard-cells with very peculiar thickenings, but which are referable to the same general type as exhibited by the modern fronds. We had a central thin region round the pore and two fusiform or crescentic patches on either side (compare the figures of *Zamites*, Pl. 19. fig. 2, and *Otozamites*, Pl. 19. fig. 3, with figures of *Zamia*, text-fig. 21, and *Stangeria*, Pl. 17. fig. 1).

We have to remember, however, that in looking at the recent stomata the most noticeable thickening is due to the lignification of the guard-cells, while in the fossil forms presumably nothing but cutin remains. In order to get preparations of the recent structures which are really comparable with the fossils, we may get rid of the woody tissue by continued boiling in Schultz's solution; in the resulting preparations little but cuticle will be left. Preparations made in this way show considerable similarity with the fossil forms (*cf.* text-fig. 32 with figures on Plate 19), the woody thickening seems to have been accompanied by a certain amount of cuticularisation which is especially seen at the poles, while on either side of the pore the thickenings on the upper surface of the guard-cells are seen as the spindle-shaped patches which

are so characteristic in the fossil forms; the thin areas between the pores and the cusp-shaped thickenings are the same as are seen in the fossil forms, especially in *Otozamites*, *Anomozamites*, and *Tæniopteris*.

The polar thickenings of the guard-cells which are seen in most of the recent forms, and which appear especially marked in *Stangeria* (where they consist of both woody tissue and cuticle), are almost entirely absent in the fossil fronds, and present a point of difference.

It is probable that the relations of the guard-cells to the surrounding epidermal cells and subsidiary cells, offer one of the most important points of comparison which can be obtained in the surface view. In the Bennettitalean fronds we have a very constant and characteristic arrangement. On either side of the guard-cells there are two large subsidiary cells of exactly the same diameter as the guard-cells and probably developed from the same mother-cell. [The small cells seen at the poles of the slit in some cases, *e. g.* in *Anomozamites*, Pl. 19. fig. 5, are probably not polar subsidiary cells but thin places above the poles of the actual guard-cells.] In the Nilssoniales group, however, and in recent fronds, the subsidiary cells are very different. They are usually six or

Fig. 32.

Stomata of *Stangeria paradoxa* as seen in surface view after prolonged maceration. Notice that the lateral upper thickenings of the guard-cells show as spindle-shaped patches, while there appear to be empty cell-like spaces at the poles. Cf. with figures of *Ptilophyllum* and *Otozamites* stomata.

more in number, and are arranged more or less radially round the guard-cells. In *Stangeria*, *Encephalartos*, and other forms, we find two large lateral and two smaller polar subsidiary cells, but they often become subdivided. Thus in this respect the Bennettitales are quite distinct from the recent Cycads. On the other hand the fronds included in the Nilssoniales may be closely compared with the modern forms in this character, the numerous overarching subsidiary cells of *Nilssonia*, with their strongly cuticularised points, sometimes coalescing to form a solid ring, being comparable with those of *Dioon* and *Cycas*. These overarching cells frequently are of the same type as those forming the remainder of the epidermis in both the fossil and the recent forms.

It is unfortunate that we cannot get more evidence as to the form and structure of the guard-cells in the Nilssoniales group, though in the case of the example of *Ptilozamites* figured (Pl. 19. fig. 16) the guard-cells seem to have been of somewhat the same type as those of the Bennettitales.

In comparing the general epidermal cells of the recent and fossil types we may notice that there is nothing among the modern forms which is really comparable to the very sinuous cells of *Ptilophyllum* and *Anomozamites*. The cell-walls of the *Stangeria* epidermis are somewhat sinuous, and this is of interest when taken with the forking nervation of its pinnæ, another point of comparison with the Bennettitales. If the sinuous cell-walls are to be regarded as a feature of any phylogenetic importance, it may indicate the relationships of the Bennettitales with the Pteridosperms and ferns. On the other hand, the straight walls of the Nilssonian epidermal cells, together with the shape of the cells, are additional points of agreement with the majority of the modern Cycads.

We may summarize generally the relationship between the fossil and modern Cycadean fronds as indicated by this enquiry, as follows. The cuticular structures indicate that the Bennettitalean fronds form an easily recognizable group containing the majority of the fronds previously described as Cycadean, which are quite distinct from the modern fronds and from those of the Nilssoniales group. The latter, which includes the genera *Nilssonia*, *Ctenis*, and *Ptilozamites*, are somewhat closely allied to the modern fronds, and may probably be regarded as the true Mesozoic Cycads and the ancestors of the modern forms.

On the other hand, the Bennettitalean stomata show some important resemblances to the Cycadean ones, and may well point to a parallel line of evolution from a common stock.

It is scarcely possible at the present time to make any detailed comparison between the epidermal structures which have been described in this paper and those of the other groups of Gymnosperms, since in no case have the latter been studied in sufficient detail. A few general comparisons may, however, be attempted. It seems probable that of the other Gymnosperms, all of which possess guard-cells with the same characteristic pear-shaped section, the Gnetales are the most closely allied in the form of their cuticular structures. In fact, it would seem that the Bennettitalean cuticles are more closely allied to those of some of the Gnetums than to any other group, the points of comparison being the frequent arrangement of the stomata in rows, the relations of the guard-cells to the subsidiary cells, and the method of thickening of the guard-cells. In *Gnetum Gnemon* the walls of the epidermal cells are often sinuous. We have been able to examine the cuticles of *Welwitschia* and *Gnetum Gnemon*, and the stomata in these forms have also been described by Takeda* and others, but other species need to be examined before the comparison can be pressed.

The stomatal structures of *Ginkgo*† are somewhat similar to those of *Nilssonia*, but differ in the smaller and more constant shape and number of the subsidiary cells and in the very slight thickening of the guard-cells. The stomata of the Conifers‡, while possessing guard-cells which are thickened in a somewhat similar way, seem to differ from those of the Cycadales and Bennettitales in the uniformity of the thickening and

* Takeda (1913), p. 350, text-figs. 1-5, pl. xxix. figs. 2-6.

† Seward (1911'), pl. v. figs. 59-62.

‡ Mahlert (1885), figs. on Taf. 1.

in the spreading out of the woody lamellæ at the poles into an almost continuous investment without the characteristic thin patches. The relations of the guard-cells to the surrounding epidermal cells also seem to be another important feature of difference.

It would seem, then, from the foregoing study, that the characters presented by the stomata and epidermal cells of the Gymnosperms can be regarded as important as indicating to some extent their relationships. In the Cycadean alliance we find some characters which have undergone comparatively little modification in the ages which have elapsed between Jurassic times and the present day, and we are consequently entitled to regard stomatal structure as being the expression of ancestral characters rather than of purely local and temporary conditions of environment. Of course, further study of other groups is needed, and we hope that it will be possible in the future to make further researches upon this line of enquiry.

In conclusion, may we express our thanks to Professor A. C. Seward for his help and advice during the course of this work, also to Professor Nathorst and other friends who have given us information and assistance.

BIBLIOGRAPHY.

- BATHER, F. A. (1908).—"Nathorst's Methods of studying the cutinised portions of Fossil Plants." Geol. Mag., Dec. v. vol. v. p. 454.
- BERRY, E. W. (1911).—Maryland Geological Survey. Lower Cretaceous. Baltimore.
- BORNEMANN, J. G. (1856).—"Ueber organische Reste der Lettenkohlengruppe Thüringens." Leipzig.
- CHAMBERLAIN, C. J. (1912).—"Two Species of *Bowenia*." Bot. Gaz. vol. liv. p. 419.
- HILL, T. G. (1912).—"Scharlach R: a Microchemical Test for Oils." New Phytologist, vol. xi. p. 72.
- KARZEL, R. (1908).—"Die Verholzung der Spaltöffnungen bei Cycadeen." Wiesner-Festschrift, p. 510.
- KRAUS, G. (1866).—"Ueber den Bau der Cycadeenfiedern." Prings. Jahrb. vol. iv. p. 305.
- LIGNIER, O. (1894).—"Structure et Affinités du *Bennettites Morierei*, Sap. et Mar." Mém. Soc. Linnéenne de Normandie, vol. xviii. Caen.
- (1912).—"Stomates des Écailles interséminales chez le *Bennettites Morierei* (Sap. et Mar.)." Bull. Soc. Bot. de France, vol. lix. p. 425.
- LLOYD, F. E. (1908).—"The Physiology of Stomata." Carnegie Institute.
- MAHLERT, A. (1885).—"Beiträge zur Kenntnis der Anatomie der Laubblätter der Coniferen, mit besonderer Berücksichtigung des Spaltöffnungs-Apparates." Bot. Centralbl. vol. xxiv. p. 54.
- NATHORST, A. G. (1880).—"Berättelse etc." Öfvers. K. Vetensk.-Akad. Förhandl. No. 5, p. 35.
- (1907).—"Paläobotanische Mitteilungen 1 & 2." K. Svenska Vet.-Akad. Handl. Bd. xlii. No. 5.
- (1908).—"Paläobotanische Mitteilungen 4." K. Svenska Vet.-Akad. Handl. Bd. xliii. No. 6.
- (1909).—"Ueber die Gattung *Nilssonia*, Brongn." K. Svenska Vet.-Akad. Handl. Bd. xliii. No. 12.
- (1912).—"Einige paläobotanische Untersuchungsmethoden." Paläobotanische Zeitschrift, Bd. i. p. 26.
- NESTLER, A. (1895).—"Ein Beitrag zur Anatomie der Cycadeenfiedern." Prings. Jahrb. vol. xxvii. p. 341.
- PHILLIPS, J. (1829).—Illustrations of the Geology of Yorkshire. York.

- PORSCH, O. (1905).—Der Spaltöffnungsapparat im Lichte der Phylogenie. Jena.
- SCHENK, A. (1867).—Die fossile Flora der Grenzsichten des Keupers und Lias Frankens. Wiesbaden.
- (1871).—Die fossile Flora der Nordwestdeutschen Wealdenformation. Cassel.
- SCHIMPER, W. P. (1869).—Traité de Paléontologie végétale. Vol. i.
- SCHWENDENER, S. (1881).—"Ueber Bau und Mechanik der Spaltöffnungen." Monatsber. d. Akad. d. Wissenschaft. Berlin.
- SEWARD, A. C. (1900).—The Jurassic Flora. I. The Yorkshire Coast. British Museum Catalogue. London.
- (1911¹).—"Jurassic Plants from Chinese Dzungaria." Mém. Comité Géologique, n. s., livr. lxxv. St. Petersburg.
- (1911²).—"The Jurassic Flora of Sutherland." Trans. Roy. Soc. Edin. vol. xlvii. pt. iv. no. 23.
- (1912).—"A petrified *Williamsonia* from Scotland." Phil. Trans. ser. B. vol. cccii. p. 101.
- STRASBURGER, E. (1867).—"Ein Beitrag zur Entwicklungsgeschichte der Spaltöffnungen." Prings. Jahrb. v. p. 297.
- TAKEDA, H. (1913¹).—"Some Points in the Anatomy of the Leaf of *Welwitschia mirabilis*." Annals of Botany, vol. xxviii. p. 347.
- (1913²).—"Development of the Stoma in *Gnetum Gnemon*." Annals of Botany, vol. xxvii. p. 365.
- THOMAS, H. HAMSHAW (1911).—"On the Leaves of Calamites." Phil. Trans. ser. B, vol. ccii. p. 51.
- (1912).—"On some methods in Palæobotany." New Phytologist, vol. xi. p. 109.
- (1913).—"The Fossil Flora of the Cleveland District of Yorkshire. I. The Flora of the Marske Quarry." Quart. Journ. Geol. Soc. vol. lxix. p. 223.
- WIELAND, G. R. (1911).—"On the Williamsonian Tribe." Amer. Journ. Sci. vol. xxxii. p. 433.
- ZEILLER, R. (1882).—"Observations sur quelques Cuticules fossiles." Ann. des Sci. Nat. sér. 6, Bot. tome xiii.
- (1906).—Bassin houiller et permien de Blanzky et du Creusot. Fasc. 2. Flore fossile. Etudes des gîtes minéraux de la France. Paris.

EXPLANATION OF PLATES 17-20.

[The figures in Plates 17-19 were drawn with the aid of a camera lucida, the magnification being 450 diameters, unless otherwise indicated. The following abbreviations are used:—*p*, polar subsidiary cells; *l*, lateral subsidiary cells; *d*, dorsal lignified lamellæ of the guard-cells; *v*, ventral lignified lamellæ; *g*, guard-cells; *s*, stoma-opening; *i*, intercalated subsidiary cells; *a*, air-space.]

PLATE 17.

- Fig. 1. *Stangeria paradoxa*. Lower cuticle in surface view, showing a stoma with its characteristically thickened appearance due to the lignified lamellæ of the guard-cells. Note the radiating striæ at the poles of the stoma, due to folds in the cutin layer.
- Fig. 2. The same. A hair-scar from the under side of the leaf in surface view, showing the radiating folds of the cutin.
- Fig. 3. The same. Upper cuticle in surface view, showing the characteristic undulated walls of the epidermal cells and the striations due to folds in the cutin.
- Fig. 4. The same. Upper cuticle in section. Note the irregular appearance of the cutin layer due to folding.

- Fig. 5. The same. Lower cuticle, showing a section through a stoma. Note the position of the guard-cells, almost at the same level as the subsidiary cells.
- Fig. 6. The same. Lower cuticle, showing a section through a hair-scar.
- Fig. 7. *Bowenia spectabilis*, var. *serrulata*. Lower cuticle in surface view, showing a stoma; the thickened appearance is due to the lignified lamellæ of the guard-cells.
- Fig. 8. The same. A stoma of the upper side of the leaf in surface view. Note the presence in this case of six subsidiary cells, two of which are polars and four laterals.
- Fig. 9. The same. Lower cuticle, showing a section through a stoma. Note the position of the guard-cells with regard to the subsidiary cells.
- Fig. 10. *Cycas revoluta*. Upper cuticle in section, showing the pittings of the lateral and outer walls of the epidermal cells.
- Fig. 11. The same. Upper cuticle in surface view, showing the pittings of the lateral and outer walls of the epidermal cells.
- Fig. 12. The same. Lower cuticle in surface view, showing the fifteen subsidiary cells surrounding a stoma-opening. The shading indicates the raising of the subsidiary cells beyond the general epidermal level (cf. fig. 13).
- Fig. 13. The same. Lower cuticle, showing a section through a stoma. Note the presence of an intercalated series of cells between the guard and subsidiary cells, and the large air-space formed by the sinking of the guard-cells and the overarching of the subsidiary cells.
- Fig. 14. *Cycas circinalis*. Lower cuticle, showing a section through a stoma. Note the sinking of the guard-cells and the absence of an intercalated series of cells (cf. figs. 5, 9, & 13).
- Fig. 15. The same. Upper cuticle in section, showing pitting of the outer walls of the cells.

PLATE 18.

- Fig. 1. *Ceratozamia mexicana*. Upper cuticle in surface view, showing thin-walled and thick-walled cells. Note the layering and pitting of the thickened walls.
- Fig. 2. The same. Lower cuticle in surface view, showing a stoma-opening surrounded by six subsidiary cells, two of which are polars and four laterals. The cell-walls are pitted and layered as in fig. 1.
- Fig. 3. The same. Upper cuticle, showing the thick- and thin-walled cells in section.
- Fig. 4. The same. Lower cuticle: a section through a stoma, showing one intercalated series of cells between the guard-cells and subsidiary cells.
- Fig. 5. *Dioon edule*. Upper cuticle in surface view, showing the variation in thickness and the layering of the cell-walls.
- Fig. 6. The same. A section of a stoma. Note the presence of three intercalated series of cells (i^1-i^3) between the guard-cells and the subsidiary cells; also the dorsal and ventral lignified lamellæ of the guard-cells.
- Fig. 7. The same. Upper cuticle in section, showing variation in the thickness of the walls. Note the heavy cutin layer.
- Fig. 8. The same. A section of the lower cuticle, showing a crystal-containing space between two epidermal cells. The space is lined and covered by cutin.
- Fig. 9. The same. The lower cuticle, showing a crystal-containing space in surface view.
- Fig. 10. *Encephalartos Lehmanni*. Upper cuticle in section, showing pitting of the lateral walls.
- Fig. 11. *E. villosus*. Upper cuticle in surface view, showing slight pitting of the lateral walls of the cells and septation of an elongated cell at *x*.
- Fig. 12. *Zamia integrifolia*. Upper cuticle in surface view, showing variation in the thickness of the walls and occasional pitting of the thin-walled cells.

Fig. 13. *Zamia muricata*. Lower cuticle, showing a stoma in section. Note the slightly sunk guard-cells.

Fig. 14. *Macrozamia cylindrica*. Section through a stoma, showing two intercalated series of cells and the long narrow air-space.

Fig. 15. The same. Upper cuticle in surface view, showing variation in the thickness of the cell-walls, also pitting and layering of the walls.

PLATE 19.

[Camera lucida drawings of stomata of fossil fronds.]

Fig. 1. *Ptilophyllum pecten*. Showing the different thickenings of the guard-cells and the thin places above the poles. Specimen from Marske. $\times 600$.

Fig. 2. *Zamites* sp. Showing outlines of stomatal thickenings. From scale-leaves on stem. Specimen in the Naturhistoriska Riksmuseum, Stockholm. $\times 500$.

Fig. 3. *Otozamites Feistmanteli*. Stoma with differences of thickening less marked. The subsidiary cells are not thickened. Specimen from Marske. $\times 500$.

Fig. 4. *Otozamites graphicus*. The sides of the pore appear to have been slightly cuticularised. Specimen from Marske. $\times 500$.

Fig. 5. *Anomozamites Nilssoni*. Showing thin subsidiary cells and thin places at the poles. Specimen from Gristhorpe. $\times 500$.

Figs. 6-8. *Dictyozamites Hawelli*. Subsidiary cells thickened: 6 and 8 show projections from upper thickenings of guard-cells. Fig. 7. Stoma seen from mesophyll side. Specimens from Marske. $\times 500$.

Fig. 9. *Teniopteris major*. Compare with *T. vittata*. Specimen from Gristhorpe. $\times 500$.

Figs. 10-12. *Teniopteris vittata*. Showing the similarity with stomata of other Bennettitalean fronds. Fig. 10 is seen from the mesophyll side. 10 and 11 from Gristhorpe, 12 from Marske. $\times 500$.

Figs. 13, 14, 18. *Nilssonsonia* sp. From a specimen found at Roseberry Topping. $\times 500$. Note in 13 the series of overarching subsidiary cells. Figs. 14 and 18 show the guard-cells as seen from the mesophyll side of the epidermis.

Fig. 15. *Nilssonsonia mediana*. Subsidiary cells with thickened tips overarching the guard-cells. Specimen from Marske. $\times 500$.

Figs. 16, 17. *Ptilozamites Leckenbyi*. Stomata as seen from upper (17) and lower side (16). In the former the guard-cells can just be seen through the opening. Specimens from Gristhorpe. $\times 500$.

Fig. 19. *Ctenis falcata*. Showing thick wall of cuticle round the stomatal cavity. Specimen from Gristhorpe. $\times 500$.

PLATE 20.

[Photomicrographs of cuticular preparations of fossil fronds.]

Fig. 1. *Ptilophyllum pecten*. Showing upper epidermis (to the right) and lower epidermis with the stomata arranged in regular rows. From Marske. $\times 25$.

Fig. 2. *Ptilophyllum pecten*. Upper epidermis more highly magnified, showing sinuous outlines of the cells. $\times 170$.

Fig. 3. *Ptilophyllum pecten*. Part of lower epidermis more highly magnified, showing stomata. $\times 200$.

Fig. 4. *Ptilophyllum* sp. Showing rows of circular hair-scars with epidermal cells below. From Marske. $\times 200$.

- Fig. 5. *Teniopteris vittata*. Part of upper epidermis, showing characteristic folded cell-walls. From Gristhorpe. $\times 200$.
- Fig. 6. *Teniopteris vittata*. Part of lower epidermis, showing stomata which were slightly sunken, and hence the walls of the epidermal cells are out of focus. $\times 200$.
- Fig. 7. *Dictyozamites Hawelli*. Lower epidermis, showing stomata. The characteristic form of the upper thickening lamellæ is seen in a few cases. From Marske. $\times 180$.
- Fig. 8. *Anomozamites Nilssoni*. Lower epidermis with stomata. From Gristhorpe. $\times 200$.
- Fig. 9. *Nilssonia* sp. From Roseberry Topping. Upper epidermis, showing characteristic papillæ. $\times 180$.
- Fig. 10. *Nilssonia mediana*. Lower epidermis, showing elongated cells with papillæ occurring below the veins, while the cells of the intermediate areas are rounded. From Marske. $\times 200$.
- Fig. 11. *Nilssonia orientalis*. Portion of upper epidermis from specimen No. 323 in the Naturhistoriska Riksmuseum, Stockholm. Found at Cloughton Wyke. $\times 200$. This specimen has been figured as *Nilssoniopteris tenuinervis*.
- Fig. 12. *Ptilozamites Leckenbyi*. Part of lower epidermis, showing straight-walled cells and stomata. From Gristhorpe. $\times 200$.
- Fig. 13. *Ctenis falcata*. Part of lower epidermis with stomata. From Gristhorpe. $\times 200$.

N. B. del.

CYCADEAN CUTICLES.

Grout, sc.

N. B. del.

Grout, sc.

CYCADEAN CUTICLES.

1

2

3

4

5

6

7

8

9

10

11

12

13

ERRATA.

Transactions of the Linnean Society, Ser. 2, Botany, Vol. VIII. Part 5.

Page

179, line 17 should read "PTILOPHYLLUM PECTEN (Phill.) Morris."

182 „ 5, for *Stangeria* read *Ptilophyllum*.

186 „ 1 should read "OTOZAMITES GRAPHICUS (Leck.) Schimp."

187 „ 17 „ "ANOMOZAMITES NILSSONI (Phill.) Sew."

190 „ 24 „ "NILSSONIA COMPTA (Phill.) Bronn."

191 „ 4 from bottom should read "NILSSONIA MEDIANA (Leck.) Fox-Strangw."

194 „ 7 should read "PTILOZAMITES LECKENBYI (Leck.) Nath."