THE ACTION OF RADIUM ON
CATARACTS *

MARTIN COHEN, M.D.

Professor of Ophthalmology, New York Post-Graduate Medical
School and Hospital

AND

ISAAC LEVIN, M.D.

Clinical Professor of Cancer Research, University and Bellevue
Hospital Medical College; Chief of the Department of
Cancer Research, Montefiore Home and Hospital

NEW YORK

Until recently there was no agent known that could in the slightest degree change or influence the natural course of the development of a cataract from its period of incipency to maturity. Surgery had reached a high degree of perfection and was followed in the majority of cases by a clinical success which meant restoration of vision. Surgery, however, does not influence or change the development of the lenticular opacification but simply removes the entire lens.

A great many attempts have been made to influence the development of cataract by chemical means, but as yet without success. In the last decade, however, a great deal of progress has been reported in radium therapy as applied to various pathologic conditions, and the field of usefulness of the biologic action of radium in the domain of therapeutics is constantly widening. Furthermore, a great many eye conditions, such as trachoma, vernal catarrh, and various tumors of the eyeball and adnexa, have been treated with radium without in any way impairing the normal structure of the eyeball or impairing its function.

One of the writers, who has done some research on the subject of chemotherapeutics of the cataract with entirely negative results, raised the question whether one might not expect that the radioactive substances may exert an influence on the development of the cataract. An *a priori* consideration of the subject showed the likelihood of such an influence by the radium rays.

As admitted by the best authorities on the subject, the capsular epithelium is most probably the starting point of cataract. Indeed, it is inconceivable that the change, or the precipitation, as it were, of certain soluble ingredients within the lens which result in the formation of the opacifications could take place without a change in the structure of the barrier membrane, which the capsule is; and since the capsule is essentially a product of cell life, the change in the barrier must be due in the ultimate analysis to an abnormality in either the structure or the function of these cells.

There is ample biologic and clinical evidence that radium produces various and very deep changes in cellular functions. In tumors and granulomas, the action of the rays consists in the inhibition of the proliferating capacity of the cells; in conditions like exophthalmic goiter, it influences and alters the functions of the cell. It is thus evident that the radium rays may alter or influence the capsular epithelium in such a manner as possibly to render the whole capsule less permeable for the abnormal agents within the general circulation or in the neighboring tissues, which later on entering the lens produce the opacifications; or else the change in the membrane may consist in its

enhanced ability to exosmose certain substances from the lens, and the lack of the latter may result in formation of opacities.

A thorough search of the literature showed that there had been no attempt made by any investigator to influence cataract with radium, with the exception of Koster.¹ This investigator applied radium to a great variety of eye diseases, apparently in a large number of cases. While he does not state definitely the number of cases of each disease treated, he reports on over twenty diseases, and as regards some of these diseases he distinctly states, "I have treated as many as twenty or more patients." Among others, he reports on treatment of cataract, with negative results, but states that in a case of cataracta caerulea—the star form in the anterior cortex—the opacities fell to pieces and disappeared entirely after five sittings of one hour weekly.

The method of radium therapy employed by Dr. Koster is not in accord with the modern conception of the subject. None the less, he apparently obtained favorable results in certain conditions, and there was no deleterious effect observed on the retina or on the other ocular structures. Dr. Koster used a glass tube filled with radium which he placed, uncovered and unprotected in any way, directly to the scleral conjunctiva. This is a dangerous method, since all of the soft rays remained unfiltered and could irritate and burn the adjacent tissue. In this connection, it is remarkable that an amount of radiation which would have surely caused a burn on the skin did not cause a burn of the scleral conjunctiva. It must be admitted that he used much smaller quantities than are employed at present. The therapeutic results obtained by Koster, notwithstanding his inadequate methods, the harmlessness of the procedure, which was shown by him as well as by the investigators who treated epitheliomas of the eyelids, and also the theoretical considerations mentioned above encouraged us to commence this investigation.

METHODS EMPLOYED BY THE AUTHORS

The methods employed were very similar to the technic of deep radium therapy which one of us is using at present in the treatment of malignant tumors and various other conditions. For a clear understanding of these methods it is necessary to explain here very briefly certain physical characteristics of radium.

Radium emits alpha rays, beta rays, and gamma rays. The first two act only as a general caustic on all the tissues found on the surface to which the radium is applied. The hardest beta, and especially the gamma rays, have a selective action which causes their efficacy in the treatment of deep conditions. The selectivity of the rays consists in the fact that they may influence pathologic structures while leaving the neighboring normal tissues practically intact. These gamma rays represent only about 1 per cent. of the total amount of radiations. When Koster employed in his tube 2 mg. in gamma radiations to do the work, there were possibly not enough soft radiations to injure the normal tissue; but, on the other hand, he had only an extremely small amount of gamma radiations to do the actual work. In modern technic, ten, twenty and more times the amount of radium used by Koster is employed. With the aid of various means which it is not necessary to discuss here, all the irritating soft

* Read before the Section on Ophthalmology at the Seventieth Annual Session of the American Medical Association, Atlantic City, N. J., June, 1919.

1. Koster, cited by A. F. Mattice: Arch. Ophth. 43: 237, 1914.

ACTION OF RADIUM ON CATARACT

Name	Age	Type of Cataract	Location	No. of Ap- plications	Date or Visits		Radium Application					Results
							Vision		Lens Opacification		Fundus Before and After	
					First	Last	Before	After Last	Before	After Last		
J. V. 1	45	Senile immature (right)	Anterior cortex	30	Mar. 20/18	Mar. 31/19	Counts fingers at 1 ft.	Counts fingers at 3½ ft.				Vision, improved Lens Opacity, diminished Fundus, no de- terioration
J. P. 2	75	Senile immature	Central and posterior cortex	32	Apr. 2/18	Mar. 31/19	20 70	20 50				Vision, improved Lens Opacity, stationary Fundus, no de- terioration
O. M. 3	33	Complicated (retinitis pigmentosa) immature	Anterior and posterior cortex	40	May 19/18	Apr. 1/19	Counts fingers at 3½ ft.	20 — 70				Vision, improved Lens Opacity, diminished Fundus, no de- terioration
J. D. 4	50	Complicated (irido- cyclitis) immature	Anterior cortex	33	June 8/18	Mar. 31/19	Counts fingers at 8 in.	Counts fingers at 3 ft.				Vision, improved Lens Opacity, diminished Fundus, no de- terioration
F. S. 5	58	Senile immature	Anterior cortex	14	June 19/18	Mar. 31/19	Counts fingers at 3 ft.	Counts fingers at 3 ft.				Vision, stationary Lens Opacity, diminished Fundus, no de- terioration
J. K. 6	47	Senile immature	Posterior cortex	23	June 19/18	Mar. 31/19	20 30 + 1	20 15 + 2				Vision, improved Lens Opacity, diminished Fundus, no de- terioration
L. G. 7	52	Senile immature	Posterior cortex	28	July 21/18	Mar. 31/19	20 100 — 1	20 30				Vision, improved Lens Opacity, diminished Fundus, no de- terioration
A. R. 8	62	Complicated (glaucoma) immature	Anterior cortex	36	July 25/18	Mar. 3/19	20 40 — 1	20 20 — 4				Vision, improved Lens Opacity, stationary Fundus, no de- terioration
F. G. 9	75	Senile immature	Posterior cortex	29	Aug. 13/18	Mar. 31/19	With correc- tion 20 — + 2 20	With correc- tion 20 — + 2 20				Vision, stationary Lens Opacity, stationary Fundus, no de- terioration
C. G. 10	62	Senile immature	Central and posterior cortex	21	Aug. 29/18	Mar. 31/19	20 100 — 1	10 200				Vision, decreased Lens Opacity, diminished Fundus, no de- terioration
S. K. 11	50	Senile immature	Central and anterior cortex	23	Sept. 3/18	Mar. 20/19	20 40	20 30				Vision, improved Lens Opacity, diminished Fundus, no de- terioration
C. R. 12	64	Senile immature	Anterior and posterior cortex	21	Sept. 5/18	Mar. 20/19	20 200	20 200 — 1				Vision, stationary Lens Opacity, diminished Fundus, no de- terioration

ACTION OF RADIUM ON CATARACT.—CONTINUED

Name	Age	Type of Cataract	Location	No. of Applications	Date of Visits		Radium Application					Results
							Vision		Lens Opacification		Fundus Before and After	
					First	Last	Before	After Last	Before	After Last		
R. K. 13	74	Senile immature	Anterior cortex	27	Sept. 7/18	Mar. 20/18	20 — 70	20 — 50				Vision, improved Lens Opacity, stationary Fundus, no deterioration
L. L. 14	53	Senile immature	Anterior cortex	8	Sept. 19/18	Mar. 20/18	20 — 200	20 — 100				Vision, improved Lens Opacity, diminished Fundus, no deterioration
I. S. 15	49	Senile immature	Posterior cortex	20	Sept. 25/18	Mar. 20/19	20 — 30	20 — 15				Vision, improved Lens Opacity, diminished Fundus, no deterioration
R. M. 16	56	Senile immature	Posterior cortex	13	Sept. 20/18	Mar. 20/19	20 — 100	20 — 50				Vision, improved Lens Opacity, diminished Fundus, no deterioration
I. W. 17	73	Senile immature	Posterior cortex	17	Nov. 7/18	Mar. 10/19	Counts fingers at 6 ft.	20 — 50				Vision, improved Lens Opacity, diminished Fundus, no deterioration
A. R. 18	62	Senile immature	Anterior cortex	15	Nov. 7/18	Mar. 16/19	20 — 30	20 — 30				Vision, stationary Lens Opacity, disappeared Fundus, no deterioration
J. V. 19	45	Senile immature (left)	Anterior cortex	4	Jan. 29/19	Mar. 25/19	20 — 40	20 — 30				Vision, improved Lens Opacity, diminished Fundus, no deterioration
P. S. 20	38	Complicated (glaucoma) immature	Anterior cortex	9	Jan. 19/19	Mar. 25/19	Counts fingers at 6 ft.	20 — 30				Vision, improved Lens Opacity, diminished Fundus, no deterioration
H. D. 21	48	Senile immature	Central and posterior cortex	7	Jan. 31/19	Mar. 20/19	20 — 40	20 — 40				Vision, stationary Lens Opacity, diminished Fundus, no deterioration
F. I. 22	56	Senile immature	Anterior and posterior cortex	6	Feb. 1/19	Mar. 20/19	Counts fingers at 4 ft.	Counts fingers at 4 ft.				Vision, stationary Lens Opacity, stationary Fundus, no deterioration
M. A 23	58	Senile immature	Posterior cortex	5	Feb. 1/19	Mar. 20/19	10 — 200 myopia	10 — 200				Vision, stationary Lens Opacity, diminished Fundus, no deterioration
P. M. 24	67	Senile immature	Posterior cortex	5	Feb. 13/19	Mar. 20/19	20 — 70	20 — 70				Vision, stationary Lens Opacity, stationary Fundus, no deterioration

rays are filtered off and only the selectively acting gamma rays are utilized. The actual technic consisted in covering the radium with brass, photographic paper and gauze, and placing it over the closed eyelid. The distance between the radium substance and the eyelid was about 2 cm. The application usually lasted two hours.

The main difficulties that an investigator encounters when he attempts to test the efficacy of any measure on the development of a cataract is the fact that the disease is frequently very slow in progressing and may remain stationary for a long period. It is even assumed by some authors that occasionally a cataract may spontaneously improve for a time. None the less, it is the experience of every ophthalmologist that by far the greater number of senile cataracts follow the usual course and mature in a period of from six months to two years.

A statistical analysis of 131 cases reported by Dr. Nevens of the Liverpool Eye and Ear Infirmary is very characteristic of the usual experience. Of his cases, seven, or 5 per cent., matured in less than one month; twenty-six, or 20 per cent., matured in a period of between one and six months; thirty, or 23 per cent., matured within one year; thirty-five, or 27 per cent., matured in two years; nine cases, or 7 per cent., in three years; seventeen, or 13 per cent., in between three and seven years; and seven cases, or 5 per cent., in between twelve and twenty years, thus giving a total of 75 per cent. that matured within the first two years and only 18 per cent. that took more than three years to mature.

It is thus clear that in a certain number of consecutive cases selected for study, not less than three fourths of such cases, in the natural course of events, would reach maturity within two years, and would consequently have to show a good deal of progression within the first year.

The material used for the present investigation consisted of all the cases of immature cataracts seen by one of us in his routine clinical and private practice during the time of the investigation, twenty-four cases being included.

Twenty patients had senile cataracts of the usual type. In the majority of the cases, both eyes were affected, but only one eye, which was less mature, was treated. Four cataracts were complicated; one was due to a retinitis pigmentosa, one to an iridocyclitis, and in two instances chronic glaucoma was present. An iridectomy was performed in the latter cases, but the cataract was not due to any trauma. None of the patients suffered from true diabetes or nephritis, with the exception of one who showed sugar, traces of albumin, and a few hyaline casts. The type, extent and location of each cataract is noted on the accompanying chart.

Three of the patients received applications of radium for one year; three for eleven months; two for ten months; two for nine months; six for eight months; two for seven months; three for four months, and three for three months.

The result of the radium treatment in these cataracts as regards vision and lens opacification is as follows: There was a combined improvement in the lenticular opacification and vision in twelve cases; lens opacification disappeared in one case without improvement in vision; opacification diminished in five cases with stationary vision (in one case vision decreased); opacifi-

cation was stationary but vision improved in three cases; both vision and opacification remained stationary in three cases. In a word, twenty-one cases, or 87.5 per cent., showed some improvement.

The method of procedure in the examination of these cases consisted of the usual methodical examination of the eye, including also an examination of the urine, and physical and blood examinations when deemed essential.

The examination of the visual acuity and the drawings of the lenticular opacities required especial attention to several details, as follows: The illumination of the room and the test charts was of the same character at each examination. Several corresponding letters, numbers, and illiterate charts were used. After the results of the visual acuity test, functional test, and the refraction were determined and recorded in each case, the pupils were dilated with a 1 per cent. homatropin solution for the purpose of examining the details of the fundus and also for a detailed drawing of the opacification. A combined focal examination was made of the lenticular opacity, namely, with a condensing lens and a corneal loop, and then a drawing of this opacity was recorded. The cataract was next observed with the Loring ophthalmoscope through its aperture, the patient always fixing a definite object in space, in order to procure a good view of the entire opacification, and a drawing of it was made. A 20-diopter lens was then inserted in the aperture of the ophthalmoscope for a drawing of the finer details of the lens opacification. In order that the latter examination should be accurate, a distinct fixed opacity of the lens was noted at a definite angle with a strong convex lens in the ophthalmoscope. Then another drawing was made of the opacification; this drawing also served as a guide for future examinations. Finally a drawing was made of the fundus details with the aid of a Loring ophthalmoscope. In a certain number of cases an electric ophthalmoscope was necessary to view the fundus, as the opacity was very dense. Four drawings were made of each case before radium was applied, and subsequent drawings were made at stated intervals.

The chart presents graphically the type of the cataract, the extent of the opacification, the length of time during which the treatment was continued, the number of treatments given, and the results obtained, so that it is hardly necessary to give in the text a detailed history of each individual case. Therefore only some of the cases which present salient features will be discussed in greater detail.

REPORT OF CASES

CASE 1.—Senile Cataract.—Examination before the application of radium gave the following results: The patient was able to count fingers at 1 foot. With focal examination the lens had a diffuse grayish appearance; with the ophthalmoscope a faint fundus reflex was obtained through the periphery of the lens. Five days after treatment began the patient could count fingers at 3 feet; ophthalmoscopic examination of the lens showed it to have a diffuse grayish aspect, while from its periphery a large fundus reflex was visible. Seven days after the beginning of treatment the patient could count fingers at 4 feet; oblique examination showed the lens to be of a grayish color, having a few whitish dots and small striae scattered near the central area; the ophthalmoscope revealed a much larger fundus reflex, with distinct sector striations radiating from the periphery of the lens. The improvement obtained remains to date, and consequently there has been no progression of the disease during a period of ten months. It is of particular interest to note that in this case

the vision today, ten months after the beginning of radium application, is better than before the radium treatment began; furthermore, a nearly complete fundus reflex is now obtainable, whereas before the application of the radium only at a narrow rim around the periphery was a faint reddish reflex transmitted from the fundus.

CASE 3.—Complicated Cataract (Retinitis Pigmentosa).—The patient gave a history of syphilitic infection occurring seventeen years previously. He received antisyphilitic treatment for several years, but had had none for two years before coming under our observation. The Wassermann blood reaction was positive. Disturbance in vision was first noticed thirteen years before; the vision gradually deteriorated, regardless of frequent syphilitic treatment. The first examination was made on May 19, 1918. The vision of the right eye was 20/200; with the left eye the patient counted fingers at 3½ feet. The field was markedly contracted concentrically; the fundus details, though cloudy, showed the typical picture of the disease. Ophthalmoscopic examination of the lens and fundus showed a distinct fundus reflex, which allowed the lenticular opacifications to be studied; they consisted of anterior and posterior cortical opacifications; the anterior had a sector arrangement, the posterior was curvilinear, and in between these opacifications a fine granular appearance was seen with here and there evidences of small globules or vacuoles. One week after the beginning of the radium application the left eye was examined, and the condition found was as follows: Vision equaled counting fingers at 7 feet. The fundus details were more clearly seen, the striations were less apparent, and the vacuoles or fine globules were more evident. No change was noticed in the papilla or in the retinal pigment. Nine days later the examination showed the vision to be 20/200. Ophthalmoscopic examination of the lens showed a larger area free from opacification; there were fewer striae, and the fine globules were more evident. Twenty-six days after the beginning of treatment the vision was 20/100—1; ophthalmoscopic examination showed more of the fundus, the striae were finer and appeared more tortuous, and the opacifications showed a distinct sector-like arrangement. Apparently the vacuoles were less numerous than at the previous examination (later they gradually disappeared entirely). This objective appearance remained stationary, but the vision improved to 20/70, Jan. 9, 1919, and has so continued to date. While this cataract is not a purely senile one, the pathologic condition in this lens is probably identical with that found in senile cataract. If the lenticular opacity were due to the syphilitic infection, one would have expected some improvement from the antisyphilitic treatment; but despite such treatment the vision deteriorated until radium applications were instituted, after which the lenticular opacification diminished and the vision improved.

CASE 6.—Senile Cataract.—The patient's left eye had been under observation for the previous eleven months. Examination before radium was applied showed the vision to be 20/30. Lens examination with the ophthalmoscope showed mainly an isolated peripheral opacification. The fundus was normal. Radium was first applied, June 20, 1918, and has been continued uninterruptedly to date. The lenticular opacity and vision gradually improved. Nov. 12, 1918, ophthalmoscopic examination with the aperture gave a faint shadow of the opacification, which had formerly been very evident with this method. Detailed study of the lens indicated that the opacification was less dense than before, owing to the disappearance of some of the striae and of some of the finer granules which bordered the larger opacifications. The vision was 20/15, and the fundus remained unaltered. This case represents a diminution in the opacification of the lens. The vision, which was useful at the beginning of the application, is now practically normal, and has been in this condition for the past ten months. A point of especial interest in this case is that the right eye, which had a matured cataract when first observed, received several applications of radium during the first month. It was then extracted without any technical difficulties, and a very good functional result was obtained; yet the patient, who is a baker, claims that he could do his

work better before, when he had one blind eye and the other with a vision of 20/30 than he can today when the right eye is aphacic and the left has a vision of 20/15.

CASE 7.—Senile Cataract.—The patient has been under observation for the past ten months. The first examination was made, July 11, 1918. The vision at that time was 20/100. The lenticular examination showed three prominent radial sectors and peripheral curvilinear opacifications. Between these striae a moderate amount of opacification was apparent. The vision gradually improved under radium treatment, being 20/30, Sept. 9, 1918; but the opacifications, which had remained unaltered to this date, did not disappear until March 12, 1919, when a redder fundus reflex was distinctly apparent in this area. The important point in connection with this case is the fact that whereas the improvement in vision began a few weeks after the first application of radium, the diminution in the opacification did not appear until eight months after the beginning of the treatment, indicating that the action of radium may be continuous and cumulative in cataracts as well as in malignant tumors. In this case also the other eye, with a mature cataract, was treated before extraction. No technical difficulties were encountered during operation, and a good functional result was obtained.

CASE 15.—Senile Cataract.—The vision was 20/30; hypermetropia, 0.75; corrected to 20/20. Lenticular examination showed a broad peripheral curvilinear opacity. The fundus was normal. The patient complained mainly of floating opacities in the vitreous, which were muscae volitantes. The first radium application was given, Sept. 26, 1918. March 18, 1919, the vision was 20/15. The lenticular opacity was less dense and defined, while the floating opacities had entirely disappeared. After treatment it is common for these patients to observe that these muscae volitantes have disappeared.

CASE 17.—Senile Cataract.—The urine examination showed sugar, albumin and granular casts. The patient could count fingers at 6 feet. The lenticular opacity consisted of an extensive opacification of varying density. The fundus examination showed a blurred disk margin. Radium was applied, Dec. 27, 1918. Feb. 22, 1919, the vision was 20/70. The lenticular opacity became less dense in its central area and also on its nasal aspect. March 23, 1919, the vision was 20/50; the lens condition and the fundus were not altered. In view of the striking improvement in vision, and by analogy with Case 7, it is quite likely that in the next few months the opacification in this case will show further diminution.

CASE 18.—Senile Cataract.—Dec. 25, 1918, the patient's vision was 20/40+1. The lenticular opacity consisted of numerous small granules in the anterior cortex. The fundus showed a few grayish spots on the nasal side of the disk, of vascular origin. Feb. 20, 1919, the vision and fundus were unaltered, but the granules were distinctly less numerous. February 27, after repeated examinations, the small granules could not be seen. The vision and fundus were unchanged. The importance of this lies in the fact that the disappearance of the small granules after two months of treatment indicates that an early small cataractous formation may be cleared up by radium.

COMMENT

It is extremely difficult to obtain a correct estimate of changes occurring in the course of the development of a pathologic process which is so extremely chronic and protracted as a cataract usually is. Nevertheless, it appears to us that some of the phenomena observed are so striking and characteristic of the influence of the radium rays on the processes that it is impossible to conceive of them as mere coincidences.

Of the twenty-four cases studied, in 87.5 per cent. there has taken place some change for the better. As a rule, this improvement occurred in the first few weeks of treatment. After that, the condition usually remained stationary during the period of observation. But it must be understood that it was the improved condition which remained stationary, and in no

instance within the 87.5 per cent. did the lenticular opacification retrograde to the original condition as found before the initiation of the radium treatment.

In studying the changes which took place in the treated cataracts, three phenomena, depending in these patients on the function of the cataractous lens, were considered: namely, the vision, the appearance of the lens, and the visibility. By vision, was meant the amount of visual perception obtained by the patient through the cataractous lens; by the appearance of the lens, was indicated the amount and character of opacification found in the lens on an examination by the aid of transmitted light and by focal illumination; by visibility, was meant the ability of the ophthalmologist to perceive a more or less clear image of the details of the fundus through the cataractous lens.

A study of the results shows that in a certain number of cases both the vision and the amount of opacification were definitely improved; in a second group of cases the opacifications diminished, but there was no improvement in vision; in a third group, the vision was improved, while there was no change observed in the amount of opacification. In a few cases of this group an impression was gained that a brighter fundus reflex was obtained after treatment.

It is self-evident that the patients in the first group were improved. It is also easy to understand that an improved or diminished lenticular opacity may not be accompanied by any improvement in vision. The amount of vision which a cataract patient possesses depends more on the location of the opaque area within the lens than on the extent of the opacification. Striae, sectors, and large granules situated in the peripheral region may not impair the vision at all; while a smaller area of opacity situated in the center of the lens may very markedly affect the vision though not a great deal of the lens is involved. It is thus possible that certain striae or granules may diminish in size or completely disappear without in any way improving the vision. The greater objection may be raised that both the improvement in vision and the brighter fundus reflex are characteristics of too subjective a nature to be considered as clinical improvements without the supporting evidence of a change in the objective findings in the lens opacification; none the less, it must be granted that it is quite conceivable physiologically that both the vision may be improved and the fundus may appear brighter, and still no objective change in the picture of the lenticular opacification be noted.

If for the purpose of our discussion we regard a normal lens as analogous optically to a transparent fluid, then we may consider a diffuse lenticular opacification as analogous to an opaque emulsion consisting of minute granules evenly distributed in a transparent fluid. When the number of these granules is very great, or in other words the diffuse opacification is very dense, then no red fundus reflex can be obtained; when the number of granules is less or the diffuse opacification is not so dense, then we observe a faint reddish reflex from the fundus. Furthermore, when an objective finding has shown that such a red fundus reflex has become brighter, such a phenomenon can only have been caused by the disappearance of a certain number of the minute granules from the opaque lens. That these minute granules may disappear under the influence of radium is evident from the fact that in a certain number of our cases much larger granules disappeared under treatment.

This diminution of the quantity of the granules must be followed by both a brighter fundus reflex and some improvement in the vision of the patient.

We therefore maintain that while the best results were obtained in those cases in which there appeared an improvement in all the three phenomena described, we must also consider as improved clinically those cases in which there was direct improvement only in vision or in visibility. On the basis of such calculation there were in all twenty-one cases improved.

These twenty-one cases, or 87.5 per cent., represent such a high percentage of the cases treated that it is impossible to consider the result a mere coincidence. Furthermore, with the exception of one case, it was possible to maintain this diminution of the opacification as well as the improvement in vision and visibility during the period of observation, which was in every case not less than a few months, and in view of the fact that the majority of cataracts mature within a period of two years, most of the cataracts treated should have shown a certain amount of progression.

The time of observation is too short to determine whether this initial improvement can be maintained under treatment at stated intervals, or whether sooner or later the radiation will cease to be effective and the cataracts go on to maturation; but the results thus far obtained are very striking.

The biologic phenomenon that the course of development of a cataract may be, temporarily, at least, altered or inhibited by the action of the rays is fairly well established by this study. Furthermore, this investigation leaves no doubt as to the harmlessness of this agent. In order to establish the fact that any amount of the most intensive radiations leaves untouched the other structures of the eye outside of the lens, many of the patients were given longer and more frequent applications of radium than were actually needed. Moreover, in the beginning of the study larger quantities of radium were employed than are being used at present, and still there was no ill effect.

It is of importance to add here that five cases of matured cataract were also treated with large amounts of radium in order to ascertain whether a preoperative radiation would interfere in any way with the technic or the functional results of a subsequent operation. These patients were then operated on. No difficulties were encountered in the course of the operations, and the postoperative results were as favorable as the results in the nontreated eyes.

We therefore submit that, since there is no other agent in the therapeutic armamentarium which can influence the development of cataract in any degree at all; since surgery at its best leaves an aphacic eye which for practical purposes is frequently less satisfactory than the partial vision of a patient with an immature cataract; in view of the further fact that it is absolutely established that radium does no injury to the eye and does produce some, if ever so slight, biologic alteration in the lens, it is imperative that these clinical investigations of radium therapy in cataracts be continued by competent investigators, the subsequent results checked, and a definite conclusion reached.

SUMMARY

1. The application of radium to the eye is harmless.
2. There is diminution of lenticular opacifications under the influence of radium.

3. Should a cataractous lens become matured subsequent to radium treatment and should then an operation be required, no difficulties will present themselves.

4. It is therefore advisable to submit a sufficiently large number of immature cataracts with useful vision to a proper course of radium treatment so that a correct estimate of the value of the method may be obtained.

ABSTRACT OF DISCUSSION

DR. ISAAC LEVIN, New York: I am merely a radium therapist, and it is not for me to pass judgment on the clinical results. In fact, I will take it for granted, for the sake of argument, that we have thus far shown nothing, and that I stand today exactly where I did one year ago when Dr. Cohen asked me whether it is permissible or advisable to try radium in cataracts. When a new agency in therapy is applied to the human organism, the first requisite is "do not dare to do any injury." That must be borne in mind. I want you to understand that radium application, in the quantity, quality and dosage as used today in therapy of malignant diseases, does not injure any normal tissues of the organism with the exception of the epithelium of the skin, of the leukocytes, of the spermatozoa and the graafian follicles. All other tissues of the body resist the rays and are not injured by it. Nerve tissue, ganglion cells of the brain and the structures of the eye, are the most resistant tissues to the action of the rays. There is ample evidence in the literature as well as in my own work, that the largest amounts of radium employed in therapy do not destroy or injure normal tissues of the eye. I shall cite a case to illustrate this contention. I have a patient at the Montefiore Hospital who is suffering from sarcoma of the right upper jaw and the antrum. For the last year large doses of roentgen ray and radium were applied. The radium tubes were buried in the tumor, right near the orbit. The bone of the floor of the orbit is destroyed by the tumor so that the radium tube was lying for twenty hours at a time on several occasions right under the right eye. The roentgen rays were applied in the mouth also close to the eye. The eye remains absolutely normal now after one year of treatment. Thus intensive radiations correctly applied do not injure the eye. The next question is, how does the radium affect the lens? There is ample experimental evidence that it does not injure a normal lens. What may its influence be on a cataractous lens? We maintain that radium arrests the development of the cataract, and consequently has a beneficial influence on the condition. But even if further investigation should show that it has no such effect on a number of cases, it still does not injure the remainder of the organ. Therefore, the radium application in incipient cataracts should be employed for a number of years and thus a true estimate of the method obtained.

DR. ALLEN GREENWOOD, Boston: For the past fifteen years I have been following out a line of treatment which I have realized lessened my operative work in cataract cases to a marked degree. In my experience lenses showing a few spicules down and in are little affected and develop slowly; this is true also of nuclear opacities occurring in the highly myopic, and cases in which, besides the spicules, there are nebulous clouds or fine granular spots throughout the lenses, which are the vast majority of cases that interfere with vision. I give a hopeful prognosis and at once institute the following treatment: One per cent. solution of dionin (ethylmorphin hydrochlorid) is dropped into the affected eye three nights out of each week, and after a few months three nights out of every other week. It is well known that the continuous use of dionin will be of little value, but used as above, this drug, which is the one which most markedly affects the lymph circulation and hence the capillary circulation, exerts a decided influence on the nutrition of the lens. The often repeated effect of this improvement in circulation will frequently dissipate lenticular opacities and retard their future development.

DR. E. L. JONES, Cumberland, Md.: Some time ago I published a pamphlet on "The Usefulness of Dionin in Early Senile Cataract," and another one on the arrest of cataract in its early stages. I have used this treatment for more than ten years and I have a number of patients whose vision after a lapse of ten, eight, and six years, still continues to show improvement, and it was brought about through treatment of this character in the early stages of the disease. It is of little use in the later stages. In cases of reading vision, a few doses of dionin will clear up the dust formation or mist; but while the cataract will clear up the spicula remain. The cataract is like a thicket, with leaves on the twigs and branches in the summer time; and after the summer, the leaves fall off and the twigs and branches remain. That is comparable to the spicula. I use dionin with mercuric cyanid, and it works well in a majority of cases.

DR. JOHN E. WEEKS, New York: We all recognize that cataract is due to an interference with the nutrition of the crystalline lens. The epithelium of the anterior portion of the capsule is defective, as shown by Hess. There is the development of granules and minute hyaline globules forming liquor morgagni, and in addition there is erosion of the fibers of the lens to some extent where the gray spiculae are found. A disappearance of the cloudiness of the lens by absorption of the granules and the globules is sometimes observed. This is all we can hope to produce. We cannot cause the opacities due to erosion of fibers to disappear. In my opinion increase in the nutrition of the crystalline lens, either by improving the patient's general condition or by increasing the circulation in the anterior segment of the globe, may cause an arrest of development of cataract in some cases, a retardation in the development in some, and will be without effect in others. For the last thirty years I have been working on this hypothesis. I have used drops consisting of glycerin, boric acid and water. The glycerin produces a hyperemia of the anterior segment of the globe at every application, and may be used indefinitely without harm. I have discarded dionin, as it produces stimulation only over a short period of time. Dr. Greenwood has permitted an interval to elapse after using it three times, so that when used again it produces a further stimulation of the circulation. He has reached results similar to my own. It is immaterial what agent you employ provided it is harmless and sufficient stimulation is produced. I have observed the results of the use of glycerin in considerably more than 1,000 cases in private practice, records of which are in my office. The results have been satisfactory in perhaps a higher percentage of cases than that reported by Dr. Cohen as having been obtained in the treatment with radium.

DR. F. PARK LEWIS, Buffalo: There is an element of danger in allowing the impression to go out that this is a remedy for cataract and that it can be used without danger. I am not sure about that, for certainly it cannot change the metabolism of the lens and at the same time do no harm. Furthermore, I do not think we should employ a remedy empirically, which does not take into consideration the origin of the disturbance. The metabolic changes which have taken place in the lens are frequently the result of toxins in the system, from focal and other infections in the teeth or tonsils. Any remedy is unscientific which does not take into account the cause of the trouble which we are trying to control.

DR. S. D. RISLEY, Philadelphia: For many years I have been interested in the etiology of cataracts. I presented my first paper on the subject in 1889. The paper was based on eighty-six cases of beginning opacity of the lens, in each of which the record was sufficiently complete for analysis. During the intervening years, following a careful study of the eyes of the schoolchildren in Philadelphia made from 1878 to 1881, I recognized fully the importance of choroidal disease, since it was present in all of the eyes which had become myopic in the schools. Therefore, when any patient came into my office, after this fact had been grasped, no matter what the age of the patient, I treated the choroidal inflammation often by persistent and prolonged use of atropin and corrected any existing refraction error. It was very soon

observed that patients with incipient cataract, were greatly benefited by this treatment. Not only were they relieved from asthenopia, but the swelling and opalescence of the lens grew less and the sharpness of vision as shown by the test card often improved. While I have never seen spicules or banks of opacity disappear, in any case, I was convinced by these studies of the importance of choroidal disease over the nutrition of the avascular tissues: the lens, the vitreous body and the cornea. All cases of refraction error had therefore been treated by more or less prolonged use of mydriatics and careful correction of refractive defects and abnormalities of binocular balance. In eighty-six cases I found incipient cataract, as I have stated. The records were sufficiently complete for analysis, and there had been found refraction defects, choroidal disease and degenerating vitreous. To my surprise and gratification the opacity of the lens had not advanced after the treatment. The vision in many cases had improved; the asthenopia had disappeared, and the eye remained well after many intervening years. This experience was first published in 1889. Many hundreds of persons have been treated since, but added experience has but served to confirm the views then expressed. The essential fact in all this was the importance of uveal disease as a factor in the impairment of the nutrition of the avascular structures of the eye: the vitreous body, the crystalline lens and the cornea. In a large percentage, constituting the majority of the cases coming under study, the choroidal disease was associated not only with commencing opacity of the crystalline lens, but with a degenerating vitreous; dilated anterior ciliary vessels and uncomfortable eyes with failing vision. So frequently were the anterior ciliary vessels dilated, that I have come to regard this condition as an indication of uveal disorder. It soon became obvious also that the uveal disease was often but a local expression of some systemic disorder, as cardiovascular disease with kidney involvement. Indeed, I believe that the uveal tract of the eye is as liable as are the kidneys to participation in systemic dyscrasias, and therefore, that the treatment of these affections, is often quite as important as any local measures which may be adopted.

In addition to the prolonged use of the mydriatic, of which I have no dread, even in aged people, I have often used at the same time twice daily, weak solutions of physostigmin (eserin) salicylate, especially where the anterior ciliary vessels were engorged. This drug has been used because I believe it contracts the blood vessels and so stimulates the circulation which would be quite in accord with the suggestion made by Dr. Weeks.

DR. G. J. SAVAGE, Nashville, Tenn: Several things can be done in incipient cataract. I endorse practically everything they have said; but there are two things they have not said, and my experience with one of these for twenty-five years makes me a strong believer in its efficacy. One of the two suggestions comes from Dr. Woodruff, and that is, "Why may not the Crooke lenses do these patients some good?" The other thing I have been using for twenty-five years in every case of incipient cataract. It is a cell-building dose of mercuric chlorid and potassium iodid. Cataract will not develop when there is normal nutrition in the anterior segment of the eye. Anything that will improve cell-activity in this part will help the cataract condition in its inception. One one-hundredth grain of mercuric chlorid and 1 grain of potassium iodid after each meal for five weeks, and then one week of rest, and then five weeks of treatment and one week of rest, and so on, will check the cataractous changes which have just begun, in many instances.

DR. A. S. GREEN, San Francisco: Following my return from India five years ago, and since then, I have been using Smith's method for the treatment of incipient cataract. This consists of a subconjunctival injection of 15 minims of mercuric cyanid, 1:4,000. Intense swelling of the conjunctiva results which subsides in from four to ten days, but the redness does not disappear inside of a month. In addition to this treatment, however, we prescribe an eyebath consisting of a 1 per cent. solution of potassium iodid, every morning and every other night. On the alternating night the patient uses one drop of a 2 per cent. solution of dionin, which is

gradually increased to a saturated solution. This treatment is kept up for six months. Smith makes this point, that there is little use in giving this treatment when vision has fallen below 50 per cent. It is a treatment that can be given by any ophthalmologist, and the improvement in our hands has been as good as those reported with radium.

DR. EDWARD JACKSON, Denver: To go back to radium, perhaps the lack of interest in that phase of the subject has been due to the feeling that it was not available for most of us to try. I think it is important that any treatment that purports to influence a cataract should be investigated, and that the profession should arrive at some conclusion at as early a time as possible, as cataract cure without operation is all that is necessary to attract public attention to any proposed means. The possibility is before any one to use radium. It is not necessary to have a radium institute, or to be where large amounts of radium are accessible. The radium emanation has all the physical, electrical and other properties of radium salt, and will very soon be available anywhere in the country. No radium reaches maturity, that is, after thirty days it remains with very little change for a long time; but the radium emanation given off from large amounts of radium has the same properties; but it loses its property very rapidly. I think in four days it loses one half of its efficiency; and at the end of one month, it is practically inert. Since the close of the war it is possible to furnish the radium emanation anywhere. Of course, the physician can arrange for its use; on a certain day he can receive his supply and use it in a considerable number of cases. In that way it becomes practicable for a large number of us to employ and to experiment with this method. One half of the radium of the country is manufactured in Denver, and it was impossible for any of us to get any of it until the close of the war.

DR. S. LEWIS ZIEGLER, Philadelphia: I think that we should place on record a warning against the statement that radium cannot injure the eye when used in close contact with the cornea. Those who have seen a roentgen-ray burn of the cornea know that it is one of the most persistent and painful lesions. I have seen but one case, that of a surgeon who was wholly incapacitated by the intense pain. Relief could only be gained by enucleating the eye, although good vision was still present. The analogous action of roentgen ray and radium should make us pause, therefore, before recommending this agent to be applied by those who may prove to be careless and ignorant in its use.

DR. MARTIN COHEN: I have been under the impression, possibly wrong, that from reports dionin and mercuric cyanid and other products have not been as efficient as they may appear. If the products do what is claimed, then radium is only an aid. The twenty-four patients were examined consecutively, and after treatment of three, four and five weeks, they have responded with objective results which I have tried to show on the screen. If radium is effective in epithelial growths and degenerations then it seems the cataract problem might be attacked from that point. We know that radium does not injure the capsular epithelium in normal frogs' eyes. As regards affecting the cornea, we believe that if properly applied to the cornea radium will produce the results mentioned and cause no ill effects to the structures of the eye. We are interested in this problem from the scientific standpoint, and stated that radium is applied only in early cases of incipient cataract, and that some benefit was obtained by the absorption of the lenticular opacities.

Is Your Community Fit?—Is malarial fever a health problem in your community? Has a survey been made by experts to advise as to proper methods for control? The Anopheles mosquito, carrier of this disease, may be breeding in collections of water which need draining, oiling, stocking with fish, or other measures for mosquito control. Control malaria in your community and you may find you have less labor shortage, the physical well-being of your people may show a marked improvement, and your community may enjoy a great economic uplift.—*Pub. Health Rep.*, April 25, 1919.