a strangulated hernia may not always be free from danger, it can rarely be so serious as the condition which it is intended to relieve, and certainly does not deserve the unqualified disapproval and condemnation that a worthy examiner at the Royal College of Surgeons bestowed upon it when he told a candidate, who had suggested that in this very case aspiration should be tried, that if he did so he would certainly kill the patient.

Strangulation of irreducible hernia; evacuation of fluid contents of sac by aspiration; relief of strangulation.—D. S—, aged seventy-two years, admitted Nov. 10th, 1873, having had a hernia for six years, which was habitually reduced by the patient himself, though on two occasions he had to come to the hospital, where it was reduced by the surgeon. On November 9th the hernia again became unmanageable, and he came the next day to the hospital. The surgeon in the out-patient's room being unable to reduce it, he was admitted into the ward. At this time he was in no pain, had not vomited or passed wind per anum. In the right groin was an elastic swelling without impulse or gurgling. Ice was continuously applied.

The next day the patient was in an almost similar condition. A little griping pain and inclination to vomit, but no more.

Nov. 12th.—The patient being under chloroform, the taxis was applied, and once or twice gurgling elicited, but without reducing the mass. The aspirator then drew off about three ounces of viscid fluid, with no special odour. The swelling being now relaxed, the lower part was found to be soft and yielding, the upper part hard and firmly fixed in the inguinal canal. While attempts at reduction were being made a bubbling was felt in the upper part, and while the patient vomited an impulse was transmitted along the canal. The next day severe diarrhoea set in, which continued through the three following days, the tumour remaining in its relaxed state, sometimes allowing an impulse to be transmitted through its length, but not often.

The man did very well after this, though the fluid part of the swelling having been reproduced, Mr. Hill evacuated it a second time with the aspirator, but again failed to restore the protrusion to the abdomen. No irritation, pain, or symptom of any kind followed these procedures. While in hospital the patient was sent to the College of Surgeons for the examination of candidates for membership, a diversion he enjoyed much, being particularly tickled by overhearing an examiner rebuke a candidate who suggested tapping the tumour by saying, "Do you know, Sir, you would kill him if you did so." The patient had already undergone tapping twice without the smallest inconvenience.

On Nov. 27th the patient, wearing a well-fitted bag truss, was discharged in good health.

BRISTOL GENERAL HOSPITAL.

ACUTE SOFTENING OF SPINAL CORD.

(Under the care of Dr. CLARK.)

FOR the following notes we are indebted to Mr. J. Howell Thomas, physician's assistant.

Ann M—, aged eleven years, was admitted Dec. 31st, 1873, with paraplegia. About a week before admission the patient was seized with shivering and pains in her limbs, but she continued to go about until three days ago, when she suddenly lost the use of her legs. She never had any convulsions, nor was there any history of an injury to the back.

On admission the patient complained of pain in her back, along the vertebral column; the pain she stated to be worse in the cervical region. Could move her head about freely. The pain was not increased on pressure. There was complete paralysis of motion of the lower extremities. No reflex action. Sensation was diminished in the right lower extremity, and there was absolutely no sensation in the left. Complete loss of power over bladder and rectum. No abnormal physical signs in thoracic viscera. Urine acid; specific gravity 1018; no albumen; lithates. Optic discs normal.

Jan. 1st.—Temperature: evening 100.4°.

2nd.—Temperature: morning 99.4°, evening 100°. There is now no sensation in the right lower extremity. Bed-sores on the buttock.

3rd.—Temperature: morning 100°, evening 101°. There is partial paralysis of the upper extremities. Patient is unable to raise her arms as high as the shoulder. She is able to pronate and supinate the forearm, and was able to hold her knife in her hand to cut some bread this morning, but could not raise it to her mouth. Sensation diminished, but not lost.

4th.—Temperature: morning 99.4°, evening 99°. Vomited this morning. Says she feels worse. Her condition remains about the same.

5th.—Temperature: morning 98.4°, evening 98.2°. The pain in the cervical region is worse this morning; it is not increased by pressure.

6th.—Temperature: morning, 100.4°; evening, 98.4°. Urine alkaline; specific gravity 1026; ammoniacal; pus, albumen, &c. There is some cystitis. Patient does not complain of pain over the bladder, except on pressure.

7th.—Morning temperature 98.4°. The paralysis of the upper extremities is increasing; patient can just move her fingers, but is unable to raise her arms. — 1.30 P.M.: Difficulty of breathing came on; lungs were found to be congested.

Continued in much the same state, the difficulty of breathing increasing, until death at 5 P.M. on Jan. 10th.

Autopsy (Jan. 12th).—The brain weighed 2 lb. 8 oz.; the veins on its surface were congested; its substance was pale, and there were several minute points of injection in it. The spinal cord weighed $\frac{3}{4}$ oz.; the membranes were healthy; there was a quantity of serous fluid in the arachnoid sac; the cervical portion of the cord was soft and disintegrated; the rest of the cord was firm, but on making a transverse section there were several points of injection in it. Lungs congested; pleuritic adhesions on the right side. The heart weighed 4 $\frac{1}{2}$ oz., and was healthy. Liver, spleen, and kidneys normal, but intensely congested. Stomach, intestines, uterus, and ovaries normal. The bladder was in a state of acute suppurative inflammation; the coat was very much thickened, and the mucous membrane lining the interior was covered with a yellowish-grey slough.

HÔPITAL ST. ANTOINE, PARIS.

CASE OF RHEUMATISM OF THE TENDONS.

(Under the care of Dr. M. PETER.)

THE patient, aged thirty-five, had been suffering for a few days from pain in the situation of the joints of the knees and ankles. No swelling or redness was to be observed in the painful situations; no general symptoms of rheumatism; no fever. The pain was developed when the finger was moved carefully along the head of the fibula, on the tibia in the situation of the aponeurotic expansion of the tendons, and behind the malleoli. Movements of the articulations were painful through the straining exerted on the inflamed tendons.

M. Peter has often observed such cases of rheumatic inflammation of the tendons and serous bursæ, unconnected with gonorrhœa. The pain is to be sought after, not with the hand, but with the finger, carefully exerting pressure on the situation of the tendons and bursæ—for example, behind the ankles, between the biceps and head of the fibula, above the aponeurotic expansion of the knee, behind the calcaneum and the olecranon, in front of the patella, in the situation of the great trochanter, &c. The disease is really an extenuated form of rheumatism, and constitutes rheumatic synovitis. It rarely presents cardiac complications. It is seldom accompanied by any great fever, though sometimes the fever may be somewhat high, and the evening temperature may rise to 39° or 40° Cent. There may be slight œdema or not in the situation of the inflamed tendons. Likewise the surface may be discoloured or not. It is much milder than articular rheumatism, and does not require any energetic treatment.

A CORRESPONDENT of the "Cork Constitution" states that in the south of Ireland a fatal disease is destroying the poultry, which are dying off by hundreds without any apparent cause. It would be interesting to get the results of a careful post-mortem examination of one of the birds.