entirely supports the validity of Darwin's theory, in the locality selected as a test.

EDWARD B. POULTON.

Oxford, February 20.

The Accuracy of the Principal Triangulation of the United Kingdom.

THERE was some discussion of this question at the last meeting of the British Association, and an investigation by Capt. H. St. G. L. Winterbotham has been published by the Ordnance Survey (Professional Papers, new series, No. 2). It appears, however, that the measurement of Lossiemouth Base, valuable as it was, did not definitely decide the question, but that a moderate amount of further computation would do so.

(1) It will be generally admitted that statistical evidence as to the precision of any kind of observation is of no great value unless it is based on a large population. For example, if we may assume the principal triangulation to have been all executed by similar observers under similar conditions, and no unjustifiable rejections to have been made, we may accept with great confidence the probable error of $\pm 1.23''$ computed from the closure of 552 triangles. But nowadays one would not think of estimating the precision of a base-line by the discrepancy between two measurements of it, though one might reject the measurements if they did not agree as well as was to be expected from the known usual probable error of a measurement. Thus discussion of a large population of discrepancies gives a good estimate of the probable accidental error of a measurement, but a small population only fixes a lower limit to that probable error. Now we have only a population of three independent discrepancies between the four bases of Lough Foyle, Salisbury Plain, Lossiemouth, and Paris, and even if it is brought up to six by the inclusion of the three bases measured with steel chains, yet it must be considered a very small population upon which to base any estimate of precision.

(2) But the original question, to what extent the strength of the figure compensates for the large probable error of an angle, is not necessarily a question for experiment, but it is essentially a question for computation. Given the probable error of an angle in a block of triangulation adjusted rigorously by least squares, if it is required to find the probable error of the distance from any point of it to any other point, the first step is to express the unknown error in that length as a linear function of the unknown errors in a number of independent angles (*i.e.* two angles of each triangle in a chain of independent triangles stretching from one point to the other, and in another chain stretching from one of the points to the nearest base). It then remains only to add another column to the least square computation by the method described in Wright and Hayford's "Adjustment of Observations," paragraph 123. May I venture to suggest that if two or three such cases could be worked out for strong figures in the United Kingdom, and for comparison two or three cases for weaker figures, either there or elsewhere, it would not only set at rest the immediate question, but would also establish results of great importance for surveyors in general.

T. L. BENNETT.

Computation Office, Egyptian Survey Department.

ONE can but agree with Mr. Bennett in insisting on a large "population" of discrepancies upon which to found a calculation of a probable error, whether it be of the measurement of a base-line or of an angle.

His example of a base-line measurement is not, how-

ever, strictly comparable with the investigation in question. In the former a large number of independent measurements must be made from which to deduce the most probable length and the individual discrepancies from this length. In the latter the measured bases may be regarded as errorless compared with the triangulation, and the actual errors can therefore be deduced with surety.

Neglecting the steel chain bases, we have four independent measures upon any one of which the triangulation can be made to depend. These four are widely distributed.

The longest line from base to base is that from the new base at Lossiemouth to the Paris base, and along this line the old steel chain bases add additional proof that no serious errors are inherent in the triangulation.

I agree that, "if we may assume the principal triangulation to have been all executed by similar observers under similar conditions and no unjustifiable rejections to have been made, we may accept with confidence the probable error of $1.23''$ computed from the closure of 552 triangles," but it must be remembered that this is the probable error of an observed angle. The date of the work makes this a matter of no surprise. The point at issue, however, is not the probable error of an observed angle, but the probable error of an adjusted angle, or, in other words, to find out how far the intricacy of the figure has compensated for the lack of precision of angular measurement.

To say that this question is essentially one for computation is not, to my mind, correct.

The probable error of the ratio of any two sides as derived from the triangulation depends upon the probable error of an adjusted angle. This calculation is possible, but from the complexity of the figure and the intricate system of weighting the angles, impracticable. Moreover, the answer to it would still be the probable, and not the actual, error.

It is, however, possible to pick out of the general figure chains of simple triangles connecting the bases. Supposing that these chains had been the only paths of calculation, and that they had shown the same errors of ratio between the bases as are actually found, we can deduce what the probable error of an observed angle would have been to have effected this result. Such an investigation has been made, and the probable error of an observed angle in these "equivalent" simple chains is $0.85''$, or approximately the same as that given by General Ferrero, in his 1892 report, as the mean figure for the probable error of an observed angle in the triangulations of those twenty countries represented on the International Geodetic Association.

Although, therefore, further investigation on the lines advocated by Mr. Bennett would be one of the greatest interest, I do not think that it promises a result commensurate with the time and expense it would entail.

H. S. L. WINTERBOTHAM.

Atomic Models

I AM indebted to Mr. Chalmers for pointing out (NATURE, February 19, p. 687), what I had indeed suspected, that the magnetic moment due to an electron moving in a circular orbit, assuming the angular momentum to be $h/2\pi$, is *exactly* five times the magnetic moment of the magneton. The original value (15.94×10^{-22}) of the latter quantity, given by Weiss, and quoted in my former letter, was based on the value of Avogadro's constant found by Perrin. If we divide the magnetic moment of the atom gram, 1123.5, by the more recent value for Avogadro's constant given by Millikan (6.062×10^{22}) we obtain as the magnetic moment of the magneton 18.54×10^{-22} , which is exactly

one-fifth of the value of the magnetic moment, 92.7×10^{-22} , for the electron moving in a circular orbit.

Sir Oliver Lodge has directed attention to the importance of all cases where commensurable numbers enter into physical problems. Mr. Chalmers thinks that the present result indicates that, in magnetic materials, there is a unit of five (or ten) atoms, which has a constant number of magnetons. Since in the last resort we must consider magnetism to be an atomic property, I should prefer to regard the result as affording further evidence for the view that the magnetic effects of the complex nucleus must be taken into consideration. The magneton may then result as a difference effect.

To make this clearer, it may be worth while discussing a simple illustrative model. Prof. Peddie has put forward some interesting suggestions as to the structure of the atom in the February number of the *Philosophical Magazine*. He supposes that the atom may be built up of concentric spherical shells of electrification, which may be in rotation round a common axis. Following this suggestion, suppose we have a uniform sphere of positive electrification of radius A rotating with angular velocity Ω . Outside this, suppose we have a single ring containing n (from 1 to 8) valency electrons. The remaining negative electrification may be relegated to a central core having no rotation. Then the magnetic moment of the rotating sphere may be taken as $\frac{1}{2}EA^2\Omega$, where E is the total positive charge, which we shall assume equal to Ne .

We have no direct evidence as to the value of $A^2\Omega$, but if for convenience we assume that it has the same value as $a^2\omega$ for a ring electron, we obtain the result that the magnetic moment of the rotating core is equivalent to $2N$ magnetons. The resultant magnetic moment for such a model would be the difference between the $2N$ magnetons of the core and the $5n$ magnetons of the ring. This is only intended to illustrate the way in which the magneton may be introduced as a unit for measuring magnetic moments without involving the necessity of a single magneton existing as an independent entity. If the core of an atom is built up of α and β particles in orbital motion, the experimental results of Weiss indicate that the resultant magnetic moment of these particles can be expressed in terms of the magneton.

H. S. ALLEN.

Wheatstone Laboratory, King's College,
London, W.C.

Origin of Structures on the Moon's Surface.

IN NATURE of February 5 Dr. Johnston-Lavis writes, "the more I compare the moon's surface with volcanic vents, in different parts of this world, the less I see a resemblance between the two," and "the more does the planetoid and meteorite projectile theory become acceptable."

G. K. Gilbert, in his address to the Philosophical Society of Washington, 1892, gives an extremely interesting summary of the various theories to account for the features of the moon's face. After a very clear description of the phenomena to be accounted for, he accepts the meteorite theory with modifications. He remarks that, if the so-called craters of the moon were due to the impact of meteors, their form would be for the most part elliptical, whereas, in fact, they are circular. His own theory is that the earth was at one time attended by a ring similar to that which encircles the planet Saturn, and that this afterwards "gradually coalesced, gathering first around a large number of nuclei, and finally all uniting in a single sphere," the moon. He attempts to show that this hypothesis accounts for the facts.

NO. 2313, VOL. 92]

In NATURE (vol. xxv., p. 243, 1862) I suggested that when, according to Sir G. H. Darwin's view, the moon broke away from the earth and commenced an independent existence, the scar left by the great catastrophe forms now the basin of the Pacific Ocean. The same idea was elaborated five years later by Prof. Pickering (*Journal of Geol.*, vol. xv., No. 1, 1907). It is evident that if the Pacific Ocean indicates the place from which the moon departed, the continents surrounding it must have been then in existence and the earth covered with a solid crust. Both Prof. Pickering and myself make this assumption. The question which I would ask, therefore, is this: Does the moon's surface bear traces of this mode of origin?

The material detached from the earth would have been partly solid, derived from the cooled crust, and partly liquid, derived from the molten substratum. The expulsion would have been probably explosive, owing to the gases dissolved in the substratum. The material would consequently have been scattered. Subsequently it would have collected about its centre of gravity, the smaller masses falling in last. The paths of the falling masses would have been radial. What the telescope now reveals would be the final effect after the mass had settled into the spherical form. May not, then, the circular so-called craters have been caused by the impact of fragments of the solid crust, and may not the mountains of the moon be also angular portions of the earth's crust projecting above the mean spherical surface of the moon?

The above suggestions do not exclude the possibility of true meteors also having fallen, and left their mark upon the moon, but it is not probable that many large meteors have struck the moon, because they are rare upon the earth, and if many *siderites* had so fallen, the moon's specific gravity would have been higher than it is, viz., about 3.

It has occurred to me to inquire at what distance from the earth's surface the centrifugal force would balance the earth's attraction, because it is difficult to see how the material, which is supposed to have been detached from the earth to make the moon, could have got away until that distance was reached.

This condition is expressed by the equation, $a\omega^2 = g \frac{a^2}{r^2}$, where a is the earth's radius, 20,926,202 ft.,

ω the angular velocity, assumed in this case to be $2\pi/18,000$ seconds, g gravity, 32 ft. a second, and r the distance from the centre required. From this I get the required distance from the earth's surface, viz., $r-a$, equal to 53,205,200 ft., or more than double the earth's radius. How this distance could be reached seems unaccountable.

Sir G. H. Darwin, in the summary of his paper on the remote history of the earth, does not refer to this point; but since his theory has been adopted without demur by Sir Robert Ball and Prof. Pickering, the difficulty which strikes me must be more apparent than real. Can some reader of NATURE explain it?

O. FISHER.

Graveley, Huntingdon.

I HAVE read with a great amount of interest the letter by Dr. Johnston-Lavis which appeared in the issue of NATURE for February 5, as regards the ray systems of the moon from a volcanologist's point of view. Even taking into consideration his explanations, however, I cannot now help comparing the moon with our own planet, especially when Laplace's nebular hypothesis, and subsequent theories in reference to the relation of the moon to the earth are considered, which seem to prove conclusively that the moon was once a portion of this world. Then with