ress to the stage where it might easily become so.

On the morning of the third day the temperature was 101.8°, pulse 112, and respiration 36. The cough still was present in severe paroxysms, during which he would become very dyspneic and cyanotic. The consolidation had spread until then it included practically the whole of the left lower lobe; the pleuritic pain was rather severe and the pulse had lost considerably in tone.

After consultation with Drs. Ball and McInnes we determined to try plain human serum, and in order to save time I obtained a specimen of his blood. Dr. F. B. Johnson tested my own blood against his, and finding them compatible, withdrew about 75 c. c. of blood, removing the serum by the centrifuge method, and obtaining about 30 c. c. of serum, which was given that evening about 5:30. In about an hour he had a rather severe chill, rise of temperature to 104°, associated with a severe coughing spell, became very nervous and cyanotic (probably from the exertion of coughing) and the pulse became very rapid. This was followed in an hour by a drenching sweat and an improvement in all symptoms. The temperature had reached normal by 11 P. M., pulse 102, and respirations 24.

I wish also to emphasize the improvement in the patient's subjective symptoms associated with the fall in temperature. I can express it no better than to quote his own words to his wife about 9 P. M. as follows: "It is hard to believe that I feel as well as I do." He had no pain, could breathe comfortably and demanded that he be taken off the back rest which he had been on constantly since the onset, and asked to be let alone and allowed to sleep, which he promptly did and slept very quietly until 4 A. M., when he was awakened by a coughing spell and had to be placed on the back rest and given codeine for the pain which returned with the cough. The temperature rose to over 102° by 9 o'clock the next morning and preparations were made for repeating the serum that afternoon. By evening, however, the temperature had reached 100°, pulse 102 and respiration 32, and there was such an improvement both subjectively and objectively that it was decided not to repeat the serum that day and by 8 o'clock the next morning the temperature was 92.2°, pulse 84, and respiration 24. They never rose above normal afterward and further convalescence was uneventful.

In the case cited above, the fact that the patient had had influenza and still had the cough he developed with that disease, it is possible at least that it may have been associated with the Type IV infection which he had, although clinically the case was one of pure lobar pneumonia.

It is not expected that conclusions regarding the value of the procedure could be drawn from the results obtained in one case. At the Naval Hospital this method was adopted in the treatment of at least twenty-five cases in mine and other serv-

ices with very encouraging results, certainly to the extent of justifying its further trial.

BRONCHO-PNEUMONIA IN ADULTS*

By L. J. LINDSEY, M.D.,
Covington, Tenn.

Text books in the past have devoted very little space to this important subject. They merely refer to this disease as being a condition of early childhood, some going so far as to say that the majority of infants under three years of age all have this type of pneumonia, while those above this age have the more common form of lobar pneumonia, and end their statement by saying that it is occasionally found in adults.

In this paper I shall not attempt to refer to the types of bronchial pneumonia, due to the inhalation of foreign substances, nor that following anesthesia, poisonous gases, etc. This paper is based upon my experience and observation in the treatment of over 600 cases of pneumonia, 50% of these, or 300 cases, being of the bronchial type. This experience and observation was made in three United States Army base hospitals in France. Perhaps the best way to present the subject will be by briefly mentioning the definition, etiology, symptoms, diagnosis, prognosis, prophylaxis, complications and treatment.

DEFINITION

Broncho-pneumonia is a bacterial infection of the finer bronchi and their relative lobules. The process begins with the inflammation of the bronchioles and smaller bronchi. A capillary bronchitis, which extends to the alveoli, and the whole lobule or a group of lobules becomes filled with exudate, cellular and hemorrhagic, and is distinctly less fibrinous than in the lobar pneumonia.

Personally, I think in many cases the process of inflammation begins in the pharynx extending down the trachea into the larger bronchi and finally into the smaller bronchi and bronchioles. This statement is based upon the findings in

*Read before the West Tennessee Medical Association, Dyersburg, Tenn., May 29, 1919.

120 cases at autopsy, at which time there was a developed case of pharyngitis, tracheitis and bronchitis present, all of which showed a different appearance both macroscopically and microscopically, showing different degrees of inflammation along the course from the pharynx to the bronchioles. The same organisms were isolated from the different parts of this inflammatory process.

ETIOLOGY

Broncho-pneumonia occurs either as a primary or secondary affection. The secondary frequently follows measles, influenza, whooping cough and typhoid fever, and occasionally almost any of the eruptive fevers.

No doubt that the lack of fresh air, the inhalation of irritating substances, as dust, etc., and being subjected to exposure, especially when fatigued, are all predisposing causes.

As to micro-organisms producing this disease, the following stand out very prominently, namely: the micrococcus lanceolatus, streptococcus pyogenes, the streptococcus which may be either of the hemolytic or the non-hemolytic variety, the staphylococcus aureus and albus, Friedlander's bacillus pneumonia, the influenza bacillus, and following diphtheria the Klebs-Loeffler bacillus.

Let me state here that there is a tendency for a combination of the above-named organisms instead of a single variety. For instance, the streptococcus is frequently associated with the pneumococcus and these are frequently accompanied by the staphylococcus.

The reason for mentioning this combination is to show later under the treatment of this disease how difficult it is to obtain a proper serum or vaccine.

SYMPTOMS

These vary according to the type of infection and virulence. I shall attempt to picture the symptoms in a mild and later the symptoms of a severe case. It is the mild and severe cases that are hardest to diagnose, especially early, and of course you will understand that the average comes between the two.

The initial symptoms of a mild case are as follows: The patient first complains of a chilly sensation, no distinct chill as a

rule as you find in lobar pneumonia, accompanied by a headache and frequently a general aching over the entire body, slight sore throat and usually slight cough with little or no expectoration. Upon physical examination in this type little, if anything, is found abnormal in the chest. As a rule inspection does not reveal anything, neither does palpation nor percussion, but by careful auscultation you may be able to elicit a few medium moist rales. The favorite site of these rales is just below the angle of the right scapula, frequently accompanied, but not necessarily, by the same signs on the opposite side. In from twelve to twenty-four hours, forty-eight in some cases, these signs increase and the breathing is apt to show prolonged expiration, broncho-vesicular or bronchial breathing over these areas. It is at this time that you get a slight dullness on percussion. I might state that the percussion note is apt to be misleading, being more or less tympanitic in character, due to the compensatory emphysema of the surrounding tissue about the small area of consolidation. Of course, as the small areas of consolidation coalesce, the typical signs of consolidation make their appearance. Unless the area of consolidation is of considerable size the respirations are but little increased, frequently not running over 28 per minute, the pulse about 90 to 100, the temperature from 99 to 102°, sometimes reminding one of the intermittent type of malarial fever. There is a little cyanosis with a moderate amount of expectoration which often shows a moderate amount of blood. The expectorated material does not have the characteristic prune juice appearance, nor the gummy, tenacious sputum that one finds in lobar pneumonia, the leucocyte count ranging from eighteen to twenty-two thousand.

Now, in contrast to these mild symptoms, let me describe a severe case.

The patient is seized with a chill or rigor, severe headache, vomiting and general feeling of weakness, a moderate amount of cough and expectoration, with pain in the chest, often pain in the right iliac region, resembling appendicitis and hard to differentiate for the first twelve or twenty-four hours, also often shows signs of meningitis to the extent that a

lumbar puncture is often resorted to in order to clear up the diagnosis. Upon examination the physical signs in the chest are practically the same as in the mild case just described, except that the areas of consolidation form much faster. Cyanosis appears early, the temperature may go as high as 104° F., the pulse around 120, respiration around 30. The respiration and pulse rarely go higher until a few hours before death. These cases frequently die from within 24 to 48 hours after the onset. The leucocyte count, as a rule, is high, but owing to the extreme toxemia may be low.

DIAGNOSIS

Signs of an acute bronchitis with a temperature above 101° lasting over three days spells B. pneumonia, for the patient is too sick for an ordinary bronchitis.

With lobar pneumonia it may readily be confounded if the areas of consolidation are large and merged together. This, in my opinion, accounts for so few cases of this disease being recognized in the past. Usually the history of the two diseases is quite different, but if careful attention is given to the mode of onset, by careful physical examination one may be able to differentiate between the two, remembering the fact that broncho-pneumonia in the far majority of cases is bi-lateral and if seen early have the typical signs of acute bronchitis. The tendency in lobar pneumonia is for the lung to become affected by lobes and not by lobules. Later, as a rule, lobar pneumonia ends by crisis within three to eleven days where bronchial pneumonia frequently ends by lysis, three to fourteen days, some of our cases extending over a period of many weeks.

The next condition to be differentiated from bronchial pneumonia is an acute miliary tuberculosis and this differentiation, unless verified by the presence of tubercle bacilli in the sputum, may be impossible for at least two weeks, the physical signs being identical.

PROGNOSIS

The mortality of broncho-pneumonia varies greatly in different epidemics ranging from 10 to 60%. The reason for this, I am sure, is due to the increased virulence of the various organisms. For instance,

we know that the passage of micro-organisms from one individual to another increases the virulence of the type, and hence the epidemics of higher mortality through transfer through one or more persons, thereby increasing their virulence to the point where it is almost impossible for one to withstand it. Of course the prognosis depends largely upon the early diagnosis and proper treatment.

PROPHYLAXIS

I am convinced that by careful isolation of all infectious diseases with attention to the mouth, nose and throat, spraying or gargling with any good antiseptic solution, such as Dobell's, will serve as a prophylaxis in this disease.

COMPLICATIONS

Of the complications we had the following:

Empyema 10%, which may occur early and have a tendency to occur in pockets, making the diagnosis rather hard and impossible unless one resort to the frequent use of the exploratory needle, together with the use of the x-ray. Early operation for empyema is attended by a higher mortality than later on, after the fluid has become creamy, with a drop in temperature, pulse and leucocyte count.

Mastoiditis appeared in over 75% of the cases at autopsy, and, strange as it may seem, a large per cent. of these never complained of any pain during their illness.

Otitis media occurred in a few cases.

Fibrinous pericarditis occurred in three cases.

Peritonitis appeared in one case.

Erysipelas appeared in one case.

Hyalin degeneration of the muscle fibers of the rectus abdominis muscles leading to their rupture was found in two cases.

Bronchiectasis occurred in many cases. A sclerosis of the consolidated areas of the lung took place in 2%, there being an overgrowth of connective tissue replacing the normal lung tissue. These areas persist for months and years and are very hard to differentiate from a tuberculous process.

TREATMENT

The frequency and seriousness of broncho-pneumonia render it a disease which

taxes to the utmost the resources of the practitioner. There is no acute pulmonary affection over which he at times so greatly despairs. On the other hand, there is not one in which he will be more gratified in saving patients who have seemed hopeless. The general measures are much the same as in lobar pneumonia.

In the beginning of this disease, as in the beginning of all infections, a dose of calomel is generally indicated, the size of the dose being from three to six grains given in broken doses, followed in about three hours with magnesium sulphate. The patient should receive plenty of fresh air and, if possible, sunshine, care being taken not to chill the body. Water should be given freely and the temperature controlled by frequent sponging. Coal tar preparations should never be given except for a dose or two at the onset, for they have a tendency to cripple the heart, the one organ that suffers most in this dreaded disease. If the pain in the chest is too severe this should be controlled by the application either of heat or cold over the area, and if this will not suffice, by a small amount of codeine or morphin given hypodermatically; but care should be taken that the opiate does not constipate or lessen the secretions. As for expectorants, ammonium chlorid, combined with syrup of citric acid, has been found to be the most reliable. Rarely, if ever, does this produce nausea when given in this combination. If the cough is loose but rather persistent, a small amount of codeine or heroin may be added to this mixture. In no disease is it so necessary that care should be taken in selecting a suitable diet so as to maintain the patient's strength and at the same time produce as little tympany as possible. Carbohydrates should be pushed, also milk, eggs, broths and fruit juices given.

No patient with either form of pneumonia should be allowed to pass a day without at least one evacuation of the bowels, small doses of calomel given at intervals of every other day with an enema, or small dose of magnesium sulphate between times has, under my observation, given the best results.

Stimulants, of which strychnine has proven the most efficient, should certainly

not be given until the heart's action begins to show some indication of its weakness. Then it should be administered at about three-hour intervals, the dose being about a 30th to 20th of a grain, and this not given over too long a period. Digitalis has also been proven to be of great value in certain cases.

A hot water bottle should be placed to the feet of all pneumonia patients. This causes a relaxation of the blood vessels of the lower extremities and thereby relieves to some extent the already engorged lung.

Special care and attention should be given to the mouth, nose and throat that they are kept clean at all times.

During convalescence the patient's heart should be looked after, nutritious food given, and the patient not allowed to go to work for several weeks.

The serum and vaccine treatments in this disease have not proven satisfactory, due to the fact that they are worthless and often-times harmful when given before the definite type has been isolated and determined, and then as mentioned at the outset, this disease is more commonly due to a combination rather than a single germ. Therefore, it is almost impossible to obtain a stock vaccine that will be of any value, since little advantage has been gained by the use of autogenous vaccines, and this is certainly the most rational treatment of the two. Of the pneumonia serums, the Type I has been the most satisfactory, but it is to be remembered that few cases of Type I pneumonia will die without the serum.

I shall not mention the treatment of the various conditions that may arise, but in conclusion will say that a restful, quiet room, a sensible, efficient nurse, a sufficient amount of fresh air and a suitable diet and proper care of the bowels will prevent high temperature, heart failure, low blood pressure, insomnia, tympanitis and toxemia in very many cases, and prevention is far better than the treatment of these serious conditions.

CONCLUSIONS

In closing I wish to stress:

1. That broncho-pneumonia is far more common in adults than text books would lead one to believe;

2. The importance of an early diagnosis;
3. The importance of early recognition of complications, especially pus pockets;
4. The free use of the exploratory needle in determining the presence of pus pockets; and
5. A warning against the indiscriminate use of vaccines and serums in the treatment of this disease.

THE CONSERVATION OF MENTAL ENERGY

By FREDERIC J. FARNELL, M.D.,
Psychiatrist to Public Schools,
Providence, R. I.

In bringing this subject before you for your consideration, it will be necessary to precede my general remarks with the statement of a few facts in regard to the so-called laws of Nature and an attempt will be made to divert your attention for a short while toward another picture of the now much-discussed topic, the value of heredity and its allied subjects. You are all aware how much emphasis and stress is placed upon the fact that all or everything that favors the well-being of our nervous system goes toward the strongest support of our general health. What is this all, this everything? Heretofore, in the study of the processes of evolution it has been assumed quite generally that those laws which govern the animal world also govern mankind, even though it is a well-known fact that "man" holds a most unique position in this world. As a result of these studies, interests have fallen into the numerous expressions now evinced by the school of eugenics. This school bases the laws of inheritance upon the study of the lower animals and plants and then applies them to mankind.

This widespread interest has made people intellectually honest in their anxiety to know how far their physical and mental conditions can influence the physical and mental states of their children, for it is a well-known fact that the average man today has a definite opinion and belief. He firmly believes that any victory, which is really luck nowadays, over his morality, gives strength to his children, and any

lapse is detrimental and weakening to his offspring. He may often go further and feel that his athletic training and his high scholastic ability will give his son or his daughter a most superior muscular strength on the one hand or a much easier acquisition of intellectual knowledge on the other.

It is this misconception, to make it emphatic, that has led me to discuss the value of mental energy and demonstrate hastily that human evolution is in relation to plant and animal evolution only up to a certain point and that from then on it is entirely under the influence of laws distinctly its own. This teaching of eugenics has led to most unfortunate results. First, that as long as it is possible to believe that the inheritance which we transmit to our offspring might be modified by our own actions, there is hope. Secondly, however, if we are forced to believe that by nothing that we can do can we influence the inheritance which we hand to our children, there is utter hopelessness. By this view we are told that the inheritance is fixed definitely when we select our wives and husbands and that nothing that we may do subsequently can change the transmission. The choice of a husband or a wife is bound up in very complex social conditions in such a way that it is determined by an indefinite number of artificial factors of which physical fitness hardly plays even a minor part. Organic inheritance can only be controlled by controlling marriages, I grant you that, but marriages will, however, continue to be determined by passion and accident rather than by fitness.

Let me refer briefly to the relation of disease to energy. So often one hears the remark, "He inherits the disease from his father," or "His father was subject to the same trouble." Nothing is more certain than the ravages certain diseases may bring in a future generation, and yet it must be kept clearly in mind that these diseases are not inherited in the same sense as the color of one's eyes or the shape of one's nose. True, they are passed from mother to child; but these diseases are communicable in type, and the child does not inherit them—it "catches it" in just the same way as one might catch diph-