This article was downloaded by: [McGill University Library]
On: 17 October 2014, At: 11:15
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 5

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah11>

XXXII.—The zoology of Barents Sea

W.S.M. D'Urban F.L.S.^a

^a Devon and Exeter Albert Memorial
Museum

Published online: 07 Oct 2009.

To cite this article: W.S.M. D'Urban F.L.S. (1880) XXXII.—The zoology of
Barents Sea , Annals and Magazine of Natural History: Series 5, 6:34,
253-277, DOI: [10.1080/00222938009458937](https://doi.org/10.1080/00222938009458937)

To link to this article: <http://dx.doi.org/10.1080/00222938009458937>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of
all the information (the "Content") contained in the publications
on our platform. However, Taylor & Francis, our agents, and our
licensors make no representations or warranties whatsoever as
to the accuracy, completeness, or suitability for any purpose of
the Content. Any opinions and views expressed in this publication
are the opinions and views of the authors, and are not the views
of or endorsed by Taylor & Francis. The accuracy of the Content
should not be relied upon and should be independently verified
with primary sources of information. Taylor and Francis shall not be
liable for any losses, actions, claims, proceedings, demands, costs,
expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

THE ANNALS

AND

MAGAZINE OF NATURAL HISTORY.

[FIFTH SERIES.]

No. 34. OCTOBER 1880.

XXXII.—*The Zoology of Barents Sea.* By W. S. M. D'URBAN, F.L.S., Curator of the Devon and Exeter Albert Memorial Museum.

THROUGH the spirited exertions of the Dutch geographers a small vessel of 79 tons, named the 'Willem Barents,' was fitted out at Amsterdam by subscription, and made two voyages to Barents Sea, between Spitzbergen and Novaya Zemlya, in the summers of 1878 and 1879*. The objects of these expeditions were to examine the state and position of the ice in Barents Sea and, if possible, in the Kara Sea, also to take deep-sea soundings with serial temperatures, and thus supply important contributions to our knowledge of that interesting region. Magnetic and meteorological observations were also to be taken, and natural-history collections to be diligently made at every opportunity. Mr. W. J. A. Grant, a gentleman well known for his skill as a photographer, accompanied both these expeditions, and each time brought back with him some bottles filled with animals dredged in Barents Sea, and preserved in spirits. These he very generously presented to the Museum under my charge. The bottles were carefully labelled with the dates, latitudes, longitudes, and depths at which the specimens were obtained; and

* Accounts of these voyages will be found in the 'Proceedings of the Royal Geographical Society' for January and November 1879. See also the 'Illustrated London News' for January 25, 1879, and January 31, 1880.

as the collections appeared to include many very interesting species, I was anxious to submit them to the best authorities in the various departments. I was so fortunate as to find several eminent scientific gentlemen who were willing to examine the specimens for me. I sent the Crustacea collected in 1878 to the Rev. A. M. Norman, and those obtained in 1879 to Prof. J. O. Westwood; the Mollusca were determined by Dr. Gwyn Jeffreys; the Echinodermata of 1878 were examined by the Rev. A. M. Norman, and those of 1879 by Mr. W. Percy Sladen; Dr. McIntosh named the Annelids; and the Rev. Thomas Hincks has made a laborious examination of the Hydrozoa and Polyzoa. The few Spongia and Actinozoa were examined by Mr. H. J. Carter. I take this opportunity of offering these gentlemen my sincere thanks for their kindness, which has enabled me to draw up a reliable list of the animals brought home by Mr. Grant. As yet nothing seems to have been published by the Dutch naturalists who accompanied the two expeditions, with the exception of a slight sketch of the zoology of the second voyage by Dr. F. H. van Lidth de Jeude, in a pamphlet published at Amsterdam, entitled "*De Verslagen omtrent den tocht met de Willem Barents naar en in de Ijszee, in den zomer van 1879 (Uitgegeven vanwege het Aardrijkskundig Genootschap, Bijblad no. 6);*" and, as nearly two years have now elapsed since the return of the first expedition, it seems desirable that the information obtained by the study of Mr. Grant's collections should be made available to the scientific public without further delay. Barents Sea was visited by the Austro-Hungarian North-Pole Expedition under Weyprecht and Payer in 1872-1874; and dredgings were made, but mostly to the north and east of the ground examined by the 'Willem Barents' expeditions. A portion of the Invertebrates obtained were catalogued and described by Dr. Emil von Marenzeller in a pamphlet published at Vienna in 1877*. Many of the species he mentions were obtained by the Dutch expedition, and, in addition, many that were not met with by the Austrians.

It is evident that Barents Sea abounds with animal life in a very marked degree. In fact all the explorers of these northern regions seem to have been deeply impressed with the extraordinary richness of the marine fauna of the Arctic seas to the north of Lapland, Russia, and Siberia. Austrians, Dutch, and Swedes alike dwell on the amazing number of

* '*Die Coelenteraten, Echinodermen und Würmen der k.-k. österreichisch-ungarischen Nordpol-Expedition bearbeitet von Dr. Emil v. Marenzeller.*'

animals, both species and individuals, which the dredge brought to the surface.

Certain spots yielded a great variety of animal life, as the following examples will testify:—

On July 18, 1878, in lat. $73^{\circ} 41' 12''$ N., long. $22^{\circ} 58' 30''$ E., at a depth of 210 fathoms, occurred *Halichondria Hyndmani*, *Otenodiscus crispatus*, *Archaster tenuispinus* and *A. bifrons*, *Ophiopholis bellis*, *Ophiacantha spinulosa*, *Nymphon hirtipes*, *Membranipora arctica* and *M. monostachys*, *Porcella struma*, *Terebratula caput-serpentis*, var. *septentrionalis*, *Pecten grœnlandicus* and *P. Hoskynsi*, *Modiolaria discors*, *Nucula tenuis*, *Leda pernula* and *L. intermedia*, *Arca pectunculoïdes*, *Astarte crenata*, and *Siphodentalium vitreum*.

On July 30, 1878, in lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., at a depth of 160 fathoms, occurred *Calycella fastigiata*, *Thuiaria articulata*, *Sertularia cupressina*, *Lafoëa grandis*, *Selaginopsis decemserialis*, *Ammothea glomerata*, *Otenodiscus crispatus*, *Ophiocten sericeum*, *Astrophyton Lamarckii*, *Ennea nodosa*, *Nephthys ciliata*, *Cystenides hyperborea*, *Terebellides Strœmii*, *Phascolosoma Strombi*, *Munnopsis typica*, *Acanthostepheia Malmgreni*, *Tritopsis Helleri*, *Unciola leucopes*, *Acanthonotosoma inflatum*, *Idotea Sabinii*, *Nymphon hirtipes*, *Flustra membranaceo-truncata*, *Mucronella scutulata* and *M. simplex*, *Crisia eburneo-denticulata*, *Diastopora obelia*, *Hornera* sp., *Alcyonidium excavatum*, *Barentsia bulbosa*, *Phylactella grandis*, *Myrionozom subgracile*, *Flustra solida* (= *Eschara palmata*), *Scalaria grœnlandica*, and *Bulla propinqua*.

On August 1, 1878, in lat. $76^{\circ} 58'$ N., long. $45^{\circ} 40'$ E., in 110 fathoms, at the edge of the pack-ice, individuals of the following species were extremely numerous:—Five species of fishes: *Centridermichthys uncinatus*, *Icelus hamatus*, *Agonus decagonus*, *Liparis vulgaris*?, and *Hippoglossoides limandoides*; but the specimens were all of small size. Four crustaceans: *Crangon boreas*, *Sabinea septemcarinata*, *Pandalus annulicornis*, and *Idotea Sabinii*. *Nymphon Stromii* and *N. robustum*. The Echinoderms were *Strongylocentrotus dröbachiensis*, *Crossaster papposus*, var. *affinis*, *Otenodiscus crispatus*, *Asterias stellionura*, *Ophiopleura arctica*, and *Astrophyton Lamarckii*. No Mollusca were brought to me from this dredging by Mr. Grant; but not far off from this spot on the previous day, in 130 fathoms, in sand and mud, *Lima subovata* was very numerous, with several other shells.

Another rich neighbourhood appears to be off Mesjuds-jarrskii Island, on the south-west of Novaya Zemlya, in from 62 to 67 fathoms, where three species of fish, fourteen species of Mollusca, seven species of Crustacea (including *Alauna*

Goodsiri), nine species of Echinodermata (including *Antedon Eschrichtii*), ten species of Annelids, four species of Polyzoa, six Hydroids (including *Myriothele phrygia*, Fab., and a very large species of *Lucernaria*), and several sponges were obtained on July 31 and August 1, 1879.

When it is considered that Mr. Grant only took such things as were not required by the Dutch naturalists, the above lists will convey some idea of the profusion of animal life in Barents Sea.

Certain animals seem almost universally distributed in this region, such as the interesting tube-worm *Cystenides hyperborea*, which came up in almost every dredging and in great numbers. The Ophiurids *Ophiocten sericeum*, *Ophiopholis bellis*, and *Ophiacantha spinulosa* also occurred almost everywhere, and sometimes choked the dredge with their masses. *Ctenodiscus crispatus* was also very generally distributed at all depths from 62 to 210 fathoms.

Off the north coast of Norway, just outside Barents Sea, the Ophiurids were scarce, and, on the contrary, Brachiopoda, which were rarely met with in Barents Sea, abounded, at least in individuals.

The annexed list of species (p. 258) is tabulated in a similar manner to that given by Dr. Marenzeller; and the two can therefore be readily compared with each other.

SPONGIA.

Mr. Henry J. Carter, F.R.S., has kindly furnished the following descriptions of the two new Sponges :—

Suberites montalbidus.

Form monticular. Colour grey-white. Surface corrugated. Vents, one large on the summit, naked, the rest small, on the sides. Texture soft, matted. Spicules of two kinds, viz. :—1, skeleton, large, subpinlike, head variable in shape; 2, flesh-spicule, minute, shaft cylindrical, straight or curved, pointed at each end and inflated in the centre. Size of specimen 11-12ths inch in diameter at the base, 8-12ths inch high.

Suberites montiniger.

Form monticular. Colour grey-black. Surface even. Vents, one large at the summit, fringed, the rest small, on the sides. Texture soft, matted. Spicules of one kind only, viz. skeleton, large, subpinlike, head oval, elongated. Size of specimen 13-12ths inch in diameter at the base, 8-12ths inch high.

Hab. Marine, on hard objects.

Loc. Barents Sea, near S.W. end of Novaya Zemlya, in lat. $71^{\circ} 6' N.$, long. $50^{\circ} E.$ 31st July, 1879. Depth 62 fathoms.

Obs. Both these sponges are monticular in form, but specifically different, as will be seen in comparing the above descriptions. *Suberites montalbidus*, however, is, with the exception of form and slight differences in the spiculation, the same as *Suberites domuncula*, Sdt., = *Halichondria suberea*, Johnston, common on the British shores. The differences in spiculation chiefly consist in the pinlike spicules of the former not having such a globular or defined head, and the flesh-spicule for the most part being pointed at each end instead of obtuse. Marenzeller mentions the following species collected by the Austro-Hungarian expedition:—

Cacospongia Schmidtii, Mar.; *Chalinula cavernosa*, Mar.; *Isodictya tenera*, Mar.; *Stylocordyla longissima*, G. O. Sars; *Thecophora semisuberites*, Schm., and *T. elongata*, Mar.; *Rinalda uberrima*, Schm.; *Halicnemis hemisphaerica*, M. Sars; *Microciona ambigua*, Bowbk.; *Cladorhiza abyssicola*, M. Sars; *Ascetta coriacea*, Mont.; *Sycaltis glacialis*, Hæck.; and *Sycandra utriculus*, Schm.

HYDROZOA.

The Rev. Thomas Hincks, B.A., F.R.S., has drawn up the following list of the Hydrozoa:—

Order HYDROIDA.

Suborder Athecata.

Genus MYRIOTHELA, Sars.

? *Myriothela phrygia*, Fabricius.

[Not *M. phrygia* of Allman and Hincks.]

This is distinct from the British species, and probably identical with *Lucernaria phrygia* of Fabricius.

Lat. $71^{\circ} 6' N.$, long. $50^{\circ} E.$ Near S.W. end of Novaya Zemlya, 62 fms. [Greenland (*Fabr.*), Norway (*Sars*); off Halifax, N. S., in 52 fms. (*Verrill*).]

Genus EUDENDRIUM, Ehrenberg.

Eudendrium ? sp.

A fragment only occurs, which does not enable me to determine the species. The stem is compound below, but simple

List of Animals collected in Barents Sea by W. J. A. Grant, Esq., in 1878 and 1879.

Name.	Date.	North latitude.	East longitude.	Depth in fathoms.	Remarks.
PROTOZOA.					
<i>Triloculina tricarinata</i> , D'Orb.	17, 7, 78	74° 4' 0"	23° 0' 0"	220	Very large.
COELENTERATA.					
SPONGIA.					
<i>Halichondria Hyndmani</i> , Bk.	18, 7, 78	73° 41' 12"	22° 58' 30"	210	See description. " " Attached to a coral.
<i>Suberites montalbidus</i> , H. J. Carter, n. sp.	31, 7, 79	71° 6' 0"	50° 0' 0"	62	
— <i>montiniger</i> , H. J. Carter, n. sp.	31, 7, 79	71° 6' 0"	50° 0' 0"	62	
<i>Isodictya</i> (species undetermined)	31, 7, 79	71° 6' 0"	50° 0' 0"	62	
HYDROZOA.					
<i>Hydroida.</i>					
<i>Myriothele phrygia</i> , Fab.	31, 7, 79	71° 6' 0"	50° 0' 0"	62	See note.
<i>Eudendrium</i> species.	15, 7, 79	72° 32' 3"	36° 29' 5"	128	A fragment only, probably <i>E. ramosum</i> .
<i>Calycella fastigiata</i> , Alder	30, 7, 78	75° 16' 6"	45° 19' 36"	160	On <i>Alcyonidium</i> .
" "	15, 7, 79	72° 32' 3"	36° 29' 5"	128	On <i>Gemellaria</i> .
" "	1, 8, 79	71° 23' 0"	49° 38' 0"	67	" "
<i>Lafoëa grandis</i> , Hincks	30, 7, 78	75° 16' 6"	45° 19' 36"	160	On <i>Myricosum</i> .
<i>Lafoëina tenuis</i> , M. Sars	31, 7, 79	71° 6' 0"	50° 0' 0"	62	Off Novaya Zemlya.
" "	1, 8, 79	71° 23' 0"	49° 38' 0"	67	On <i>Cellularia Peachii</i> .
<i>Filellum serpens</i> , Hassall.	14, 7, 78	74° 6' 0"	18° 5' 0"	25	Off Bear Island; very fine, on <i>Balanus</i> .
" "	17, 7, 78	74° 5' 0"	23° 0' 0"	220	On <i>Sertularella</i> .
<i>Halecium muricatum</i> , Ellis & Sol.	14, 7, 78	74° 6' 0"	18° 5' 0"	25	Amongst <i>Gemellaria</i> .

<i>Sertularia tricuspidata</i> , <i>Alder</i>	14, 7, 78	74	6	0	18	5	0	25	Very abundant.
"	17, 7, 78	74	5	0	23	0	0	220	
<i>Sertularia quadricornuta</i> , <i>Hincks</i> , n. sp.	14, 7, 78	74	6	0	18	5	0	25	Off Bear Island.
<i>Thuiaria articulata</i> , <i>Fallas</i>	30, 7, 78	75	16	6	45	19	36	160	
<i>Selaginopsis decemserialis</i> , <i>Mereschkowsky</i> ..	14, 7, 78	74	6	0	18	5	0	25	Off Bear Island.
"	17, 7, 78	74	5	0	23	0	0	220	
<i>Sertularia cupressina</i> , <i>L.</i>	30, 7, 78	75	16	6	45	19	36	160	One specimen.
<i>Calycozoa.</i>									
<i>Lucernaria campanulata</i> ? <i>M.-Ed.</i>	1, 8, 79	71	23	0	49	38	0	67	Very large.
ACTINOZOA.									
<i>Ammothea glomerata</i> , <i>Carter</i> , n. sp.	29, 7, 78	74	6	42	45	1	42	160	See description.
"	30, 7, 78	75	16	6	45	19	36	160	On <i>Tubularia</i> .
"	1, 8, 78	76	58	0	45	40	0	110	
"	31, 7, 79	71	6	0	50	0	0	62	Attached to shells.
<i>Gerssemia florida</i> , <i>Rathke</i>	1, 8, 79	71	28	0	49	38	0	67	
"	15, 7, 79	72	32	3	36	29	5	128	
ECHINODERMATA.									
HOLOTHUROIDEA.									
<i>Eurygus scaber</i> , <i>Lütken</i>	31, 7, 79	71	6	0	50	0	0	62	Off S.W. coast of Novaya Zemlya.
ECHINOIDEA.									
<i>Strongylocentrotus dröbachiensis</i> (<i>O. F. Müller</i>)..	1, 8, 78	76	58	0	45	40	0	110	
"	19, 7, 79	75	23	5	38	39	5	88	
"	31, 7, 79	71	6	0	50	0	0	62	Off S.W. coast of Novaya Zemlya.
ASTEROIDEA.									
<i>Asteracanthion greenlandicus</i> , <i>Stp.</i>	31, 7, 79	71	6	0	50	0	0	62	Off S.W. coast of Novaya Zemlya.
<i>Crossaster papposus</i> (<i>Linck</i>), var. <i>affinis</i> , <i>Brandt</i> ..	1, 8, 78	76	58	0	45	40	0	110	8 rays.
<i>Ctenodiscus crispatus</i> , <i>Ratz.</i> , = <i>corniculata</i> (<i>Linck</i>)	18, 7, 78	73	41	12	22	58	30	210	
"	24, 7, 78	72	5	6	37	57	18	140	
"	30, 7, 78	75	16	6	45	19	36	160	
"	1, 8, 78	70	58	0	45	40	0	110	

List (continued).

Name.	Date.	North latitude.	East longitude.	Depth in fathoms.	Remarks.
<i>Ctenodiscus crispatus</i> , Retz., = <i>corniculata</i> (Linné)	15, 7, 79	2° 32'	3° 36' 39"	128	
"	31, 7, 79	71° 6'	0° 50' 0"	62	
<i>Archaster tenuispinus</i> , Düb. & Kor.	18, 7, 78	73° 41'	22° 58' 30"	210	
"	29, 7, 78	74° 6'	42° 45' 1 42	160	
<i>Archaster bifrons</i> , Wy. Thomson	18, 7, 78	73° 41'	22° 58' 30"	210	Young.
<i>Asterias stellionura</i> , Perrier	1, 8, 78	76° 58'	0° 45' 40' 0"	110	One specimen.
OPHIUROIDEA.					
<i>Ophioglypha Sarsii</i> , Lütken.	31, 7, 79	71° 6'	0° 50' 0"	62	
— <i>robusta</i> (Ayres)	31, 7, 79	71° 6'	0° 50' 0"	62	
<i>Ophiocten sericeum</i> (Forbes)	14, 7, 78	74° 6'	0° 18' 5' 0"	25	
"	29, 7, 78	74° 6'	42° 45' 1 42	160	
"	30, 7, 78	75° 16'	45° 19' 36"	160	Very abundant.
"	31, 7, 78	76° 31' 18"	45° 36' 30"	130	
"	15, 7, 79	75° 23' 5'	38° 39' 5'	128	Very abundant.
"	31, 7, 79	71° 6'	0° 50' 0"	62	
<i>Ophiopholis bellis</i> (Linné)	18, 7, 78	73° 41' 12"	22° 58' 30"	210	
"	29, 7, 78	74° 6' 42"	45° 1 42	160	
"	31, 7, 78	76° 31' 18"	45° 36' 30"	130	
"	15, 7, 79	72° 32' 3'	36° 39' 5'	128	Very abundant.
"	1, 8, 79	71° 23' 0'	49° 38' 0'	67	
"	1, 8, 78	76° 58' 0'	45° 40' 0"	110	
<i>Ophiopleura arctica</i> , Duncan	18, 7, 78	73° 41' 12"	22° 58' 30"	210	
<i>Ophiacantha spinulosa</i> , Müll. & T.	29, 7, 78	74° 6' 42"	45° 1 42	160	
"	31, 7, 78	76° 31' 18"	45° 36' 30"	130	Very abundant.
"	15, 7, 79	72° 32' 3'	36° 39' 5'	128	" "
"	31, 7, 79	71° 6' 0'	50' 0' 0"	62	" "
<i>Astrophyton Lamarckii</i> , Müll. & T.	30, 7, 78	75° 16' 6"	45° 19' 36"	160	Very young specimen.
"	1, 8, 78	76° 58' 0'	45° 40' 0"	110	Young (see note).

CRINOIDEA.	1, 8, 79	71	23	0	49	38	0	67	Off S.W. coast of Novaya Zemlya.
<i>Antedon Eschrichtii (Müll.)</i>									
VERMES.									
ANNELIDA.									
<i>Evadne impar (Johnst.)</i>	31, 7, 79	71	6	0	50	0	0	62	Off S.W. coast of Novaya Zemlya.
<i>Antinos Sarsi, Knabg.</i>	31, 7, 79	71	6	0	50	0	0	62	
<i>Lagisca rarispina, Sars</i>	31, 7, 79	71	6	0	50	0	0	62	
<i>Ennosa nodosa, Sars</i>	30, 7, 78	75	16	6	45	19	38	160	
<i>Northia hyperborea, Hansen</i>	31, 7, 79	71	6	0	50	0	0	62	In shelly tubes; off S.W. coast of Novaya Zemlya.
<i>Nychia cirrosa, Pallas</i>	31, 7, 79	71	6	0	50	0	0	62	In shells of <i>Trochus</i> .
<i>Nereis zonata, Mgrn.</i>	31, 7, 79	71	6	0	50	0	0	62	
<i>Nephtys ciliata, O. F. Müller</i>	30, 7, 78	75	16	6	45	19	38	160	
"	15, 7, 79	72	32	3	36	29	5	128	
"	31, 7, 79	71	6	0	50	0	0	62	
<i>Ephesia gracilis, H. Rathke</i>	15, 7, 79	72	32	3	36	29	5	128	
<i>Trophonia Whiteavesii, Müll. & T., variety</i>	15, 7, 79	72	32	3	36	29	5	128	
<i>Brada arenosa, McIntosh (= granosa, Stp.)</i>	15, 7, 79	72	32	3	36	29	5	128	
<i>Terebellides Stroomii, Sars</i>	30, 7, 78	75	16	6	45	19	38	160	
<i>Cystenides hyperborea, Mgrn.</i>	24, 7, 78	72	5	6	37	57	18	140	Plentiful.
"	29, 7, 78	74	6	42	45	1	42	160	
"	30, 7, 78	75	16	6	45	19	38	160	
"	31, 7, 78	76	31	10	45	36	30	130	
"	19, 7, 79	75	23	5	38	39	5	88	
"	31, 7, 79	71	6	0	50	0	0	62	
<i>Sabella spitsbergenensis, Mgrn.</i>	31, 7, 79	71	6	0	50	0	0	62	
<i>Dasychone</i> species	31, 7, 79	71	6	0	50	0	0	62	
GEPHYREA.									
<i>Phascolosoma Strombi, Mont.</i>	29, 7, 78	74	6	42	45	14	2	160	In broken tubes of <i>Cystenides</i> .
"	30, 7, 78	75	16	6	45	19	38	160	
"	19, 7, 79	75	23	5	38	39	5	88	

List (continued).

Name.	Date.	North latitude.	East longitude.	Depth in fathoms.	Remarks.
ARTHROPODA.					
CRUSTACEA.					
<i>Cirripedia.</i>					
<i>Balanus balanoides</i> , L.	14, 7, 78	74 6 0	18 5 0	25	Very abundant off Bear Island.
— porcatus, <i>Da Costa</i>	23, 8, 78	73 30 0	54 30 0		On dead <i>Fusus</i> -shell on shore, Matotschkin Shar.
<i>Edriophthalma.</i>					
<i>Anonyx nugax</i> (<i>Phipps</i>)	31, 7, 79	71 6 0	50 0 0	62	
<i>Acanthonotosoma inflatum</i> (<i>Kröyer</i>)	30, 7, 78	75 16 6	45 19 36	160	
<i>Gammaracanthus loricatus</i> (<i>Sabine</i>)	31, 7, 79	71 6 0	50 0 0	62	
<i>Amphithoe læviscula</i> , <i>Bell</i> ?	15, 7, 79	72 32 3	36 39 5	128	
<i>Acanthostephia Malmgreni</i> (<i>Goës</i>)	30, 7, 78	75 16 6	45 19 36	160	
<i>Tritopsis Helli</i> , <i>Boeck</i>	14, 7, 78	74 6 0	18 5 0	25	Off Bear Island.
"	30, 7, 78	75 16 6	45 19 36	160	
<i>Munnopsis typica</i> , <i>Sars</i>	30, 7, 78	75 16 6	45 19 36	160	
<i>Unciola leucopes</i> (<i>Kröyer</i>)	30, 7, 78	75 16 6	45 19 36	160	
<i>Hyperia cyaneæ</i> (<i>Sab.</i>)	15, 7, 79	72 23 3	36 39 5	128	Immature.
<i>Idotea Sabini</i> , <i>Kröyer</i>	30, 7, 78	75 16 6	45 19 36	160	Small specimens.
"	31, 7, 78	76 31 18	45 36 30	130	On sand and mud.
"	1, 8, 78	76 58 0	45 40 0	110	Large and numerous.
<i>Podophthalma.</i>					
<i>Alauna Goodsiri</i> , <i>Bell</i>	31, 7, 79	71 6 0	50 0 0	62	
<i>Hippolyte macilentæ</i> , <i>Kröyer</i>	24, 7, 78	72 5 6	37 57 18	140	
— <i>Belcheri</i> , <i>Bell</i>	31, 7, 79	71 6 0	50 0 0	62	
<i>Pandalus annulicornis</i> , <i>Leach</i>	1, 8, 78	76 58 0	45 40 0	110	

Crangon boreas, <i>Phipps</i>	1, 8, 78	76 58	0	45 50	0	110	Very numerous.
Sabinea septemarinata, <i>Sabine</i>	1, 8, 78	76 58	0	45 40	0	110	Very fine and numerous; ♀ loaded with eggs.
"	15, 7, 79	72 32	3	36 39	5	128	Small specimens.
"	31, 7, 79	71 6	0	50 0	0	62	
"	1, 8, 79	71 23	0	49 38	0	67	Off Bear Island.
Eupagurus pubescens, <i>Krøyer</i>	14, 7, 78	74 6	0	18 5	0	25	Off Novaya Zemlya, in shells of
"	31, 7, 79	71 6	0	50 0	0	62	<i>Natica</i> , <i>Buccinum</i> , &c.
<i>Podosomata.</i>							
Nymphon hirtipes, <i>Ball</i>	18, 7, 78	73 41	12	22 58	30	210	
"	30, 7, 78	75 16	6	45 19	36	160	
"	1, 8, 79	71 23	0	49 38	0	67	
Nymphon Strömii, <i>Krøyer</i>	1, 8, 78	76 58	0	45 40	0	110	Very plentiful.
— robustum, <i>Ball</i>	1, 8, 78	76 58	0	45 40	0	110	" "
— gracile, <i>Leach</i>	31, 7, 79	71 6	0	50 0	0	62	
— grossipes, <i>L.</i>	1, 8, 79	71 23	0	49 38	0	67	
MOLLUSCA.							
MOLLUSCORDEA.							
<i>Polyzoa.</i>							
Gemellaria loricata, <i>Busk</i>	14, 7, 78	74 6	0	18 5	0	25	Very fine and abundant off Bear Island.
"	15, 7, 79	72 32	3	36 29	5	128	Dark-coloured variety.
Cellularia Peachii, <i>Busk</i>	31, 7, 79	71 23	0	49 38	0	67	Off Novaya Zemlya.
Menipea ternata, <i>Ellis & Sol.</i>	14, 7, 78	74 6	0	18 5	0	25	Off Bear Island.
"	31, 7, 79	71 6	0	50 0	0	62	Off Novaya Zemlya.
"	1, 8, 79	71 23	0	49 38	0	67	Abundant.
Menipea ternata, <i>Ellis & Sol.</i> , forma gracilis, <i>Smith.</i>	15, 7, 79	72 32	3	36 29	5	128	
Bugula Murrayana, <i>Johnst.</i>	15, 7, 79	72 32	3	36 29	5	128	
Fusula membranacea-truncata, <i>Smith</i>	30, 7, 78	75 16	6	45 19	36	160	
Membranipora monostachys, <i>Busk</i>	18, 7, 78	73 41	12	22 58	30	210	On stones, &c.
— lineata, <i>L.</i>	18, 7, 78	73 41	12	25 58	30	210	" "
— craticula, <i>Alder</i>	14, 7, 78	74 6	0	18 5	0	25	" "

List (continued).

Name.	Date.	North latitude.	East longitude.	Depth in fathoms.	Remarks.
Membranipora sophiæ, <i>Busk</i>	14, 7, 78	° 74	' 6	25	On stones &c.
— arctica, <i>Smit</i>	14, 7, 78	74	6	0	Very common on stones, shells, &c.
"	18, 7, 78	73	41	22	25 {
Microperella ciliata, <i>Pallas</i>	14, 7, 78	74	6	0	210 {
Portia tubulosa, <i>Norman</i>	14, 7, 78	74	6	0	25
Myriozoum subgracile, <i>D'Orb.</i>	14, 7, 78	74	6	0	25
Schizoporella sinuosa, <i>Busk</i>	30, 7, 78	75	16	6	On stones.
— hyalina, <i>L.</i>	24, 7, 78	72	55	0	On stones.
— plana, <i>Dawson</i>	14, 7, 78	74	6	0	On stones.
(Myriozoum crustaceum, <i>Smit</i> .)	14, 7, 78	74	6	0	On stones.
Porella struma, <i>Norman</i>	18, 7, 78	73	41	12	25
Phylactella ? grandis, <i>Hincks</i> , n. sp.	30, 7, 78	75	16	6	210
Mucronella scutulata, <i>Busk</i>	30, 7, 78	75	16	6	160
— simplex, <i>Hincks</i> , n. sp.	30, 7, 78	75	16	6	160
Eschara (Flustra) solida, <i>Stimpson</i> (= Eschara palmata, <i>Sars</i> , and Escharella palmata, <i>Smit</i> .)	30, 7, 78	75	16	6	On stones.
Eschara glabra, <i>Hincks</i> , n. sp.	30, 7, 78	75	16	6	Abundant.
Cellepora, ? n. sp.	15, 7, 79	72	32	3	160
Crisia eburneo-denticulata, <i>Smit</i>	30, 7, 78	75	16	6	On <i>Endendrium</i> .
Diatopora obelia, <i>Johnst.</i>	30, 7, 78	75	16	6	160
Hornera, ? sp.	30, 7, 78	75	16	6	On <i>Flustra solida</i> , <i>Stp.</i>
Lichenopora verrucaria, <i>Fab.</i>	14, 7, 78	74	6	0	160
Alyconidium excavatum, <i>Hincks</i> , n. sp.	30, 7, 78	75	16	6	On <i>Balanus</i> .
Arachnidium simplex, <i>Hincks</i> , n. sp.	31, 7, 79	71	6	0	Abundant, attached to tubes of <i>An-</i>
Buskia nitens, <i>Alder</i>	15, 7, 79	72	32	3	nelids.
Barentsia bulbosa, <i>Hincks</i> , n. sp.	30, 7, 78	75	16	6	On <i>Endendrium</i> .
Loxosoma singulare, <i>Keferstein</i>	31, 7, 79	71	6	0	160
					On <i>Menippea</i> .

<i>Brachiopoda.</i>										
<i>Terebratula caput-serpentis</i> , <i>L.</i> , var. <i>septentrionalis</i>	18, 7, 78	73 41 12	22 58 30	210	Very large.					
<i>Terebratula spitzbergensis</i> , <i>Davidson</i>	19, 7, 79	75 23 5	38 39 5	88						
MOLLUSCA PROPER.										
<i>Conchifera.</i>										
<i>Pecten greenlandicus</i> , <i>G. B. Sowerby</i>	18, 7, 78	73 41 12	22 58 30	210						
" <i>Hoskynsi</i> , <i>Forbes</i>	19, 7, 79	75 23 5	38 39 5	88						
"	17, 7, 78	74 5 0	23 0 0	220						
<i>Lima subovata</i> , <i>Jaffreys</i>	31, 7, 78	76 31 18	45 36 30	130						
<i>Modiolaria discors</i> , <i>L.</i>	14, 7, 78	78 6 0	18 5 0	25						
"	18, 7, 78	73 41 12	22 58 30	210						
<i>Nucula tenuis</i> , <i>Mont.</i> , var. <i>inflata</i> , <i>Hancock</i>	14, 7, 78	78 6 0	18 5 0	25	Very fine.					
"	18, 7, 78	73 41 12	22 58 30	210						
"	20, 7, 78	74 6 42	45 1 42	160						
"	19, 7, 79	75 23 5	38 39 5	88						
<i>Leda limatula</i> , <i>Say</i>	19, 7, 79	75 23 5	38 39 5	88						
"	31, 7, 79	71 6 0	50 0 0	62						
" <i>Müll.</i>	14, 7, 78	78 6 0	18 5 0	25						
<i>Leda pernula</i> , <i>Müll.</i>	18, 7, 78	73 41 12	22 58 30	210						
"	24, 7, 78	72 5 6	37 57 18	140						
"	29, 7, 78	74 6 42	45 1 42	160						
"	31, 7, 78	76 31 18	45 36 30	130	Very fine.					
"	19, 7, 79	75 23 5	38 39 5	88						
"	31, 7, 79	71 6 0	50 0 0	62						
"	14, 7, 78	78 6 0	18 5 0	25						
<i>Leda intermedia</i> , <i>M. Sars</i>	18, 7, 78	73 41 12	22 58 30	210						
"	19, 7, 79	75 23 5	38 39 5	88						
<i>Leda abyssicola</i> , <i>Torell</i>	29, 7, 78	74 6 42	45 1 42	160						
"	31, 7, 78	76 31 18	45 36 30	130						
<i>Arca glacialis</i> , <i>Gray</i>	14, 7, 78	78 6 0	18 5 0	25						
"	24, 7, 78	72 5 6	37 57 18	140						
"	24, 7, 78	72 5 30	37 57 18	145						
"	29, 7, 78	74 6 42	45 1 42	160						

List (continued).

Name.	Date.	North latitude.	East longitude.	Depth in fathoms.	Remarks.
<i>Area glacialis, Gray</i>	31, 7, 78	56 31 18	45 36 30	130	
"	19, 7, 79	75 23 5	38 39 5	88	
<i>Area pectunculoides, Seacchi</i>	18, 7, 78	73 41 12	22 58 30	210	Large specimens.
<i>Cardium islandicum, L.</i>	19, 7, 79	75 23 5	38 39 5	88	Young.
"	31, 7, 79	71 6 0	50 0 0	62	"
<i>Astarte crenata, Gray</i>	18, 7, 78	73 41 12	22 58 30	210	
"	24, 7, 78	72 5 30	37 57 18	145	
"	19, 7, 79	75 23 5	38 39 5	88	
<i>Astarte fabula, Reeve</i>	19, 7, 79	75 23 5	38 39 5	88	
"	31, 7, 79	71 6 0	50 0 0	62	
<i>Astarte borealis, Chemnitz</i>	31, 7, 79	71 6 0	50 0 0	62	
<i>Tellina calcarea, Chemn.</i>	24, 7, 78	72 5 30	37 57 18	145	
"	19, 7, 79	72 23 5	38 39 5	88	
"	31, 7, 79	71 6 0	50 0 0	62	
<i>Saxicava rugosa, L., var. præcisa</i>	14, 7, 78	78 6 0	18 5 0	25	
"	24, 7, 78	72 5 6	37 57 18	140	
"	24, 7, 78	72 5 30	37 57 18	145	
"	31, 7, 78	76 31 18	45 36 3	130	
"	19, 7, 79	75 23 5	38 39 5	88	Very large.
<i>Solenocoelia.</i>					
<i>Siphonotalium vitreum, M. Sars</i>	17, 7, 78	74 5 0	23 0 0	220	
"	18, 7, 78	73 41 12	22 58 30	210	
"	29, 7, 78	74 6 42	45 1 42	160	
<i>Gastropoda.</i>					
<i>Lepeta caeca, Müll.</i>	24, 7, 78	72 5 6	37 57 18	140	
"	24, 7, 78	72 5 30	37 57 18	145	
"	15, 7, 79	72 32 3	36 39 5	123	

<i>Trochus greenlandicus, Chemn.</i>	14, 7, 78	78	6	0	18	5	0	25	
<i>Trochus varicosus, Mighels</i>	29, 7, 78	74	6	42	45	1	42	160	
"	19, 7, 79	75	23	5	38	39	5	88	
"	31, 7, 79	71	6	0	50	0	0	62	
<i>Trochus cinereus, Couthouy, var. arctica</i>	31, 7, 79	71	6	0	50	0	3	62	
<i>Scalardia greenlandica, Chemn.</i>	30, 7, 78	75	16	6	45	19	36	160	
<i>Natica affinis, Gmelin</i>	14, 7, 78	78	6	0	18	5	0	25	
"	29, 7, 78	74	6	42	45	14	2	160	
<i>Natica greenlandica, Beck</i>	29, 7, 78	74	6	42	45	14	2	160	
"	15, 7, 79	72	32	3	36	39	5	128	Young.
"	31, 7, 79	71	6	0	50	0	0	62	Young.
<i>Cancellaria viridula, Fabricius, var.</i>	15, 7, 79	72	32	3	36	39	5	128	Egg-capsule and spawn.
<i>Buccinum undatum, L.</i>	14, 7, 78	74	6	0	18	5	0	25	Young.
"	31, 7, 78	76	31	18	45	36	30	130	Young.
<i>Buccinum glaciale, L.</i>	31, 7, 79	71	6	0	50	0	0	62	"
— ciliatum, Fabr., var.	31, 7, 79	71	6	0	50	0	0	62	"
<i>Fusus tortuosus, Reeve</i>	19, 7, 79	75	23	5	38	39	5	88	"
"	31, 7, 79	71	6	0	50	0	0	62	"
<i>Fusus despectus, L., var.</i>	23, 8, 78	73	30	0	54	30	0	..	[Shar.
<i>Pleurotoma bicarinata, Couth.</i>	29, 7, 78	74	6	42	45	1	42	160	Dead shell, on shore, Matotschkin
— turricula, Mont., var. nobilis	17, 7, 78	74	5	0	23	0	0	220	
— pyramidalis, Ström	31, 7, 79	71	6	0	50	0	0	62	
<i>Bulla propinqua, M. Sars</i>	29, 7, 78	74	6	42	45	1	42	160	
"	30, 7, 78	75	16	6	45	19	36	160	
<i>Bulla striata, Brown.</i>	19, 7, 79	75	23	5	38	39	5	88	
VERTEBRATA.									
PISCES.									
<i>Centridermichthys uncinatus, Reinh.</i>	1, 8, 78	76	58	0	45	40	0	110	At edge of pack-ice.
<i>Icelus hanatus, Kröyer</i>	1, 8, 78	76	58	0	45	40	0	110	"
<i>Agonus decagonus, Bl. Schn.</i>	1, 8, 78	76	58	0	45	40	0	110	"
"	31, 7, 79	71	6	0	50	0	0	62	Young.
<i>Aspidophoroides monopterygius (Block)</i>	31, 7, 79	71	6	0	50	0	0	62	One specimen in a bad state.
<i>Liparis vulgaris?, Flem.</i>	1, 8, 78	76	58	0	45	40	0	110	
<i>Blennius species</i>	31, 7, 79	71	6	0	50	0	0	62	
<i>Hippoglossoides limandoides (Bl.)</i>	1, 8, 78	76	58	0	45	40	0	110	At edge of pack-ice.

towards the extremity, and of a rather dark horn-colour. Branches are given off irregularly, which are ringed at the base and at intervals above it; they bear alternate polypiferous ramules, similarly annulated. The form is nearly allied to *E. ramosum*, if not identical with it.

Lat. $72^{\circ} 32' 3''$ N., long. $36^{\circ} 29' 5''$ E., 128 fms.

Suborder Thecaphora.

Genus CALYCELLA, Hincks.

Calycella fastigiata, Alder.

Lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms., on *Acyonidium*; lat. $72^{\circ} 32' 3''$ E., long. $36^{\circ} 29' 5''$ E., 128 fms., on *Gemellaria*, &c.; lat. $71^{\circ} 23'$ N., long. $49^{\circ} 38'$ E., 67 fms.

In the Barents-Sea specimens the pedicels are long and distinctly annulated; in the British examples which I have seen they are smooth.

Genus LAFOËA, Lamouroux.

Lafoëa grandis, Hincks.

Lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., on *Myrizoum*, 160 fms. [Davis Straits, 100 fms. (*Wallich*); Atlantic coasts of America (*Verrill*).]

Genus LAFOËINA, M. Sars.

Lafoëina tenuis, Sars.

Lat. $71^{\circ} 23'$ N., long. $49^{\circ} 38'$ E., on *Cellularia Peachii*, in 67 fms., and lat. $71^{\circ} 6'$ N., long. 50° E., in 62 fms., off Novaya Zemlya.

Genus FILELLUM, Hincks.

Filellum serpens, Hassall.

Lat. 74° N., long. 23° E., in 220 fms., on *Sertularella tricuspidata*. Also very fine on *Balanus*, off Bear Island, in 25 fms.

Genus HALECIUM, Oken.

Halecium muricatum, Ellis and Solander.

Fragments with fine capsules amongst a mass of *Gemellaria loricata*, off Bear Island, in 25 fms.

Genus SERTULARELLA, Gray.

Sertularella tricuspidata, Alder.

Extremely abundant. Lat. 74° N., long. 23° E., in

220 fms.; and also off Bear Island, in 25 fms. As is usual in Arctic specimens, the capsules are present in profusion.

Sertularella quadricornuta, n. sp.

Bear Island, in 25 fms.

Genus THUIARIA, Fleming.

Thuiaria articulata, Pallas.

Lat. 75° 16' 6" N., long. 45° 19' 36" E., in 160 fms. Obtained by Austro-Hungarian expedition in lat. 76° 14' N., long. 58° 54' E., 100 metres. [British Seas.]

Genus SELAGINOPSIS, Allman.

Selaginopsis decemserialis, Mereschkowsky.

Lat. 74° N., long. 23° E., in 220 fms., and also off Bear Island, in 25 fms. [Northern Pacific Ocean (*Mereschkowsky*).]

Genus SERTULARIA, Linn.

Sertularia cupressina, L.

Lat. 75° 16' 6" N., long. 45° 19' 36" E. A single specimen from 160 fms. [British Seas; Labrador (*Packard*); Massachusetts Bay (*Agassiz*).]

Marenzeller mentions the following species which were not in Mr. Grant's collection:—*Corymorpha glacialis*, M. Sars; *Salacia abietina*, M. Sars; *Laföea dumosa*, Flem.; and *L. fruticosa*, M. Sars.

ACTINOZOA.

Mr. H. J. Carter considers the *Ammothea* which occurred abundantly in Barents Sea, and of which Mr. Grant brought me many specimens, distinct from *A. Luetkeni* of Marenzeller. He has drawn up the following description:—

Ammothea glomerata.

Cauliflower-like, consisting of more or less verruciform polyps aggregated into tuberoso, globular masses, arising from a longitudinally corrugated stem. Colour, in spirits, white. Texture cartilaginous. Surface irregular, on account of the variety in size and variable amount of projection of the polyps. Composed of subcartilaginous material charged with calcareous spicules. Polyps octotentacular, the largest about 1-24th inch in diameter; tentacles tubular, coated both inside

Ann. & N. Mag. Hist. Ser. 5. Vol. vi.

19

and out with spicules; spicules comparatively long, slightly curved, slightly clavate, more or less covered irregularly with small tubercles, 1-100th by 1-750th inch in their greatest dimensions.

Hab. Marine, growing on the stems of *Tubularia* and attached to shells, 62 to 160 fms., Barents Sea.

Obs. This differs from *Ammonothea Luetkeni*, Marenzeller (Cœlent., Echinod. u. Würmer der k.-k. österreichisch-ungarischen Nordpol-Exp. p. 16, Taf. iii. fig. 1: Wien, 1877, 4to), in the more or less agglomerated condition of the polyps, which in *A. Luetkeni* are separate like bunches of grapes. The latter has been found on the west coast of Greenland, and by Capt. Feilden in Smith Sound. With reference to the disputed point about the "retractile" nature of the polyps in *Ammonothea*, there can be no doubt that they are verruciform and composed of tubular tentacles coated all over, both inside (that is, towards the internal cavity of the polyps) and outside, with spicules, as above stated; while in *Lobularia* they are attached to a retractile tube *within* the radiated aperture of the surface. Hence, as stated by Kölliker (*apud* Marenzeller), they are *not retractile*. In *Ammonothea* the only tentacles are those seen on the outside of the cell, which is not imbedded in the matrix.

Marenzeller mentions the following Actinozoa as having been met with by the Austro-Hungarian expedition:—*Ammonothea Luetkeni*, Mar.; *Gersemia florida*, Rathke, and *G. loricata*, Mar.; *Umbellula encrinus*, L.; *Paragorgia arborea*, L.; *Urticina felina*, L.; *Phellia*, sp. ?; and *Zoanthus arcticus*, M. Sars.

Dr. van Lidth de Jeude mentions a species of *Actinia* as having been found in three dredgings. On one occasion two specimens were obtained attached to a large whelk(?)—shell. One or two corals brought home by Mr. Grant have not yet been determined.

ECHINODERMATA.

Crossaster papposus (Linck), var. *affinis* (Brandt).

The only specimen brought me by Mr. Grant had but eight rays.

Ophiopleura arctica, Duncan, in Ann. & Mag. Nat. Hist.
August and September 1878.

Only two specimens of this rare Ophiurid were brought me by Mr. Grant,

Astrophyton Lamarekii, Müll. et Tr.

Respecting the young specimen of this species taken in lat. 76° 58' N., long. 45° 40' E., in 110 fms., the Rev. A. M. Norman remarks as follows:—"The tubercles on the ribs of the disk make this, at first sight, to look like *A. Agassizii*; but the arms are not granular as in that species, and I therefore conclude that with advancing growth the tubercles on the ribs would disappear, instead of being developed into the irregular spines of *Agassizii*. I therefore suppose this specimen to be the young of *A. Lamarekii*."

Marenzeller mentions the following species, of which there were no examples in Mr. Grant's collections:—*Haplodactyla arctica*, Marenzeller; *Psolus Fabricii*, D. & K.; *Pteraster militaris*, O. F. Müller; *Stichaster albus*, Stp.; *Corethaster hispidus*, Wy. Thom.; *Amphiura Sundevalli*, M. & T.; *Astrophyton eucnemis*, M. & T.; *Antedon celtica*, Barrett; and *A. Sarsii*, D. & K.

Dr. van Lidth de Jeude alludes to the following:—*Molpadia*, sp., *Psolus*, sp., in great numbers on 15th July, 1879, in 138 fms.; *Cribella sanguinolenta*, on 17th July, in 127 fms.; *Ophiocoma nigra*, *Asteronyx Loveni*?, and a *Comatula* on 5th and 17th July, in 100 and 127 fms.

VERMES.

Dr. van Lidth de Jeude mentions the following species of worms, of which there are no representatives in Mr. Grant's collections:—*Ampharete arctica*, *Scione lobata*, *Onuphis Eschrichtii*, *Chlymene lumbricalis*, *Brada villosa* and *B. granulata*, and *Phyllodoce grælandica*.

Dr. Marenzeller enumerates the following species which were apparently not obtained by the Dutch expeditions:—*Cerebratulus angulatus*, O. F. Müller; *Scalibregma inflatum*, Rathke; *Cistenides granulata*, L.; *Ampharete Goesi*, Mgrn.; *Amphicteis Gunneri*, M. Sars; *Melinna cristata*, M. Sars; *Amphitrite cirrata*, O. F. Müller; *Thelepus circinatus*, F.; *Euchone tuberculosa*, Krøyer; *Chone infundibuliformis*, Krøyer; and *C. Duneri*, Mgrn.; *Spirorbis lucidus*, Mont. (this is probably the species which occurs on some of the Polyzoa brought home by Mr. Grant); *Hyalopotamus Claparedii*, Mar.; *Eucrante villosa*, Mgrn.; *Nephtys longosetosa*, Örst.; *Phyllodoce Luetkeni*, Mgrn.; *Syllis fasciata*, Mgrn.; *Nereis pelagica*, L.; *Northia conchylega*, M. Sars; *Glycera capitata*, Örst.; *Phascolosoma Örstedii*, Kef.; and *Echiurus forcipatus*, Rein.

The Gephyrean *Phascolosoma Strombi*, Mont., was found inhabiting repaired tubes of *Cystenides hyperborea*.

CRUSTACEA.

Dr. van Lidth de Jeude mentions *Hippolyte polaris* and *H. Sowerbyi* as having been dredged (the latter in great numbers) on 17th July, 1879. He also mentions *Pagurus Bernhardus* as being frequent at Matotschkin Shar, and a species of *Hyas*.

POLYZOA.

The Rev. Thomas Hincks has furnished the following list the species collected by Mr. Grant:—

Subclass HOLOBRANCHIA.

Group a. ECTOPROCTA.

Order GYMNOLÆMATA.

Suborder Cheilostomata.

Genus GEMELLARIA, Savigny.

Gemellaria loricata, L.

Very abundant off Bear Island, in 25 fms. A dark-coloured variety occurs, with much elongated internodes, in lat. 72° 32' 3" N., long. 36° 29' 5" E. [St. Lawrence, Dawson.]

Genus CELLULARIA, Pallas.

Cellularia Peachii, Busk.

Lat. 71° 23' N., long. 49° 38' E., 67 fms., off S.W. end of Novaya Zemlya. [St. Lawrence, Dawson.]

Genus MENIPEA, Lamouroux.

Menipea ternata, Ellis & Solander (normal form).

Extremely abundant, as in all northern dredgings.

Menipea ternata (forma *gracilis*, Smitt) = *M. gracilis*, Busk.

Lat. 72° 32' 3" N., long. 36° 29' 5" E., 128 fms. [F.-Pierce Bay, lat. 79° 29' N., last British Arctic expedition.]

Genus BUGULA, Oken.

Bugula Murrayana, Johnston.

On *Gemellaria loricata*, lat. 72° 32' N., long. 36° 29' 5" E., 128 fms. [St. Lawrence, Dawson; lat. 79° 29' N., last Brit. Arctic exped.]

Genus FLUSTRA, Linnæus.

Flustra membranaceo-truncata, Smitt.

An essentially Arctic form. In this species membranous processes are given off from the back of the cells, terminating in branched fibrils, by which the zoarium is attached. This structure, which is not shared by other members of the genus, points to some peculiarity in the conditions of life.

Genus MEMBRANIPORA, Blainville.

Membranipora monostachys, Busk.

On stones &c., lat. 74° N., long. 24° E., 220 fms.

Membranipora lineata, L.

On stones off Bear Island, 25 fms.

Membranipora craticula, Alder.

On stones off Bear Island, 25 fms. [St. Lawrence, *Dawson*.]

Membranipora Sophia, Busk.

On stones off Bear Island, in 25 fms.

Membranipora arctica, Smitt.

Extremely common. This is the most abundant species amongst the dredgings from Barents Sea. It occurs creeping over stones, shells, &c., and also erect and free, forming expansions composed of a single lamina or of two (*Hemeschara* or *Eschara* auctt.). In the *Hemescharine* state the dorsal surface of the zoecia is seen to be thickly covered with minute white disks. Lat. 74° N., long. 23° E., 220 fms.; and off Bear Island, 25 fms.

Genus MICROPORELLA, Hincks.

Microporella ciliata, Pallas.

On stones off Bear Island. The zoecia are much calcified and often coarsely grooved, the furrows extending from the margin to the umbo, which overhangs the pore. The latter is suborbicular and strongly dentate, and is placed in a hollow between the umbo and the lower lip. The avicularium is very generally wanting.

Genus PORINA, D'Orbigny.

Porina tubulosa, Norman.

On stones off Bear Island, 25 fms. [St. Lawrence, *Dawson*.]

Genus MYRIOZOOM.

Myriozoom subgracile, D'Orb.

Lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms. Lat. $75^{\circ} 36' 3''$ N., long. $57^{\circ} 6' 7''$ E. (*Marenzeller*).

Genus SCHIZOPORELLA, Hincks.

Schizoporella sinuosa, Busk.

On stones, lat. $72^{\circ} 55'$ N., long. $37^{\circ} 57' 18''$ E., in 150 fms. [St. Lawrence, *Dawson*.]

Schizoporella plana, Dawson (= *Myriozoom crustaceum*, Smitt).

Off Bear Island. [St. Lawrence, *Dawson*.]

Schizoporella hyalina, L.

On stones off Bear Island, 25 fms.

Genus PORELLA, Gray.

Porella struma, Norman.

Lat. $73^{\circ} 41' 6''$ N., long. $22^{\circ} 58' 30''$ E., in 220 fms.

The habit of growth in the only Barents-Sea specimen is Hemescharine; and the dorsal surface of the zoecia is furnished with one or more spinous projections, often of some length.

[North of the Shetland group; Bergen (*Norman*).]

Genus PHYLACTELLA, Hincks.

Phylactella? *grandis*, n. sp.

Lat. $75^{\circ} 16' 6''$ N., long. $48^{\circ} 19' 26''$ E., 160 fms.

Genus MUCRONELLA, Hincks.

Mucronella scutulata, Busk.

On stones, lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms.

Mucronella simplex, n. sp.

On stones, lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms.

Genus *ESCHARA*, auctt.

Eschara (*Flustra*) *solida*, Stimpson (= *Eschara palmata*, Sars, and *Escharella palmata*, Smitt).

Abundant in lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms. Lat. $79^{\circ} 13' 1''$ N., long. $63^{\circ} 21' 7''$ E. (*Marenzeller*). [Greenland; St. Lawrence.]

Eschara glabra, n. sp.

Lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms.

Genus *CELLEPORA*, Fabricius.

Cellepora, ?n. sp.

On *Eudendrium*, lat. $72^{\circ} 32' 3''$ N., long. $36^{\circ} 29' 5''$ E., 128 fms.

Suborder **Cyclostomata**.Genus *CRISIA*, Lamouroux.

Crisia eburneo-denticulata, Smitt.

Lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms.

Genus *DIASTOPORA*, Lamouroux.

Diastopora obelia, Johnston.

On *Eschara palmata* (*Flustra solida*, Stp.), lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms. [St. Lawrence, *Dawson*.]

Genus *HORNERA*, Lamouroux.

Hornera, ?sp.

Lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 36''$ E., 160 fms.

Genus *LICHENOPORA*, Defrance.

Lichenopora verrucaria, Fabricius.

On *Balanus*, off Bear Island, 25 fms.

Suborder **Ctenostomata**.Genus *ALCYONIDIUM*, Lamouroux.

Alcyonidium excavatum, n. sp.

Attached to tubes of Annelids, lat. $75^{\circ} 16' 6''$ N., long. $45^{\circ} 19' 5''$ E.

Genus ARACHNIDIUM, Hincks.

Arachnidium simplex, n. sp.On other Polyzoa, lat. $71^{\circ} 6' N.$, long. $50^{\circ} E.$, 62 fms., off Novaya Zemlya.

Genus BUSKIA, Alder.

Buskia nitens, Alder.On *Eudendrium*, lat. $72^{\circ} 32' 3'' N.$, long. $36^{\circ} 29' 5'' E.$

Group b. ENTOPROCTA.

Order PEDICELLINEA.

BARENTSIA, nov. gen.

Barentsia bulbosa, n. sp.Lat. $75^{\circ} 16' 6'' N.$, long. $45^{\circ} 19' 36'' E.$, 160 fms.

Genus LOXOSOMA, Keferstein.

Loxosoma singulare, Keferstein.On *Menipea*, lat. $71^{\circ} 6' N.$, long. $50^{\circ} E.$, off S.W. end of Novaya Zemlya, in 62 fms.

The foregoing list of Barents-Sea Polyzoa embraces thirty-two species, of which six appear to be new. It compares very favourably with the results in this department of the Arctic expedition under Sir G. Nares, by which only seventeen species were obtained.

Marenzeller mentions, in addition to the species in Mr. Grant's collection, the following seven species:—*Hornera lichenoides*, L.; *Alcyonidium gelatinosum*, L.; *Scrupocellaria inermis*, Norm.; *Membranipora Flemingii*, Busk; *Eschara cervicornis*, Pallas (probably = *Porella compressa*, Hincks); *Discopora coccinea*, Abildg., form *ventricosa*, Hass. (= *Mucronella ventricosa*, Hassall); *Cellepora ramulosa*, form *avicularis* (= *Cellepora avicularis*, Hincks).

BRACHIOPODA.

Brachiopoda were rarely met with in Barents Sea. Dr. van Lidth de Jeude mentions *Terebratula cranium* as having occurred off the north coast of Norway.

PTEROPODA.

Dr. van Lidth de Jeude records *Clio borealis* and *Limacina arctica*.

GASTROPODA.

The following species are mentioned by Dr. van Lidth de Jeude which were not found in Mr. Grant's collection:—*Siphonentalis affinis* and *Hero formosa*, Lovén, the latter in the Matotschkin Shar, where also other species of Nudibranchiata occurred.

CONCHIFERA.

Dr. van Lidth de Jeude mentions *Pecten islandicus*, *P. septemradiatus* and *P. abyssorum*?, *Cardium ciliatum*, and *Panopea norvegica* as occurring in Barents Sea.

On examining the list of animals collected by Mr. Grant it will be found that there are enumerated 153 species, namely—Protozoa 1, Spongia 4, Hydrozoa 13, Actinozoa 2, Echinodermata 16, Annelida 15, Gephyrea 1, Crustacea 24, Polyzoa 32, Brachiopoda 2, Mollusca 36, Pisces 7.

Considering the limited means at Mr. Grant's disposal, and the confined space on board the little vessel, in which fourteen persons were cooped up, it will probably be acknowledged that he has achieved satisfactory results by his efforts to secure specimens for the Exeter Museum.

XXXIII.—On new Hydroida and Polyzoa from Barents Sea.

By the Rev. THOMAS HINCKS, B.A., F.R.S.

[Plate XV.]

A FULL list of the Hydroida and Polyzoa obtained by Mr. W. J. A. Grant in the Arctic seas, during the expedition of the Dutch exploring-vessel the 'Willem Barents,' is included in Mr. D'Urban's report, published in the present Number of the 'Annals.' This paper will contain a detailed description of the new forms which occur in the collection.

Subkingdom CŒLEENTERATA.

Class HYDROZOA.

Order HYDROIDA.

Suborder THECAPHORA, Hincks.

SERTULARELLA, Gray.

Sertularella quadricornuta, n. sp. (Pl. XV. figs. 1, 1a.)

Stem almost straight or very slightly sinuated, irregularly