The regular slope of the plains of Northern India up to the Siwalik Hills, which rise suddenly from the flat ground, leads me to infer that the sea must have continued to reach at least as far as the foot of the Siwalik Hills for some time after their upheavement.

The rise of the Tibetan plain has not been caused by the granitic eruption of the line of snowy peaks. That the greenstone rocks, that abound in many parts of it, have equally not caused it, is proved by the peculiar nature of the valleys among the hills to the west of the lakes, which must have been laid out level under water; from which it is to be inferred that these eruptive rocks are older than the tertiary beds of the plain. The same thing is shown by the occurrence of worn pebbles of greenstone in the surface of the plain in the vicinity of some of the detached hills of that rock.

The former extension of the glaciers far beyond their present limits is a phenomenon that may be noticed almost everywhere in these mountains, and may give rise at first sight to an idea that there may here also have been some special period of cold corresponding to the glacial epoch of Europe. But it seems, I think, more probable that this is here only the result of a change of climate consequent on the upheaval of the great plains of Northern India.

The existence of ancient moraines on the tertiary plain of Tibet proves that the extension of the glaciers is post-tertiary. Now, if we conceive that after the rising of this plain to nearly its present elevation, the sea still continued to wash the foot of the Siwalik Hills, as I have already said that I considered likely, it is clear that the climate of the Himalaya would have been far more moist, and that the quantity of snow that fell on the highest parts of the mountains would have been greatly in excess of what now falls there, causing a great extension of the glaciers beyond the limits to which they have now receded.

5. *NOTICES of the GEOLOGY of the STRAITS OF SINGAPORE.* By J. R. LOGAN, Esq., F.G.S., M.R. Asiatic S., Corr. M. Ethnological Soc., and Batav. Soc. Arts and Sc.

[PL. XVIII.]

Introduction.—Under this name I shall describe not only the rocks in the Straits of Singapore and along its shores, but also all those comprehended within a rhomboidal space, about sixty miles long and thirty miles broad, of which the northern boundary is a line stretching E. by N. across the Malay Peninsula, from the Straits of Malacca to the China Sea, and touching the most northerly part of the Old Strait of Singapore; the southern boundary being a line in the same direction, touching the most southerly part of Tilo' Sumpat, on the north coast of Bintang, and stretching across that island and all those west of it to Phillip's Channel. The investigation of the geology of the Malay Peninsula could not begin at any other place so favourable for observation, for instead of the dense jungle which everywhere else

obstructs our progress and hides the nature of the ground, the whole breadth of the Peninsula is here depressed and broken up, exposing natural sections in various directions, which are preserved from the obliterating influence of the rapid decomposition and exuberant vegetation of the region by the strong currents which the China Sea, on the E., and the Bay of Bengal, on the W., impel through the depression. Of these littoral sections the best are those of the islets in the Strait of Singapore, and the points of Singapore Island, the S.E. promontory of the Peninsula, Bintang, and Batam, which advance into it, and, receiving the full force of the main current, are worn by more powerful waves than those which invade the sheltered portions of the Straits, such as the narrow channel which separates Singapore from the continent, and the shallow bays of Batam.

I. STRUCTURE AND COMPOSITION OF THE DISTRICT.

1. *General configuration.*—The configuration of the district presents several features worthy of remark. The land-portion of it is low, and is composed of hills generally in ramified systems, the direction of the principal branches forming a small angle with that of the geographical axis of the Peninsula (N.W. by N., S.E. by S.), that is, deviating about 20° or 25° to the W. of it, or from N.W. by W. to N.W. by $W.\frac{1}{2}W.$ * The mean height of the hills must be under 100 feet, the greater number falling short of that elevation, while only a few points attain to heights varying from 500 to 750 feet†. The

* This direction, however, agrees with that of the whole western side at least of Johōr, the Peninsular coast from the latitude of Pulo Lumut, a little to the north of Parcelar Hill, running S.E. by E. to its south-western extremity at Tanjung Burn, a distance of about 160 miles.

† The following are the heights of the principal eminences of the ranges in the neighbourhood of Singapore Town, above the level of low-water, at spring tides, as determined trigonometrically by J. T. Thomson, Esq.

	Feet.		Feet.
Lessuden	72·3	Lady Hill	108·0
Mount Erskine	88·3	Mount Victoria	100·8
Ansersabs Hill, or Mount Wallich	144·3	Mount Elizabeth	82·2
Mount Parsee or Palmer	119·3	Cairn Hill	113·2
Duxton	68·2	Emerald Hill	71·9
Craighall	65·0	Mount Emily	135·1
Guthrie's Hill	106·2	—— Caroline	108·1
Raeburn	45·2	—— Louisa	112·1
Everton	34·2	—— Sophia	108·3
Spottiswoode Park	73·5	Bukit Chermin	106·6
Line Hill	124·2	Hooden Field	70·2
Mount Farquhar	107·3	Saint James	54·3
Rosemary Hill	115·9	Monastery	72·5
Sisters Caroline & Catherine Hill	110·1	Mount D'Alguilar, south	80·6
Sri Menanti	81·5	—— ———, north	75·5
Mount Harriet	103·5	Bain's Hill	78·6
Nassim's Fine Hill	93·6	Briars	91·3
Broad Field	75·3	Swiss Cottage	80·5
Bonny Grass	85·5	Mount Helen	80·8
Annan Bank	80·5	Mount Arthur	80·0
Pavilion	108·3	Duncarn	75·5
Kiltiney House	94·4	Green Hill	87·8

[Perth

ranges and branches of hills are always narrow and separated by depressions, which form salt-water creeks at their mouths and flat alluvial valleys towards the interior. A subsidence of a few feet would submerge them and restore the original outline of the land,—a system of narrow, elongated, branching peninsulas, alternately swelling and contracting, the several ranges and branches always tending to a parallelism with the axis of the mainland. Singapore, for instance (excluding the western triangular portion, which is nearly separated from it by the inlets of the Pandan and the Kranji and forms a curious peninsula by itself), would then present five peninsulas running from N.W. by W., where they are connected, to the S.E. by E. for about six miles, when the four to the west would be cut off along a line nearly N. by E.; while the broadest and most easterly would continue its course in the direction of Tanah Merah Besar, sending off branches to the north-east and forming the bulk of the eastern division of the island*.

The general direction of the Strait is transverse to that of the hill-ranges, but forms an acute angle with it. In reality, however, the Strait consists of a succession of depressions, each of which nearly coincides in direction with the axes of elevation of the land. The most western of these, about 22 miles long (N.W. to S.E.) and 18 miles broad, stretches from the south-west promontory of the Peninsula to the northern islands of the Bulan Archipelago, and is bounded on the east by the western coast of Singapore and the chain of islets which stretch from its extremity across the Strait to the N.W. point of Batam. The second basin lies between the S.E. coast of Singapore and the N. coast of Batam, both of which it deeply penetrates; the former

	Feet.		Feet.
Perth Hill	82·2	Mount Faber	300·7
Kaynan's Hill.....	98·7	Hill in front of Mount Faber ...	265·4
Whampoa's Hill.....	93·0	Red Hill.....	112·0
Monks Hill.....	78·4	Ulong's Hill	129·3
Draycott.....	84·2	Mount Zion	45·4
Sheriff's Hill, Sri Minanti.....	82·2	Peak Island	101·4
Sri Tanglin	86·5	St. John's	189·8
Mount Mary	102·7	Blakang Mati little Hill	202·9
Angus Back Hill	129·4	Pulo Braui Hill	161·5
Angus House	69·4	Bukit Timah	519·1
Pearl's Hill.....	170·9	Hill behind Green Hill	124·5
Institution Hill	121·9	Claymore Chinese Burying	
Government Hill	156·5	Ground	74·5
Blakang Mati large Hill, or Bukit		Hannifasah's House	71·2
Serapong	301·7		

The elevations of some other hills have been determined also by Mr. Thomson:—

	Feet.
On the mainland—Gugong Bau	749
Marubukit.....	661
Bukit Santi	645
On Pulo Bentan (Bintang Island)—Gunong Bentan...	1212
Bukit Kijang.....	759

* The map shows the configuration of this part of the island well. All the portions represented there as alluvial are flat, and have been silted up in recent periods.

in the alluvial plain forming the districts of Rochor, Gelang, Kalang, Paya Lebar, and Siglap, with its long branches or inlets, extending between the hill-ranges to the middle of the island; and the latter in Batam and Bulan Bays*. Its greatest length, from the extremity of Batam Bay to that of the Singapore Plain, is about 15 miles†; its least, from Tanjong Siglap to T. Singkwang, about $8\frac{3}{4}$ miles; its breadth is nearly 10 miles. The third basin is the middle of a prolonged submarine valley which stretches northward into the Peninsula, where it forms the estuary of the River Johör; on the south it penetrates the Johör Archipelago, where it forms the Strait of Rhio separating Batam from Bintang. The fourth or most easterly basin, —about 18 miles long, 20 miles broad at its two extremities, and 14 miles in its middle,—lies between the south coast of the S.E. promontory of the Peninsula and the N. coast of Bintang.

2. *Geological Topography.*—In giving a general sketch of the geology we shall consider the District as divided into four zones nearly corresponding with the four basins of the Strait.

1st zone.—The first, or most westerly basin, has on its southern side a deep channel, free from islands, which opens into the second basin between Pulo Sambo Kichi and Pulo Sikijang. To the south of this channel a dense clump of islands commences, which stretches to the S.E., and includes Bulan with its conspicuous peak, from which the Dutch hydrographers have named the group the Bulan Archipelago. It remains to this day unsurveyed and unexplored either by the Dutch or English, and in some charts still figures as one island, although it must consist of several scores‡. It is separated by a narrow channel, called Silat Batu Haji, from the west coast of Batam, which has a general direction coinciding with that of this basin. The northern part of the basin is shallow and contains numerous islands; and the same configuration is continued in the western section of Singapore, which is a congeries of narrow peninsulas, separated by mangrove creeks. The prolongation of the zone in the mainland, including the S.W. promontory, is flat and alluvial towards the W., and hilly towards the E.

The whole of this zone consists of stratified sedimentary rocks, sandstone appearing to predominate towards the west, and sandstone and conglomerate, intermixed with clay and shale, towards the middle. Towards the eastern side a broad band of shale, chiefly chocolate-coloured, occurs; and in the extreme east, where it abuts on the syenite of the 2nd zone, this shale is interstratified with sandstone and conglomerate.

The western portion is nearly all covered by the sea or alluvium, the only islands, except those belonging to Bulan Archipelago, being Pulo Rungkum (Alligator I.), Pulo Sinang (Barn I.), Situmbu Kichi,

* These bays are deeper than they appear on the map, and the numerous creeks and inlets which pierce the island between the hill-ranges are not laid down, no survey of the bays having been yet made.

† About twenty-one miles, if we include the valleys which have been inlets of the sea since the land attained its present level.

‡ I have examined the northern part, where I found a perfect labyrinth of islets.

and Bésar (Rabbit and Coney I.), all of which appear to consist of sandstone and shale.

In the next parallel to the east we find in Pulo Salu strata of clay-stone and chert, highly indurated, crystalline, fissile, bluish, greenish, and greyish; soft clay; quartz-veins; quartzo-ferruginous walls; a subcrystalline grit of a granitiform aspect and in some places with a greenish base;—between Salu and P. Sudong, reefs of iron-rock*;—in P. Kapal, fine, purplish clay; steatitic clay, indurated, white, with seams and walls of iron-rock; fine shale, indurated, purple;—in P. Piel Ayëm, soap-stone of variable purity, indurated, coarsely foliated; hard steatite; iron-rock in seams and veins; the steatite at some places containing iron-pyrites, and locally iron-masked and cellular;—in P. Blakang Padang, coarse conglomerate, some of the pebbles of which are steatite, either pure or imbedding small quartz-grains; finer beds with much steatitic clay, generally of a light greenish colour, occasionally violet and red; broad beds of fine-grained sandy marl.

Returning to the north, we have in P. Marambong clay with quartz and iron-rock remarkably developed;—in the north-west division of Singapore at Tanjong Gul, clay, fine (as at Marambong), with quartzo-ferruginous walls; at P. Sambulan and Tanjong Kling, clays traversed by ferruginous and jaspideous walls; much laterite-gravel;—at P. Damar laterite soil, gravel, and fragments; the coast of Singapore facing it is lateritic; on the west bank of the mouth of the Jurong clays and sandy marl; iron-rock swelling up through it in mammillary and botryoidal bosses; ferruginous and jaspideous walls; large laterite blocks composed of small pebbles and fragments cemented together.

Following this band of strata to the S.E., we observe in the Sikra Islands clays chiefly, with ferruginous walls; in Sikra Laut numerous highly indurated, ramifying, mammillated, iron-walls; radiating and concentric laminæ of colour; thin beds of bluish-black and greenish clays. These rocks are continued across to the Old Strait. In P. Busong we find indurated sandstones and clays with irregular ferruginous walls; reticulations of quartz-veins; much disturbance of stratification;—in P. Bukum, clay with indurated walls; light-chocolate-coloured clay; laminæ of colour;—in P. Simakan and P. Siking, iron-rock; hard laterite;—in P. Jong, clay, white with a faint green tinge; a few conglomerate-beds; numerous iron-walls; concentric and radiating laminæ of colour highly developed;—in P. Sabaru conglomerate, sandstone, clay; numerous iron-dykes and laminæ of colour as in P. Jong.

The next parallel is composed chiefly of sandstone and conglomerate, but with intervening beds of clay, mostly chocolate and purplish. At particular places much iron-rock occurs, and ferruginous lamination is frequent, although not on the great scale of the Sabaru group. The principal exhibition of this band is in the high range of hills on

* This term is used for brevity in this general sketch. The most common form is the hydrous peroxide of iron; and the iron-rock throughout the district, I shall afterwards show, is the common rock of each particular situs disguised by a greater or less injection of iron.

the S.W. coast of Singapore, stretching from Suŋge Pandan to Batu Blyër (where the highest sea-cliffs in the district occur), and continued in the island of Blakang Mati. Proceeding along the same line of elevation to the S.E., we find it again in P. Sikukur, the two Sikijangs and Subar. It reappears on the other side of the Strait, and on the same line, in the two Sambos (where we find sandstone, grit, and conglomerate, for the most part highly indurated and semi-crystalline; clay, chocolate, light purplish, violet, greenish, and yellow; and iron-rock), P. Mirlam and the eastern portion of the N.W. point of Batam.

To the east of this line the chocolate and violet clays, which have hitherto either shown themselves as isolated beds or in narrow bands amongst the other strata, become continuous, occupying a breadth of about one mile and stretching right across Singapore in the direction of N.W. by W. This band includes P. Brani and the eastern portion of Blakang Mati, in both of which the colours vary much,—chocolate, purplish, different tints of violet, greyish, lilac, green, and in which sandstone-layers are intercalated. It is then lost for about six miles in the Strait, but is found again on the east side of Sambo Kichi and on the coast of Batam, at Pulo and Tanjong Daŋgas.

2nd zone.—The second zone consists almost wholly of soft amorphous clay and sandstone with conglomerate; the latter, however, containing a good deal of clay either mixed with it or in separate beds. The clay in Singapore forms a compact tract of about sixty square miles, occupying the main body of the island*. On the south it is separated from the plain of Singapore by a narrow margin of sandstone, and then stretches across the island in a N. by E. direction to the Old Strait. On the west its boundary runs in a north-westerly line, nearly parallel to the chocolate clay band, for about eight miles, when it is deflected towards the N. in the direction of S. Kranji, in the vicinity of which it meets the Old Strait. In Bukit Timah, solid greenish granite and syenite, passing in some places into compact laminated felspar, are seen. On some of the other hills of the clay-tract, protruding blocks of syenite and granite are occasionally found. Where deep sections have been made through the clay, its structure and composition are found precisely to resemble that of adjacent blocks, while on the surface it has a remarkable uniformity of character†, and is easily distinguished from the sedimentary clays. The whole of this tract, therefore, is decomposed plutonic rock. The clay in some places, and particularly in the bottoms of valleys, where it is covered by vegetable mud and has undergone a natural bleaching, is a pure white kaolin, but it generally imbeds quartz-granules (in rare cases abundantly), and a few feet below the surface is mottled

* It is almost a square of eight miles, coinciding with the Peninsular axis; that is, two of its sides run N.W. by W. (the northern of the two giving its direction to the Old Strait from T. Pongal to S. Tambrau), and the other two sides running N.E. by N. (the northern of the two giving its direction to the Old Strait from S. Sinoko to the creek of S. Batang Hari, and the southern giving its direction to the ancient W. coast of Singapore Bay).

† The variable proportions of iron and quartz, however, affect this character.

with red in various degrees, in isolated or connected ramifying blotches or in parallel curvilinear streaks. Fresh sections exhibit various tints of red, lilac, purple, yellow, &c. Pure white streaks and zones also occur frequently, exhibiting a kind of irregular, broken reticulation. It is sometimes altogether of a dark red hue. When this is the case, masses of a half-decomposed iron-rock are generally found in it; and in most localities, where cuttings have been made, a similar rock occurs here and there in dykes and ramifying veins. On the sides of these veins laminae of a red colour are sometimes seen, either parallel to the vein, radiating from points in it, or forming systems of concentric lines. The iron-rock is also found either strewn over the surface in amorphous cellular blocks and small pebbles, or below the surface, at depths varying from a few inches to a few feet, in layers of similar pebbles. With these pebbles others of jaspideous and porcellaneous rock are frequently intermixed, and veins or strings of the latter are found in the deeper sections or in the half-decomposed crumbling granite which in a few localities is exposed at the surface. In some places patches of a vesicular jaspideous rock are frequent in the decomposed mass. These are sometimes considerably ferruginous. Fragments of semi-decomposed syenite and granite, ferruginous in different degrees, are common. When highly ferruginous, they are more or less vesicular, and the only unchanged constituent of the solid rock is the quartz. Even this is sometimes penetrated by the iron. The upper soil, or that which has been completely subjected to atmospherical action, is a clay with a greasy lustre, but often inclining to a dry friable appearance, in colour generally yellowish brown, but with lighter and darker hues in particular localities. In many hills it is highly ferruginous and of a deep red colour. This, with the smaller proportion of quartz, distinguishes it from the granitic clays of Pinang higher up on the west coast.

The easily accessible localities where ferruginous blocks are most exposed are the hills along the boundary between the plutonic and sedimentary tracts near Singapore Town, such as Mount Victoria (58)*, Sri Menanti (39), the hills connecting it with Mount Zion (38), the hill forming the extreme S.W. point of the granitic tract (33), the western hill of Dr. Oxley's nutmeg-plantation (36), Cairn Hill (50), the coast of the mainland from Tanjong Tanguloh to Pulo Nanas (at Tanjong Passier Mera and near S. Tukong), &c.

The sandstone and shale surround the plutonic tract on the W. and S., being found occupying the space to the N.W. of it up to the Old Strait, rising in the hills near Singapore Town to the S.E. of it, and, lastly, forming, with a great predominance of sandstone, grit, and conglomerate, all the eastern portion of the island, with the exception of the hills at Chaŋggèi. At several places there is much ferruginous rock, as in the hills running south from the Mosque in the town to Tanjong Malang, and continued in the reefs off that point. On the opposite side of the flat of the Singapore River, and where the very same line, prolonged to the north, crosses the two bifurcations of the Tanglin Range, we find a considerable development

* The figures refer to those on the map.

of laterite-rock in Government Hill (35), Mount Sophia (48), and Mount Seligi (47), where the clay is intercalated amongst the sandstone, which latter is highly indurated.

On the Batam side of the Strait the sandstone is found from Suñge Ledi in Batam Bay to Bulan Bay, and at some places contains iron-rock. The east side of Bulan Bay and the whole N.E. promontory of Batam with Pulo Nongsa is granite, which differs from that of Singapore in colour and in being larger-grained.

3rd zone.—On the east coast of Batam, in contact with the granite, are found iron-rock and clay, and, in the granite, iron-dykes containing casts of shells. The third zone includes the island of Pulo Ubin, consisting of various forms of granite and syenite (often approaching to and sometimes passing into greenstone) and remarkable for its tabular and concentrically laminar structure and its extraordinary grooves, resulting from the action of decomposition on its rocks;—the eastern promontory of Singapore has a similar composition; Pulo Nanas or Kirimkin, and the mainland opposite are composed of greenstone-porphry; the hills on the mainland at Tanjong Tanguloh of decomposed plutonic rock, similar to the prevailing Singapore clay, and, like it, containing ferruginous and jaspideous fragments; the small islets of P. Sijahat and P. Hantu III, consisting of foliated claystone and chert, varying from subcrystalline to highly crystalline, generally very hornblendic, and traversed in P. Sijahat by a dyke of greenstone-porphry; Pulo Tikong Besar and Pulo Tikong Kichi are chiefly sandstone, highly indurated, and with iron-rock; in the sea off the S.W. point of Tikong Besar rocks of granite occur, called Batu Kapala Tua*; Marbukit (Johor Hill) is composed of clays and conglomerates, indurated, passing into crystalline chert, &c., and of quartz and iron-rock;—the coast of Bintang, from Tanjong Kalumpang to Tanjong Subong, is probably granitic.

4th zone.—In the fourth zone we find on the east bank of the Johor River, at Johor Lama†, granite; to the south of it, grit and conglomerate, from Tanjong Iladong to Tana Runto (in the conglomerate of which fragments, partially decomposed, of chloritic granite, some very little rounded, are found), and probably as far as Tanjong Boe; in Gunong Bäu and the lower hills behind it on the S. Karang, greenstone, varying from compact to granular, the latter approaching the fine-grained granites of Pulo Ubin; along the south coast of the S.E. promontory of the Peninsula at Tanjong Tiram greenstone-porphry; at T. Telumpang, granite; at Tanjong and Pulo Rumnia, shales, further to the E. passing into subcrystalline rocks; at T. Penyusoh (Point Romania), Peak Rock, South Island, and Pedro Branco (not included in the map), syenite; North Rock, shale; along the east coast, facing the China Sea, at Tanjong Pungai and Tanjong Kinawar, talcose sandstone and shale, sometimes semi-crystallized, with abundance of iron-rock in large rounded shining blocks, and quartz‡;—

* J. T. Thomson, Esq.

† This Bugis town is further north than the map extends.

‡ See my "Notices of the Geology of the East Coast of Johore." *Journ. Ind. Arch.* vol. ii. p. 625.

at Tanjong Lompatan, talcose shale, quartz, and iron-rock;—along the north coast of Bintang, from Tanjong Subong to the N.E. Point, syenite*; at the Point, talcose clay and shale with iron-rock†, and quartz (similar to T. Kinawar)‡.

3. *Character and Extent of each kind of Rock.*

1. *The Sedimentary Rocks.*—Viewing the district as a whole, we observe that amongst the sedimentary rocks the shales greatly predominate, the chocolate and purplish being the most common. The soft shales and clays are of far greater extent than the indurated, and are either quite soft, yielding to the hand, or have various degrees of consistency, up to so much hardness and toughness as to require the hammer to break them. They are in great measure a mixture of clay and fine siliceous granules, to which the term *marl* may be applied, but as descriptive of their consistency only, for none of them are calcareous. Those in which clay is in excess, so as to be completely plastic, are not frequent. The steatite, when it occurs, is generally in large proportion, but of limited extent. I have only observed completely indurated argillaceous and argillo-siliceous rocks of a cherty and jaspery nature, and considerable in extent, at Pulo Salu, P. Kapal, P. Piel Ayem, Tanjong Putri (western), Pulo Sijahat, Pulo Hantu III, and Tanjong Pinrang §.

The sandstones and conglomerates are more often indurated than friable. In many cases the former have a subcrystalline texture, and the latter a compact siliceous basis, traversed by veins of quartz-crystals. The most numerous pebbles in the conglomerates are of a dull whitish quartz. Greyish quartz-pebbles are also common. Black, red, and yellowish pebbles occur sparingly, although in some localities more abundantly than in others. Felspathic pebbles, sometimes with a crystalline, but generally with an earthy fracture, are common, violet tints prevailing. The basis of the conglomerates, when not indurated, varies from clay, often chloritic or steatitic, to a friable earth or marl, sometimes also chloritic, containing an abundance of quartz-granules. The highly indurated and subcrystalline sandstones chiefly occur in Mount Palmer (24), Government Hill (34), Mount Sophia (48), and Mount Liligi in Singapore, and in the promontory terminating at Tanjong Singwang in Batam.

2. *The Plutonic Rocks.*—Taking a similar broad view of the pluto-

* J. T. Thomson, Esq.

† Mr. Thomson says, that "Laterite in a fused state appears to have run into and over this stratum in several places so as to crack it."

The lava-like appearance of the hydrous peroxide of iron here and in numerous other localities is exceedingly deceptive. I was misled by it for a considerable period, and the views which I have communicated to the Asiatic Society respecting the geology of this region were more or less coloured by this notion. The ejected-fluid-aspect and blackish colours are entirely owing to the peroxidation and concreting process of the iron. Is not the iron-dyke in Mavrosplia, which puzzled M. Virlet d'Aoust (*Expédition Scientifique de Morée*, pp. 54, 55), capable of a similar explanation?

‡ Specimens from J. T. Thomson, Esq.

§ P. Pisang (N.W. of P. Kocob, beyond the limits of the map) and the east coast of Krimun Kichi are striking examples of this class of rocks.

nic portions of the district, the most general character that we observe is the predominance of felspar and the prevalence of hornblende,—quartz and mica being generally in small proportions, and the latter rare. Well-developed granite of the common type is found in very few localities on the northern side of the Straits, the plutonic tract of Singapore being very deficient in quartz and mica, most of the red clay hills which occur in it being probably decomposed greenstones and highly hornblendic syenites. Even Pulo Ubin, although it presents several decidedly granitic bosses, tends, on the whole, more to the felspatho-hornblendic character, which exclusively prevails to the eastward of it, in Pulo Kirimkin (Nanas), the mainland opposite, Pulo Sijahat, Gunong Bau, and the adjacent hills. The only considerable display of proper granite is that on the southern side of the Strait, in the north-east section of Batam and the north of Bintang*.

With respect to the granulation and composition of the plutonic rocks, every variety may be found, from the most compact hornblende and felspar, through granular greenstones, minutely grained grey-stone, syenite, and eurite, to large-grained granite. In Pulo Ubin we find the following gradations (the hornblende being in general either in nests of small granules mixed with felspar, disseminated in minute particles, or in long thin streaks): light reddish granite, with quartz, predominating; granite of a bluish grey tinge; syenite with felspar in excess; syenitic greenstone; greystone; grey felspar, transparent quartz with some black mica interspersed, and nuclei with the mica in excess; a quartzo-felspathic basis (pegmatite) with minute particles of mica and hornblende disseminated, in some places sparingly, in others abundantly, so as, on partial decomposition, to give the rock the appearance of finely-ground pepper; spherical nuclei of mica with a little felspar, surrounded by concentric laminæ; greenish grey saccharoid felspar, with crystals of quartz and nests of mica sparingly disseminated; whitish felspar, black mica, and hornblende; well-crystallized felspar, with dark green hornblende disseminated in granules, cloudy spots, and fibres; blackish greenstone or hornblende rock; felspar and hornblende confusedly aggregated; opaque white felspar, tinged green, and blackish green hornblende in very large and distinct granules; very finely grained blackish greenstone approaching to basalt; basalt; opaque bluish grey and white saccharoid felspathic and quartzo-felspathic bases, inclosing crystals and grains of hornblende and translucent crystals of felspar (closely resembling one variety of Vesuvian lava), in some places becoming compact, the hornblende granules sometimes so minute as to appear like fine black dust sprinkled on snow; in the group of rocks close to Ubin called S'kodo (from the supposed resemblance of one of the most conspicuous to a frog), there is found a regular large granitic crystallization, the hornblende being in well-defined crystals of various sizes and mostly fibrous.

3. *The Ferruginous Rocks.*—The iron-rock and quartz which are found so generally, and in particular localities so strongly developed, occur occasionally in broad irregular patches and bands, but usually

* I have not personally examined Bintang.

in walls, veins, and seams. Where they are most abundant the principal walls follow the direction of the beds, but are most frequently sinuous, and connected by branches with those adjacent. The beds themselves, in such cases, are generally bent in a similar direction. The sides of the beds, where not obliterated by these walls, are often indurated and acquire a shining mammillated surface. Sandstones are often divided into little cubical and rhomboidal compartments, being pervaded by a complete reticulation of a thin black ferruginous crust or wall, which, from the washing-out of the sandstone by the action of the waves, projects, like the sepiments of a honey-comb, from the surface.

In clays and shales (and sometimes in fine-grained sandstones) red-coloured laminae are frequently seen in connection with the ferruginous walls and veins, and arranged either in lines radiating from points or in concentric curvilinear figures. The most remarkable examples of this laminar coloration occur in the islands of Jong, Sabaru, and Bukum. Another kind of lamination is often seen in ferruginous walls. In this the original laminae of the rock are preserved, but converted into iron-ore. Sometimes they are swollen into a globular form. This seems to be the origin of most of the mammillated and botryoidal forms.

The associated iron and quartz are found developed on the largest scale at Marambong and Tanjong Kinawar; and more or less quartz is found in almost every highly-ferruginous wall.

The localities where the iron predominates over the quartz are very numerous. Amongst the more striking instances may be mentioned, in sandstones, T. Sirimbun, Mount Palmer, the coast of P. Tikong Bësar; and in shales and clays, P. Dammar, the coast of Singapore opposite, Tanjong Pénjuru, P. Siking, P. Sabaru, the lateritic portion of Government Hill, the north-east part of P. Tikong Kichi, and Tanjong Pungäi.

The iron-rock is generally a hydrous peroxide and different forms of brown hæmatite, compact and ochry. In most localities the proportion of iron varies much, and is often too small to entitle the rock to be called an ore, but patches and masses of ore occur abundantly, and some walls are almost wholly composed of it. Regularly crystallized ores are not frequent in considerable patches, but hæmatitic crystals occur in the body of pieces of compact ore; and the thin brownish-black and black ore penetrating ferruginated rocks in seams, forming the external walls or crust of mammillary and botryoidal, and the lining of tubular and cavernous, masses of laterite and iron-masked rocks, is sometimes composed of fine fibrous crystals. I have found large-crystallized hæmatite in ferruginous sandstone and shale near the boundary of the plutonic and sedimentary tracts to the westward of Bukit Timah, and in violet-coloured shale midway between the Chinese Tokong, on the Pandan, and Bukit Timah.

We have already mentioned the mode in which ferruginous rock occurs in the plutonic tract. The following is an analysis, made in England, of a specimen of the common ore of the dykes occurring in this tract.

Peroxide of iron (containing 56 parts of metallic iron)	80
Silica, lime, and alumina	5
Water	15

Sp. gr. 3.12.

Pyrites occur very rarely, as I have only observed them in a considerable nidus amongst the shale of Pearl's Hill in a mass of anthracite, in steatite at P. Piel Ayem, and in the granite of Pulo Ubin.

4. *Alluvium*.—All the valleys are filled up to a greater or less height with alluvium, the bulk of which has been deposited from salt water, the freshwater deposits being generally superficial, although in some places of considerable depth. As this formation, in all its varieties, is still going on before our eyes, I will here mention causes and effects together. It has been remarked, that, when the district attained its present level, the sea must have flowed up the numerous long inlets which everywhere penetrated the land. All the different rock-formations must then have been accessible to the sea. Let us revert to this period in Singapore, where we can easily trace our way back to it. At that time, in crossing from Batu Blyer to Tanjong Tangeloh eight arms of the sea had to be passed, each with its shores winding round the points and into the concave recesses of the ramifying ranges which separated them, and receiving from the multitude of hillocks the clay which every shower of rain removed from their surfaces. The whole land was so constructed as to allow the greatest amount of waste to take place under ordinary atmospherical erosion. Scarcely any flat ground existed. Everywhere acclivities rose from the sea only to attain a certain height and be again bent into declivities which sunk into the sea on the other sides. The upper part of the valleys filled rapidly with clay, mud-flats were formed there and along the margins of the inlets, on which the mangrove and other salt-water trees flourished, and by their decay gave a carbonaceous character to the mud. In the shade of these maritime forests the *kapiting* and *udang kiteh** built up their pyramids of mud and helped to elevate and consolidate the floor, while their excrements and remains, with those of numerous crabs and testaceous mollusks, contributed with the vegetable matter to give a new character to the clay. In the meantime the same process went on in the eastern part of the island, but the sediment there was more mixed with sand. The points at the mouths of the inlets along the whole northern coast of the second basin of the Straits then consisted of sedimentary strata, exposed to the action of waves and currents, and those in the east, in particular, formed of yielding sandstone, supplied a large quantity of alluvial matter, while the shales of the hills near the site of Singapore town freely gave their loose red soil to the sea. When the mangrove-marshes, with their mud-flats in front, had gradually approached the mouths of the narrow inlets, a new condition of things arose. The sand, no longer sinking into deep water beyond the influence of the powerful surface-agitation of the sea, was thrown upon the mud-flats,

* Burrowing crabs. The *udang kiteh* is shaped like a scorpion, and greatly resembles, if it be not identical with, the *Thalassina scorpionoides*.

and, during the north-east monsoons, the heavy waves from the China Sea piled it up in long belts in front of the mangroves.

The clayey sediment of the plutonic tract now accumulated more exclusively around the mangroves within these belts. As the sands extended, the waves no longer penetrated into the inlets. In the higher part of the valleys, and in the recesses of the lower parts, the water became brackish, the mangrove and its associates died out, and a new and more exuberant order of vegetation, in which palms and ferns abounded, took their places. A black peaty matter now formed rapidly, filling up all the hollows, spreading over the ancient mud-flats, and burying the remains of the mangrove-forests. The alluvium now advancing beyond the inlets into the gradually shoaling bay of Singapore, the three processes which we have described proceeded for the future simultaneously. One sand-bank after another was thrown up, and the streams from the valleys were deflected into lagoons between them, in which a succession of events analogous to that of the old inlets took place. In the upper part of each successive lagoon were brackish marshes, in which dense forests grew; and in the lower, and along the margins of the creeks as far as the water was salt, the mangroves kept their place, forming, with the sand, a barrier against the waves. Bank after bank arose, lagoon after lagoon was formed, until the whole of the ancient bay was converted into a marsh, covered with forest and containing a succession of dry banks of sand or *pérmatangs*, the last of which has formed the present lagoon of the Gelang, projecting its extremity, like a tongue, into the harbour of Singapore at Tanjong Ru (Sandy Point), and has given a common estuary to the streams from every valley that opens into the Singapore plain.

This history we narrate with as much confidence as if there had been human witnesses and chroniclers of its whole progress, because not only has it left its own records of all its stages, but its current events are repetitions of these, and afford the fullest explanation of each of them*.

The valleys have now perfectly level surfaces, covered with forest where plantations have not been formed; and the layers of which they are composed are the pure plutonic clay,—a dark bluish marine clay which is seen at the heads of the valleys on both sides of the island up to the foot of Bukit Timah,—a black, soft, vegetable peat, or rather mud, in some places of great depth where hollows have

* “The lower parts of the valleys are mostly swampy, consisting of sand, clay, and black peaty mud; of the latter there are considerable tracts constantly moist and exhibiting an extraordinary rankness of vegetation. Looking on one of these swamps covered with tall but slender trees, and dense underwood growing up rapidly, and from the looseness of the deep bed of black vegetable matter,—the accumulated remains of their short-lived predecessors—destined soon to fall in their turn, and considering the deposits of clay and sand which accompany and give rise to it, it is impossible to doubt that we see nature repeating the precise process by which the materials of most of the ancient carboniferous strata were brought together. Towards the sea these forest marshes give place to mangrove swamps.”—*On the Local and Relative Geology of Singapore, including Notices of Sumatra*, &c. (by J. R. Logan, Esq.), Journ. Beng. Asiatic Society, vol. xvi. p. 525.

existed,—and sand. The plain consists of the same matter, sand and vegetable mud prevailing towards the surface in all the eastern part, where alone a considerable extent of either is found. The inner part of the plain, before it was cleared and partially drained, formed there a large marsh called Paya Lebar (broad marsh) with a deep bed of black vegetable mud in which a dense forest grew. The outer part, towards the range of sandstone hills forming the eastern margin of the plain, is a sandy flat, which, as it extends to the west, divides into long p̄rmatangs, with flats of clay and vegetable matter between them. The upper or eastern part of these flats often contains a considerable proportion of sand. The more inland banks continue across the inner part of the plain, making the first continuous coasts of the bay. Others, deflected in different directions, but generally trending more to the west and to the south of west, traverse the plain; the penultimate one extending along the Chaṅgie road up to Kampong Glam, and the latest forming the present coast terminating at Tanjong Ru. Although the force of the north-west monsoon has laid down the sandy debris of the eastern cliffs in these long p̄rmatangs extending to the western side of the plain, deflected all the eastern streams in the same direction, and is even bending to the west the mouths of the western streams themselves, and driving them all up into the extreme S.W. corner, where there will probably be ultimately but one outlet for all the waters that fall into the harbour, yet there are records of a time when the western side of the plain and the western inlets had their own p̄rmatangs. One of the most distinctly marked runs transversely to these inlets, or from S.W. to N.E., and it must have been formed while the greater part of the plain traversed by the p̄rmatangs from the east was covered by the sea. The only point where the Siglap sandstone-ranges are now wasting is at the Red Cliffs, but the hills stretching to the west of this for about a mile and a half, and gradually retiring from the present line of coast, are sharply scarped at all their points which project on the sandy plain, and the scarping of those forming the S.W. angle of the range was probably coæval with the formation of the p̄rmatang in question*.

Shells are found, often very abundantly, in the alluvial clay. Beds of Coral are also occasionally met with. The infrequency of deep excavations is probably the reason why, in localities inhabited from very ancient times†, human remains are so seldom brought to light. In the inner part of the Singapore plain a piece of coir-rope was found 6 feet below the surface, and a piece of wood bored through, and having the hole filled with the twisted fibres of a piece of rope, was found in the town of Singapore at a depth of 40 feet. Mr. Thomson bored through 10 feet of blue mud and 30 feet of ferruginous

* The p̄rmatangs of Singapore are inferior both in length and elevation to those of the large plain of Province Wellesley, facing Pinang, which have been formed by the waves of the Bay of Bengal. For further information on this subject consult the section headed *Rivers and Alluvial Formations* in my "Sketch of the Physical Geography and Geology of the Malay Peninsula," Journ. Ind. Arch. vol. ii. pp. 116 to 136.

† Journ. Ind. Arch. vol. i. p. 300, 302, and vol. ii. p. 517.

earth before this piece of wood was brought up. Other 40 feet of hard ferruginous earths and clays were pierced at this locality (near the Hindu Temple).

At the Gaol, 55 feet of blue mud, mixed with shells, were found resting on red earth mixed with the bark of trees.

At Teluk Ayer, near the beach, the layers bored through were,—

Soft blue mud with shells	40 feet.
Bright red clays alternating with brown and yellow clays	60 "

At the foot of Government Hill, near the Convict Lines, there were found,—

White clays	20 feet.
Red clays and laterite resting on a very hard white clay	30 "

In digging "Jocksing's Well," 9 feet of alluvial sand were found to rest on ferruginous gravel similar to that which occurs frequently on the hills near Singapore*.

We have not mentioned Coral as contributing to the alluvium of Singapore plain, because there is no evidence that it did so in a considerable degree. The eastern cliffs wasted too rapidly to allow of its growth in front of them, and mud and sand-banks, both unfavourable to the Polyps, were constantly increasing within the bay. The fresh water which it received from the interior would also be hostile to them. We shall at another place notice the localities where they flourish and the office they perform.

4. *Structure and Disposition in the Mass of the Rocks*†

1. *Sedimentary rocks*.—With the exception of the sandstones in the eastern part of Singapore and those on the east bank of the River Johor below Johor Lama, which are nearly horizontal, the strata rise at various angles, generally high, and frequently vertical.

Their strike over the whole district approximates to the direction of the Malay Peninsula, or N.W. by N. It deviates, however, about $22\frac{1}{2}^{\circ}$ on both sides, or N. by W. and N.W. by W., and this not merely at places distant from each other, but in limited localities.

Strata are very rarely found lying in other quarters of the compass, and such aberrations never extend far. The most striking exceptions are Kiltiney, adjoining Government Hill in Singapore, where the strata run N.N.E., as they also appear to do in the next hill to the west, Institution Hill.

* The above paragraph is from the paper cited in the preceding note. It is added, on the authority of Dutch writers, that in Banka, which is a part of the Peninsular band of elevation, and where tin pits are often dug to a considerable depth through the alluvium, there has been found a bed of turf, with wood which had been cut, below 20 feet of clay; a boat, differing from the kinds which have been in use from time immemorial, at a depth of 16 feet; and traces of a rice-field in another deep pit.

† I have obtained sections and measurements of the strata of many of the islands, but I reserve these for notices of these islands.

Amongst the most continuous lines that I have observed are that of Bukum, Jong, and Sabaru, nearly N.W. $\frac{1}{2}$ W. throughout; the Batu Blyer sandstone-cliffs along a considerable portion of their length; some of the shale-beds extending from Tiloh Blangah and Pulo Brani to Blakang Mati; the most southerly beds of Sikukur which are continued in the West Sikijang. But the greater number of the islands and points, where the stratification can be observed, present irregularities. Thus, to notice the most marked instances, proceeding from north to south, at Marambong we not only find the more ordinary oscillations, such as N.N.W., N.W. $\frac{1}{2}$ N., &c., but bends and curves, so that at some places the beds run N.E. by E., N.N.E., &c. At Busong similar irregularities occur, as we see N.E., N.E. by N., N.N.E., N., N. $\frac{1}{2}$ W., N.N.W. $\frac{1}{2}$ W. At Tanjong Lompatan the strike is N.E., E. by S., &c. The more common variations are such as those of Batu Blyer Cliffs, N.N.W., N.W. by N., N.W.; Blakang Mati, N.N.W., N.W. $\frac{1}{2}$ W., N.W. by W.; P. Brani, N.N.W., N.W. by N., N.W.; the hills east of the village of Tiloh Blangah, N. by W., N.N.W., N.W. by N.; West Sikijang, N. by N.W., N.N.W., N.W. by N.; East Sikijang, N. by N.W., N.W. $\frac{1}{2}$ W., W. by N.; Tanjong Piñger, N. by W., N.N.W., N.W. by N.; N.W., N.W. by W. at Tanjong Malang; N.W. by W. and N. by W. in Mount Wallich (25); N.W. by W. and N.W. by N. in Pearl's Hill (28). The most prevalent strikes recur in almost every locality where the strata are well exposed. N. by W. beds are seen in Sambo Kiehi, Tanjong Piñger, Tiloh Blangah, and both Sikijangs, Mount Palmer (24), Mount Wallich (25), and Tanjong Pagar; N.N.W. in T. Piñger, T. Dangas, Marambong, Sikra, Batu Blyer, Blakang Mati, Pulo Brani, Tilo Blangah, Sikukur, West Sikijang, Tanjong Pagar, Tanjong Malang, Government Hill (35); N.W. by N. in T. Piñger, Batu Blyer, T. Blangah, P. Brani, and West Sikijang; N.W. in Piel Ayem, Tanjong Gul, Batu Blyer, Hodin's Hills, Pulo Hantu II. (12); Pañti Chirmin (13), Pulo Brani, and Blakang Mati, Tanjong Malang, and the east point of Bintang; N.W. $\frac{1}{2}$ W. in Bukum, Jong, B. Blyer, B. Mati, West Sikijang, and Tanjong Puñgai; N.W. by W. in Sabaru, Batu Blyer, Hodin's Hills, T. Blangah, P. Brani, Blakang Mati, Timukul, Mount Wallich (25), Dickinson's Hill (29), and Pearl's Hill (28); in the shale-beds between Tilo Blanga and the Sepoy Lines (30), N.W. in the same beds to the west of Bukit Timah.

The variations in dip, like those of strike, have a general character of uniformity, with considerable occasional deviations. Vertical or approximately vertical strata occur in P. Blakang Padang, Sambo, T. Piñger; T. Busong, Jong, Bukum, the Batu Blyer Range, Hodin's Hills, the N.W. point of P. Blakang Mati, the Tiloh Blangah Hills, Mount Palmer. Oscillations may be observed in Marambong; in Busong; in the Batu Blyer Range of about 70°, or from 25° W.* to vertical and a few degrees from vertical to the E.; in Tanjong Piñger about 75°, or from 20° W. to vertical and a little to the E.; in P. Blakang Mati of about 45°, or from 50° W. to a little E. of vertical; in West Sikijang of about 90°, or from 45° W. to 45° E.; in

* W. and E. are here used for westerly and easterly.

East Sikijang about 35° , or from 45° to 80° E.; in the range terminating at Tanjong Batu (Singapore) about 85° , or from 50° W. to 45° E.

A high inclination being very common, a decided tendency to a dip in one direction is not to be anticipated. In several localities the rocks dip away from vertical on both sides, indicating the existence of synclinal axes, which in a few instances are seen, as in the N.E. point of Blakang Mati, Mount Palmer, &c. The only distinct anticlinal axis that I have observed is in a hill forming the south point of West Sikijang, where the strata are arched.

Notwithstanding this tendency to verticality, the westerly dip on the whole predominates in the western part of the district, for, although beds having an easterly dip are found in P. Mirambong, P. Blakang Padang, P. Busong, P. Jong, P. Sabaru, the Batu Blyer Range, P. Hantu, Blakang Mati, Sikukur, West and East Sikijang, T. Piñger, Mount Wallich, Mount Palmer, Kiltiney Hill (34), &c., and in some cases the dip is exclusively to the east, yet in most of these localities the beds that dip to the east are associated with a greater extent of beds that have a westerly dip. In the eastern part of the district, however, the dip is generally to the eastward, as at the Tikongs, Tanjong Puñgai, and the N.E. point of Bintang; but it is not uniformly so, for at Tanjong Lompatan it is about 45° W. In the next considerable exposure of rocks on the east coast, those of Sidili Point, to the north of the district, the dip is again easterly.

The strata, although where iron-rock and quartz occur often much disturbed, and sometimes in patches, broken up and mixed, forming a hard ferruginous breccia, are nowhere completely shattered, and the fragments confusedly mingled, except in considerable portions of Government Hill (34), Mount Sophia (48), and Mount Siligi (47), where no beds are distinguishable, the hard sandstone being scattered through the clay, at some places in angular blocks from a few inches to several feet in diameter, and in others as a fine breccia.

2. *The Plutonic rocks.*—The prevalence of felspar and hornblende renders these rocks very decomposable when kept exposed to the influence of water under ground. Hence while blocks rising above the soil suffer little waste, the mass below is almost everywhere converted into clay to a considerable but unknown depth. The only locality in the district where rocks have escaped decomposition in sufficient number and size to display their structure is Pulo Ubin. This has probably been owing to the steepness and narrowness of the range of which it consists, and the action of the sea along both its sides, which have combined to degrade and wash away the decomposed portions, thus leaving the more resistant masses to emerge from the soil and stand out above the influence of decomposition*.

All the numerous exposed blocks of Pulo Ubin have a large laminar or zoned structure. The laminæ are in different forms, flat, spherical, and concentric, and variously curved; but the systems of circles

* When an exposed rock is attacked, the decomposing portion is washed or falls off, and the decomposition is arrested for the time. Under-ground decomposition tends to spread unchecked on all sides.

and curves seen in horizontal sections are generally portions of a connected system, of which the external laminae are rectilinear, or nearly so. The principal rectilinear planes of division approximate in their general direction to N.E., S.W., or at right angles to the axis of elevation of the region.

Transverse and approximately horizontal planes of division exist, so that the rock tends everywhere to break into spherical, cuboidal, and tabular masses. Its composition in many places varies with the structure; and the unequal wasting thereby occasioned has given rise to very singular external forms in the exposed masses. The most common are cuboidal rocks, from a few yards to twenty and upwards in diameter; having their sides traversed by concave vertical grooves, varying in depth from a few inches to about six feet; the sides of the larger concavities being often grooved in their turn. These grooves generally extend from the base to the summit of the rock, a height in some cases of about 40 feet, although generally much less. On the upper surfaces, and on the sides where they are inclined, a succession of hemispherical and pear-shaped cavities are frequently seen; and, where they are wanting, less regular furrows, corresponding with divisional planes, generally supply their place*.

The true explanation of all these phenomena I believe to be that which I offered towards the conclusion of the paper mentioned in the preceding note. It is as follows:—"The blocks, protruding from the hills or ranged along the shores of Pulo Ubin, are more solid and less decomposable masses and nuclei, of which the forms, with the directions of their sides and axes, have, in almost every instance, been determined by structural planes, and which remain after the surrounding rocks have been disintegrated and washed away. With respect to the latter, it is obvious that while the island has been extending by the growth of alluvium in its bays, its more open coast has been slowly retreating, so that what was once a part of the solid land is now a band on its border washed by the sea, but still exhibiting numerous rocky remnants. The larger masses still evidently occupy their original positions. Frequently their seaward face is curved. Sometimes another mass stands behind, merely separated from that in front by a chasm whose sides are parallel. With respect to the decomposition of the rocks on the hills, the soil is entirely derived from this source, with the exception of a very slight superficial mixture of vegetable matter, which in many places is absent. In general, however, the blocks that remain are decomposing with exceeding slowness. One exception I noticed in the N.W.-S.E. side of a block about 20 feet in height. The laminae being inclined inwards, in disrupting by their own weight fall some feet in front of the base, where a long mound of earth has consequently accumulated. I have now only to revert to the grooves. The circumstances attending them which any hypothesis of their origin must explain are these: their general prevalence; the existence, however, of exposed rocks devoid

* I have described the rocks of Pulo Ubin in detail in a paper communicated to the Batavian "Genootschap van Kunsten en Wetenschappen," in 1847, and published in the 22nd vol. of their Transactions.

of them; their being commonly confined to the sides facing the exterior of the island, although sometimes found on other and even on all sides of a rock; their great depth and regularity; their general coincidence with divisional lines; their conformity to the course of rain; and their antiquity. It is this last circumstance which, presenting at the outset a great difficulty, leads, on further consideration, to what I consider the true explanation. That meteoric influences have been the great agents of erosion I have already suggested. But the antique and permanent character which is impressed on the great majority of the rocks, their vegetable coatings, the hardness and sharpness of the external edges of the grooves, prove that the rocks must have existed under very different conditions from the present, to enable atmospheric forces to produce results of such magnitude. The considerations which have hitherto occupied us in the concluding portion of this paper appear to me to indicate what these conditions were. The composition and structure of the external rocks, unveiled by the action of the sea on the beach, show zones of soft rock*, rows of globular decomposing masses, and of harder ferruginous spheroids, &c., susceptible of being detached, and a general tendency to perpendicular division. If, therefore, we conceive the external layer of the island, when it first became exposed to decomposition, to have resembled in character the zone that has been laid open for our inspection along the beach, it is easy to comprehend how the wasting away of the more decomposable parts might at last leave exposed masses, including bands of the less stubborn material already partially softened or disintegrated under ground, and that the action of the atmosphere and rain-torrents would gradually excavate the more yielding portions, until the solid remnants exhibited their present shapes."

II. METEOROLOGICAL AND HYDROLOGICAL INFLUENCES.

We have necessarily anticipated much belonging to this head. The mean annual temperature of the district is about $81^{\circ} 25'$ (that of the hottest and coldest months differing about $2^{\circ} 76'$); the range from 6° to 7° ; the fall of rain about 92 inches distributed over every month in the year, with a considerable excess from October to January.

All the climatic forces which operate on the land are affected by the two monsoons which prevail in the Indian Ocean and China Sea. During the S.W. monsoon from April to October, the Straits of Singapore being to the leeward of the mountain-range of Sumatra, equable weather is experienced in the district. Showers are frequent but seldom long-continued, and the monthly mean of the thermometer ranges from 81.21 to 82.31 .

During the N.E. monsoon from November to March, while the

* Some rocks may be seen along the beach with chasms 2 or 3 feet wide, the sides being quite hard and the bottom a soft decomposed substance. In such a case a zone of rock differing in composition from that adjoining has evidently been gradually decomposed and washed out.

Straits of Malacca are sheltered by the mountains of the Malay Peninsula; the eastern coast of our district is exposed to the full force of the monsoon, and the hills that stretch along it are not sufficiently high to destroy, although they considerably modify its influence in the Straits. During the greater part of this monsoon rains fall abundantly, the streams are frequently swollen, and sometimes overflow their valleys, the sun is obscured more frequently and for longer periods, the temperature is lowered, strong winds often prevail, and the sea is more agitated. It is, therefore, in this monsoon that the erosions and abrasions of the land referable, mediately or immediately, to the climate are greatest.

The heat and humidity of the climate affect the solid inorganic mass of the land chiefly by their favouring the decomposition of the rocks, the growth of vegetation, and the degradation of the soil. These operations are much less marked than those of the sea, but they are incessantly progressing over the whole land, and their effects, when accumulated by time, must be very great. The soil is always kept moist, which causes the decomposition of the plutonic rocks to descend to a great depth; and the rain, the most constantly acting and universal mechanical agent of geology, and here gaining a maximum of effect from the configuration of the ground, falls on an average about half the number of days in the year. When it is considered that every considerable shower in Singapore is the parent of innumerable little rills,—each turbid with fine clay and propelling quartz-granules, &c.,—which pour down the sides of hundreds of hills into the valleys, the great beds of alluvial clay appear less disproportioned to the size of the streams. These are very numerous, but, owing to the low level of the country, insignificant; the broad and deep salt-water creeks, which occur at short intervals along all the coasts* and seem to be the estuaries of considerable rivers, seldom extending beyond a few winding reaches, at the top of which they dwindle to petty rivulets of fresh water. None of the Singapore streams have courses longer than six miles, and it is only after heavy showers that they pour down a considerable body of water. The continental part of the district includes one large river, the Johor, the course of which, in a direct line from its source in Gunong Blumut to Tanjong Tikong, is about fifty miles; but it may be said to fall into the sea to the north of the district, because the long and broad estuary as far up as Tanjong Gidong is an arm of the sea. It has caused the formation of a large alluvial tract extending along the west side of the estuary. The other rivers of the mainland falling, with one exception, into the Old Straits, such as the Pulau, Sakodai, Tambräu, Libam, and Santi, have all elevated the beds of their valleys to some distance from the Strait to a level between low and high water-mark, and covered them with marine forests. Thus in every valley of the district some of the alluvial processes before-described are constantly proceeding.

The Straits, as we have seen, are sheltered from the direct influence

* The eastern coast of the mainland and the S.E. coast of Singapore must be excepted.

of both monsoons, and it is only when the China Sea is agitated by strong or continued winds during the N.E. monsoon that heavy waves roll into the Strait, and attack all the exposed shores of the three most easterly basins*,—the Peninsular coast from Point Romania to Marbukit, the Batam coast as far west as T. Siŋgkwang, and the south-east coast of Singapore.

The east coast of the Peninsula and the north coast of Bintang, lying on the China Sea, are at all times exposed to the action of its waves; and the abrasion at all the points, notwithstanding the solidity of the rocks, has been great. It is particularly observable at the promontory terminating in Tanjong Puŋgai and Tanjong Kĭnawar, the first of which points is covered with massive globular blocks of iron-ore gleaming like gigantic balls of polished metal, the indestructible fragments of a hill, one-half of which has been destroyed, and of which the cliff behind, notwithstanding its masses of iron-rock, is year after year yielding more spoil to the waves. The hill at the second or northerly point, being less ferruginous, has been ground down to the level of the sea, and nothing is now left but the long quartz-penetrated ledges of its foundation. The extremity of the Siglap and Tana Mera hills on the east coast of Singapore, being composed of soft sandstone and conglomerate, are annually suffering abrasion; large fragments falling down during every N.E. monsoon. The waste here must have been enormous, for the abraded sand has completely blocked up the whole S.E. coast of ancient Singapore, so that for about ten miles the only spots where the sea touches it are the two hills, the abrasion of which forms Tana Mera Bĕsar and Tana Mera Kichi (the Red Cliffs). This long sand-bank, which has completely dammed in all the water of the plain of Singapore, is the best measure which the Straits afford of the power of the waves raised by the N.E. monsoon, and it is worthy of notice as an instance in which the compensatory effects of abrasion far exceed the destructive, as respects the superficial proportion between land and sea.

"The sedimentary ranges to the west of Singapore plain, which are variously indurated, charged with iron, and highly inclined, have opposed greater resistance to the sea; but the quantity of soft clays and shales which they contain has facilitated its action, and long ribs and ledges of iron-masked rock, stretching across the banks, mark at once how difficultly destructible they are in themselves, and how unavailing they have proved to save the hills which they bound together. At one place the sea has cut through some highly indurated strata and divided an island (Blakang Mati) in two, and a strong current rushing through this narrow gateway into New Harbour, its old circuitous route around the northern end of the island has gradually been filled up with mud, and the detached northern half thus united to Singapore. The islands in the Strait are all greatly abraded†," and many of them have long ledges running out

* The tidal current sometimes runs for eighteen hours of the twenty-four into the Strait during springs, and for two or three days without interruption during neaps.

† This paragraph so far is copied from my Essay on the Geology of the Peninsula cited before.

from their extremities or exposed at their sides. When the sea has worn down a portion of one of these islands below its level, the coral-line polyps that incrust its outer margin, where it sinks into deep water, begin to advance upon it towards the land and preserve it from further abrasion. Were it not for the causes to be mentioned, this curious organic shield would effectually defend the rocky foundations of the islands from the attacks of the sea; and in process of time, when the Strait was denuded of all the hills that now rise above its surface, each of them would be represented by a coral-reef of an irregular annular shape*. But in many localities, after the corals have flourished over a considerable horizontal space for some time, yielding strata of sandstone come under the influence of the waves, which grind them down and spread the sand over the coral-field. The speedy consequence is that the polyps perish, and in their place a bed of sand and dead coral is left, which sometimes cements into a hard calcareous layer, but generally becomes a prey to the sea. Living coral-fields are found in all parts of the Strait, but most abundantly in the north-eastern portion of the most westerly basin in the shoals around and between its numerous islets. Beaches from which corals have perished occur frequently; amongst the largest being those in front of Blakang Mati village, the east coast of Sambo Bësar, and Bukum. On coasts of soft wasting sandstone they are seldom found; and in the proper mangrove localities, that is, in sheltered places where mud accumulates, they are either wanting or are scattered in small and weakly patches on spots where sand occurs. Their proper habitat is a beach or shelf of indurated rock abraded to a level below that of the ebb-tide, sinking at its outer margin into deep water, and with a free exposure to the sea so as to be constantly scoured by the waves. Such shelves seldom preserve all these conditions up to the land, so that there is generally an inner space from which the waves retire at ebb-tide. It is still moist, full of small pools, and dotted with pieces of living coral, but the continuous coral-beds growing luxuriantly in

* If the Strait were to remain perfectly undisturbed by subterranean forces, neither undergoing subsidence nor elevation, it is obvious, from what we now see to be going on, that these two antagonist powers, water and the coral-polyps, would, in the course of time, leave the Strait without any vestige of land, save some reefs and rings covered and shielded by the coral. These two simple and ever-operating agencies, mechanical and organic, appear sufficient to explain the fact, that patches of land are found over vast spaces of ocean at the same level. Let the present condition of things last long enough, and an aggregation of mountain-groups, like that of the Malay Peninsula, will be abraded, broken into islands and coral-reefs, until island after island is worn down beneath the level of the sea, those only remaining at that level in which the conditions for the continued existence and growth of coral are maintained throughout the process of degradation. In endeavouring to follow out such a process from the facts presented by a coral-line sea, like that of the Straits of Singapore, the shores where coral is absent are as instructive as those where it is present; for whatever successive changes the configuration and disposition of the land may undergo in the progress of denudation, the same causes will continue to favour and oppose coralline growth so long as wasting shores and streams of fresh water exist. The process would accelerate as it advanced, and a condition of things approximating to that of the archipelagoes in the open ocean be reached, in which both the mechanical and organic powers are greater than in the Strait of Singapore.

all their variety of rich colours and beautiful forms, giving a gorgeous pavement to the sea, are not found until this space is passed.

In some places solid beds of calcareous conglomerate are forming on the beaches; and ferruginous conglomerates, breccias, and sandstones are found wherever highly ferruginous rocks exist behind sandy shores.

III. RELATION OF THE ROCKS TO ANIMAL AND VEGETABLE LIFE; AND THEIR ECONOMICAL USES.

Although the soils of the district have not the fertility of the volcanic and calcareous soils which occur in many parts of the Indian Archipelago, they are covered with an indigenous vegetation of great vigour and luxuriance, supporting numbers of animals of different species. The hills of plutonic rock support dense and continuous forests, composed of more than 200 species of trees*, many of which are of great size. So long as the iron is not in such excess as to recombine the clay into stone, or render it hard, those soils which contain most iron are the most fertile. The purely or highly felspathic are the worst. But even felspathic soils, when they have a sufficient proportion of quartz, are, in this climate, capable of producing an abundant vegetation.

Although it is obvious to every observer that there is no kind of soil in the District for which nature has not provided plants that flourish luxuriantly in it, yet it must not be hastily concluded, as some have done, that this exuberant vegetation indicates a general fertility in the soil. It is found, on the contrary, when the native plants are destroyed, and the land employed for agriculture, that there are very few soils in which cultivated plants not indigenous to the region, but whose climatic range embraces it, will flourish spontaneously. While the cocoa-nut, betel-nut, sago, gomuti, and the numerous Malayan fruits succeed well with little care, the nutmeg and clove are stunted and almost unproductive, unless constantly cultivated and highly manured; yet the climate is perfectly adapted to them. Place them in the rare spots where there is naturally a fertile soil, or create one artificially, and their produce is equal to that of trees in the Molucca plantations. With respect to the indigenous plants, gambier, pepper, and all the fruit-trees flourish on the plutonic hills, provided they are not too deficient in iron and quartz. The hills of violet-coloured shale, where they are not too sandy, are equal to the best plutonic soils,—those, namely, in which there is a sufficient proportion of hard granules to render them friable, and sufficient iron to render them highly absorptive of water without becoming plastic. The sandstone and very arenaceous shale soils are the worst. Of the alluvial soils, the sand, particularly when it contains a mixture of vegetable matter or triturated shells, is the proper soil of the cocoa-nut, and the vegetable mud is best suited to the sago. When the country has been better and longer drained and cultivated, the latter soil will become a rich mould; at present it is every-

* My list contains at present 217 trees, but it is not complete.

where too wet and sour to make a fertile soil. Rice is grown on some patches of it. The bluish sea-mud contains good ingredients, but clay is in excess, and the animal matter appears to assist in rendering it hard and untractable when it is not saturated with water. Even for such a soil nature has provided plants useful to man, for the betel-nut and some of the indigenous fruit-trees grow well in it with little cultivation. Although there are cultivated plants adapted for every kind of soil in the District, and it has indigenous tribes who can live exclusively on its yams, sago, fish, and wild animals, it is incapable of feeding a population of the more civilized races; and the latter must always be dependent on other countries for the great necessary of life, viz. rice.

The rocks which are used for economical purposes are not numerous. The only edible one is the fine clay called *ampo*, which is made into thin cakes, smoked, and kept for use.

The plutonic rocks, and the indurated sandstones and conglomerates, are used for the foundations of houses. Lateritic stones are sometimes used by the Malays as pedestals for the posts on which their small houses rest. Granite is used for steps, milestones, tombstones, &c. Of the blue alluvial clays the bricks and tiles are made, of which the town of Singapore is built. The fine kaolin which abounds has been found the best adapted of any in India for the manufacture of porcelain, but no manufactory has ever been established*.

TV. TRACES OF THE GEOLOGICAL HISTORY OF THE DISTRICT.

Having thus given a general description of the actual constituents and arrangement of the rocks of the district, we have next to inquire whether they preserve any evidences of its past geological history, and I think the following conclusions may be safely adopted.

1. Ceasing for a moment to view the district as isolated, it is established that it is a portion of the zone of elevation extending from the Himalayas (or their vicinity) to Banka, or rather to the granite patch in Java; approximately parallel to the plutonic zones of Burmah, Siam, Cambodia, and Anam, which seem to terminate in Borneo, and surrounded by the great volcanic band of the Indian Archipelago, the most active western portion of which, Sumatra, is semi-plutonic, and is parallel and adjacent to the Malay Peninsula; separated from it only by the broad and shallow submarine valley of the Straits of Malacca†.

2. The sedimentary strata of the district were deposited prior to the elevation of the Malay Peninsula, and therefore prior, probably, to the elevation of the whole south-eastern region of Asia.

3. They were formed from the abrasion of a tract composed chiefly of clays, and probably partially elevated above the level of the sea, or at least so near its surface as to be subject to strong currents. These appear to have prevailed along some of its wasting shores in

* See Dr. O'Shaughnessy's Report of Experiments made by him for Government.

† See "The Physical Geography and Geology of the Malay Peninsula," Journ. Ind. Arch. vol. ii. p. 89-93.

the vicinity of Singapore, for the water-worn pebbles of the conglomeratic beds entirely resemble, in their variable size and occasional deficiency of abrasion, the gravel at present accumulating in the vicinity of the sea-cliffs. The mode in which they are disposed shows that they were from time to time driven and irregularly heaped by violent currents or waves.

All the sedimentary beds of the district form one series accumulated from the debris of the same rocks, and the region underwent no violent upheaval during the period of their deposition. This is an inference from the generally uniform mineral character of the beds,—particular kinds of clay or shale prevailing everywhere, and the conglomerate layers, wherever they reappear, exhibiting the same peculiar clayey matrix and the same prevailing pebbles. No instance of unconformable stratification has been observed.

The ancient land consisted chiefly of two rocks; 1st, a fine, tough, easily frangible clay, of which the prevailing colour was a dull violet; 2nd, a granite composed of quartz, chlorite, white felspar, and a fine silvery talc, the last sometimes wanting, and all the ingredients apparently varying greatly in their relative proportions at different places. The first rock, reduced to sediment, produced the shales and clays of the district, and, in rolled fragments, furnished a portion of the pebbles of the conglomerates. The second rock, disintegrated and partially decomposed, produced by its debris the great bulk of the conglomerates, of which the matrix is generally quartz-granules mixed with more or less of chloritic or steatitic earth with greenish and yellowish hues. It also furnished a few shale-beds in which chlorite is contained. The quartz-pebbles, generally whitish, but sometimes of black and other colours, were probably derived from this granite also; the quartz in some specimens tending to segregate. It probably existed likewise in contemporaneous veins and nodular masses. The granite, where free from chlorite or the larger developments of quartz, furnished the granules of the sandstone-layers; or these may have resulted from a more thorough decomposition of the granite, and the washing of the chlorite and felspar to a distance from the place where the sand accumulated, just as we see, at present, partially angular and rolled fragments accumulating in the vicinity of wasting points, the more finely disintegrated sand carried to some distance and accumulated by itself, and the clay suspended and deposited over a far wider surface, extending to a considerable distance.

4. The rocks of this ancient tract differed from the existing plutonic rocks of the district, and the latter were not in existence at the time of the deposit of the sedimentary beds, for no fragments resembling them have been found in these beds, and many of the rocks which most abound in them nowhere exist in the present plutonic rocks. The most conclusive fact is the entire absence of iron-masked pebbles or fragments in the sedimentary strata. The abundance of ferruginous walls in the latter and in the decomposed plutonic rocks, and the constantly recurring layers of ferruginous pebbles and blocks in the modern debris on the surface of the land and along the shores of the wasting points of both formations, prove that beds derived

from their decay will be principally characterized by their profusion of such pebbles and blocks.

5. The upheaval of the sedimentary rocks was in general attended with great violence, as is shown by the irregularity in the strike and dip of the beds in numerous circumscribed localities, and the breaking up and mingling of those in Government Hill, &c. Where there are most evidences of violence, there is the greatest development of iron or quartz. The line of disturbance, for instance, which goes through Government Hill, and is continued in the hill south of it, ending in Tanjong Malang or Batu, is marked in the latter by iron-masked blocks and gravel on the surface, sandstone indurated and pervaded by an iron-honeycomb, iron-mammillated crusts on the sides of the beds, &c., and in Government Hill by the clayey beds adjacent to the disrupted sandstone being converted into the typical laterite, and by the sandstone at some places near the junction being broken into a mass of small fragments recemented by iron and clay*.

6. The formation and intumescence of the plutonic rocks of the district caused this upheaval, and gave the prevailing direction to the stratification and to the ranges of hills, this direction agreeing with that of the plutonic zone of the Peninsula.

7. The partial metamorphosis, induration, and *ferrugination* of the sedimentary strata were effects of this plutonic action.

8. The vast abundance of iron evolved during the plutonic intumescence, and rising into the strata, is the distinguishing peculiarity of the Malayan zone of elevation†, and its wide-spread effects so strongly strike the eye that the other changes are less observed; but the latter evince even more decidedly the potency and the unity of the elevatory agent, and are particularly worthy of attention as assimilating that agency to the more common kinds of plutonic metamorphosis. The soft strata in some places have been converted into Lydian stone, porcellanite, clay-slate, hornblende slate, steatite, and chert. If we extend our observations a little beyond the district, we find similar changes on a much greater scale. Thus the eastern coast of Pulo Krimun Kichi (Little Carimon), which rises only eighteen miles to the west of the last of our sandstone islets in that direction, is composed of a conglomerate converted into the hardest chert. In many of the islands in the China Sea off Pahang the metamorphosed rocks are so indistinguishable from greenstone, that we can only recognise them as sedimentary by the presence of pebbles which some of them contain. In Pulo Tioman the junction of granite with a green, grey, and black hornblendic rock is well displayed, the granite sending veins into the hornblende, and masses of the latter being sometimes imbedded in the former, clearly proving that the hornblende, whether volcanic or (as I believe) metamorphic, is the more ancient rock.

* I trust to be able ere long to give a full account of the mechanical effects of the upheaval in different parts of the district.

† But common to it with the adjacent belts rising from the same platform—that of S.E. Asia—and the approximately parallel and probably contemporaneous Peninsula of Southern India. This peculiarity is found in S. India, Bengal, the Malayan belt as far north as it has been examined, Borneo, and the N.W. part of Australia.

9. One of the most striking characteristics of the district is the protean nature of the plutonic transformations, remarkable alike in the diversity of the plutonic rocks themselves, and in the numerous degrees and kinds of change which the plutonic agency has produced on the strata. There are considerable tracts in which the granite undergoes little variation, but which differ decidedly from each other. There are tracts where the plutonic rock graduates or passes abruptly from large-grained granite to the most finely grained, or, losing its quartz and mica, passes into greenstone or into compact felspathic or hornblendic rocks. Amongst the sedimentary rocks we find that some, near the line of junction with the plutonic, have merely become ferruginous, while others, apparently distant from it, have become highly indurated. The same rock, completely iron-masked at one place, exhibits no change whatever a few feet off. We have seen that the mechanical action to which the strata have been subjected participates in this want of uniformity.

10. Is the explanation of this character of the district,—the absence of all other sedimentary formations but the existing limited one, the entire disappearance of the ancient rocks which composed the district when the present strata were being formed, and the mode in which the plutonic exhalations have freely risen through the latter,—to be found in the circumstance of the plutonic action having extended to the surface of the ancient land (whether beneath the sea or not is another question) and assimilated it all, excepting a thin crust of the newest and highest beds, or rather bands and patches of that crust, which now lie scattered over the great plutonic mass of the Peninsula? It is clear that granite does not require a greater pressure for its formation than the so-called volcanic rocks into which it passes. In Pulo Ubin and other localities, we see, in the same rock, at one place a perfect granite, composed of large crystals of quartz, felspar, mica, and hornblende, and at another a compact or finely granular greenstone or basalt, so that the granitic form does not necessarily depend on the depth at which crystallization or plutonic metamorphism takes place*.

11. At the time of the upheaval of the strata they were new, and had neither been consolidated by great pressure nor indurated by any prior plutonic action. Where they have escaped the influence of the ferruginous and siliceous exhalations, and the heat given out during their elevation, the more aluminous beds are still soft clays, and the arenaceous slightly coherent sand, sometimes not distinguishable from the beds of clay and sand forming along the shores at present.

If the now visible plutonic rocks had been formed at a great depth and under a vast equal pressure, they would have been more uniform in their composition and structure, and in their action on the immediately superincumbent strata. They would have more slowly cooled down, and the adjacent strata, long exposed to the heat, would have acquired a certain general induration. If the strata had still been accumulated over the plutonic mass to a great thickness when the latter ceased to reduce them, and been subsequently pared down to

* See account of the rocks of Pulo Ubin, p. 22, Trans. Batav. Soc. Arts and Sc.

their present remnant by denudation, they would not have been broken up in the manner in which we see them to be, but in a mode resembling that in which the great aqueous formations of Europe and America are arranged around or on the flanks of plutonic mountains, and in which the strata of the ancient land may perhaps yet exist in some parts of the flanks of the great intumescence or submarine elevation of the Malayan Archipelago where it rises from the bottom of the Indian Ocean.

12. The prevalent layers and heaps of volcanic-like gravel were not ejected. They are merely the accumulated debris of walls and veins arising from the decomposition and washing away of the softer parts of the containing rock, and remaining by virtue of their ferruginous and jaspideous composition, which resists atmospherical action.

I have found all the forms of the gravel in the places where they were generated,—that is, in the walls traversing the solid or partially disintegrated rocks.

13. The iron was not injected in a fluid state into the strata, as lava is into fissures, but was either imbibed, or, as is most probable, conducted by vapours, gases, or electric currents. The ferruginous and quartzo-ferruginous rocks, *including laterite*, wherever minutely examined, prove to be the original rock of the situs metamorphosed, and not a foreign rock injected from beneath; and there are evidences in all classes of the rocks affected, that the iron, although ascending in abundance wherever the rocks were fissured or weakly coherent, was thence transfused without its exerting any mechanical force, and without the slightest disturbance of the previous arrangement of the particles of the rock*.

14. It is probable that the plutonic rock was not in a molten state up to its contact with the strata, because if it had been, the fissures, which must have been caused by the irregular force exerted on the strata and their frequent disruption, would in that case have given rise to true dykes and veins, *i. e.* those formed by fluid plutonic or volcanic rock being pressed up into rents.

15. The numerous ferruginous dykes which ramify in the plutonic rocks were contemporaneous, because if we suppose them to have been true dykes produced subsequent to the intumescence and consolidation of these rocks, the enormous force, necessary for rending and cracking them in so many directions, would have extended the fissures into the superincumbent strata, and the injected dykes would

* The entrance of the iron was, however, in many cases simultaneous with mechanical strainings and ruptures of the rocks, and, when rising abundantly into the main walls and forced thence into the lateral ones, must have exerted considerable pressure. The mechanical straining and the exhalations were both effects of the same cause, the plutonic intumescence. The only case which I have observed of a direct connection between the matter of the walls and a mechanical alteration in the adjacent rock is seen when quartz is largely developed in lumps in the quartzo-ferruginous dykes. The adjacent unaltered layers are pressed together and bent so as to make room for the quartz. Even here I do not think this is attributable to a large accession of new matter, but mainly to the expansion of bulk attending the conversion of the siliceous particles of the original rock into crystalline quartz.

have been continued into these strata also. The contemporaneity of the ferruginous dykes is further proved by the mode in which they are connected with the adjacent granite in localities, such as the coast of Batam and the Water Islands near Malacca*, where they can be examined in the undecomposed rock.

16. The more highly ferruginous parts of the walls in the plutonic and sedimentary rocks are identical in appearance, the proportion of peroxide of iron which they hold being so great as entirely to disguise the original rock. The peroxidation of the iron tends to obliterate all traces of its original condition, and to give a uniformity of aspect to all the rocks which I have designated as iron-masked†, so that we shall not be able to see in what form the iron exists prior to the exposure of the rocks to atmospherical influences, until we have much deeper sections than are anywhere exposed at present. It is obvious, however, that the effect of decomposition has been to saturate the water in the rock with oxidized iron, and diffuse it more uniformly through the arenaceous and aluminous matter of the walls. If the oxidized crust of the district, which probably descends to a considerable depth, were pared off, the peculiar aspect and structure of the iron-masked rocks would disappear; the deep rusty, dark brown and black colours, and the scoriform, vesicular, and tubular structure of the walls and lateritic patches, which now so powerfully arrest the attention, would be nowhere recognized, and in place of them we

* Malacca, including Naning, in all its main features, is identical with the district of the Straits of Singapore. All the phenomena of elevation, ferrugination, &c. which we see here are there repeated. See 'Five Days in Naning,' by J. R. Logan, Esq., Journ. Ind. Arch. (for May 1849) vol. iii. p. 282.

† "The interest which the discussions respecting laterite have given to that rock tends to invest it with undue importance geologically. The ferruginous emissions have penetrated all rocks indiscriminately, and their action on sandstones, grits, and conglomerates is as well-marked as that on clays, marls, and shales, although the latter only produces proper laterite. Even in the clays, laterite denotes one only of many degrees and forms of alteration. To express the origin of these rocks, and its unity, to record the cause of the difficulties which they have presented, and to distinguish them from true metamorphic rocks, I would propose, avoiding any new technical names, to term them simply the *iron-masked* rocks of the Indo-Australian regions. This term will include the principal or plutonically ferruginated rocks, which, without being either completely reduced or metamorphosed, have been either wholly disguised or partially altered by ferruginous emissions, which have saturated them in the mass, or only affected them in fissures and seams, or have been interfused between portions of the rocks not actually separated by fissures, but intersected by planes of mere discontinuity, the sides of which have an imperfect cohesion, or having a common border of inferior density and increased porosity, caused either by interruptions in the original deposition of the matter of the rock, or by unequal stretching, or incipient cleavage. The term may be also extended, perhaps, to those sedimentary beds in which the iron-saturation, although coeval with the deposit of the other constituents of the rock, has served to obscure or conceal their true nature as well as the derivation of the beds themselves. These beds appear to have been sometimes formed by superficial layers of gravel, &c. being permeated by iron-solutions. With these must not be confounded the broad bands lying over and beside the heads of iron-masked dykes, and which, having been in a loose gravelly or fragmentary state at the time when the plutonic emissions passed through them, became converted into hard, and occasionally scoriuous, ferruginated conglomerates, &c., and are therefore proper plutonically iron-masked rocks." (Journ. Asiatic Society (Calcutta), vol. xvi. p. 521.)

should see in the walls of the sedimentary rocks the ordinary matter intermixed with pyrites and other unperoxidized forms of iron-ore, and, in those of the plutonic rocks, similar forms of iron, accompanied probably by a predominance of those common plutonic minerals in which iron is a large ingredient, owing to its accumulation there at the time of the crystallization. The ready decomposition of iron-pyrites would account for the great depth to which the walls and veins have been peroxidized.

17. It is of great importance to determine, if possible, the state in which the iron was originally deposited, and why it was given off so abundantly by the plutonic intumescence. It appears probable that it was volatilized in combination with sulphur.

Is its abundance owing to the ferruginous character of the old land out of which the plutonic rock was formed, and do the highly hornblendic portions of this rock indicate a more than ordinary prevalence of iron in their parent sedimentary rock? The chocolate-coloured shale-band would probably produce such a granite. Sandstone appears along the western boundary between the granite and this shale; but near the present Arrack Distillery (33), where they are in contact, the granite is very ferruginous, decomposing into a dark-red soil, and containing half-decomposed masses approaching to iron-ore.

18. I have searched for the continuation of the ferruginous walls of the sedimentary rocks into the plutonic rocks. Unfortunately the line of junction of the two formations has not anywhere been laid bare, all the points exposed to the action of the waves being exclusively of the one or the other, and the deep superficial bed of earth conceals the undecomposed rocks in the interior. Last year a cutting of a few feet at a point (the south angle of Sri Menanti nutmeg-plantation at the junction of the River Valley and Tanglin roads (39)) where the formations meet, exposed a portion of a stratum of coarse-grained sandstone lying in an irregularly convex form on the decomposed plutonic rock, which appears to have been syenite. This has covered it in some places, and probably the whole or the greater part of the mass was originally imbedded in it. The surfaces are irregular and uneven, but have a general correspondence with the plane of the bed. It varies in thickness, having in some places 4 or 5 feet of well-preserved sandstone, while in others it shows only patches and ferruginous plates in the syenitic clay. It varies from a fine sandstone to a coarse grit, and some parts are conglomeratic. In some places it has been converted into a mass of hard and difficultly frangible iron-ore. Where least altered the rock is hard but brittle; the granules and pebbles of quartz are all preserved, but they have a dull whitish and yellowish colour, and rest in a basis of hydrous peroxide of iron, varying in colour from dark-brown to black. The microscope shows this basis to be not compact but vesicular, fine scales or plates surrounding the quartz-granules and partly filling the interstices between them. Where the clay meets the sandstone, the latter has generally a continuous crust or plate of shining, black, compact ore. Irregular systems of a similar crust are sometimes seen with the clay between them. At other places the whole is cellular, and a perfect

laterite is produced. Sometimes this structure is seen continued from the bed of sandstone into the clay, where the ferruginous ramification and blotches become softer and admit of being cut by the hoe.

In some places the iron has not pervaded the sandstone, but only penetrated it in layers or veins. In these cases it has disintegrated into a soft ferruginous sand. Where the iron has been diffused the sandstone is undecomposable, and remains hard but brittle. This locality exhibits, 1st, solid, compact, shining ore; 2nd, cellular ore filled with decomposed syenite; 3rd, variously iron-masked sandstone and grit.

I have hitherto proceeded on firm ground, but, in extending these deductions much further, must take a direction where many may not be disposed to follow, and where we cannot yet hope to obtain a sure footing. I venture, however, to suggest as hypotheses deserving of deeper investigation the following additional views which my observations strongly tend to support.

Whether the plutonic rocks are simply the product of a heat above the melting-point, or were formed at a lower temperature by chemical and electrical action induced by the heat, the sedimentary rock in contact with and raised upwards by them must have been partially reduced into the plutonic mass. In either case, the upper portion, at least, of that mass is a recomposition of the lower part of the sedimentary rock, with an addition of new ingredients from below; because, so long as the heat was so intense as to produce perfect mixture, agitation, and motion *inter se* of the reduced matter and the plutonic mass into which it entered, the upper limit of the bubble must still have been distant. So long as this level was not reached, each successive portion or layer of the sedimentary ceiling must have passed through similar changes before it became obliterated by complete absorption into the plutonic intumescence,—first becoming affected by the mechanical pressure, and then by the increasing heat and exhalations of the labouring mass below. Therefore, at whatever level the plutonic force, whatever its precise nature may have been, ceased to assimilate the superincumbent rock, this rock, to a certain distance above that level, must have passed through the same stages of alteration, exclusive of the ultimate one, which the plutonic mass in contact with it had itself gradually undergone. Whatever changes we can now observe in the remaining sedimentary rocks adjacent to the plutonic bosses, we may conclude that the *plutonization* of the latter commenced in the same way; and this we may do with perfect confidence, when we find that these changes are seen over a great region. If these postulates be well-founded, the phenomena of the district of the Singapore Straits, which are repeated throughout the whole chain of the Malay Peninsula and its prolongation to Banka and Billiton, disclose to us this fact, that the conversion of the sedimentary into plutonic rocks began with their being pervaded, in fissures and lines of inferior cohesion, with ferruginous exhalations, producing ferruginous walls and veins.

Since in the now solid plutonic rocks we find similar walls and

veins, the presumption is that these are the same walls and veins by which the metamorphism commenced. This is a simpler theory than that which would require the process of plutonic assimilation, as it advanced, to have first obliterated those originally formed, and then created a new system so remarkably similar to the old one. In what way could such systems of ferruginous walls and veins have been produced *de novo* in a plutonic mass thoroughly melted down into uniformity? If it were clearly demonstrated as a general theory that plutonic rocks are congealed from a state of fluidity, it might, notwithstanding these and other difficulties attending its application to the rocks of the district *, require us to reject the simpler theory which these rocks inculcate. But plutonic geology is in too imperfect a state to entitle us to assume any general theory as established, and nothing but the assumption that granite is necessarily, everywhere, a product of complete igneous fusion, requires us to suppose that two successive systems of ferruginous walls were produced, one at the commencement, and another before the conclusion, of the plutonic change.

If we are satisfied to suppose that only one system was produced, until some proof be adduced that it could not possibly have remained unobliterated during the plutonic process, these walls and veins become objects of the highest interest and importance, and all the phenomena connected with them demand attention.

I shall only here advert to one, the concentrically laminar coloration, frequently seen on a small scale, and so splendidly developed in Sabaru, Jong, and Bukum. The very same zoned coloration is observable in some of the sections cut into decomposed plutonic rock, such as that where the public road crosses Mr. Hewetson's Hill, where similar zones exist on the sides of a ferruginous dyke. The neighbouring solid greenstone possesses the same structure as this decomposed rock, nests of hornblende, corresponding to the red patches, and hornblendic lines, often very minute, to the curved zones. When we examine the great developments of solid plutonic rock, we find that they present similar phenomena. The granite of Pulo Ubin, for instance, when its coherence is weakened by exposure to the action of the sea, exhibits connected systems of concentric and variously curved laminæ, in which one system is interrupted and modified by another, and accommodates the shape of its curves to it, precisely as we see in those of the coloured lines of the sedimentary rocks.

If we do not adopt the theory of the necessity of granitic fluidity, may we not suppose that, as the plutonic action first invaded the superincumbent strata in the ferruginous walls and veins, it operated principally through them wherever they existed; that even when complete assimilation took place they still remained, however modi-

* Before I had ascertained the connexion between the ferruginous walls in the sedimentary and those in the plutonic rocks, I endeavoured to explain the phenomena of composition and structure presented by Pulo Ubin, in accordance with the theory of congelation. See the 'Rocks of Pulo Ubin,' *loc. cit.* pp. 26 to 40.

fied* ; that the concentric ferruginous zones to which they gave birth determined the structure of the ultimate plutonic rock, and, to a certain extent, the arrangement of hornblendic and other ferruginous minerals ; and were perhaps the channels by which an electro-chemical operation, emanating immediately from the walls, was made to pervade all the adjacent parts of the rocks, until crystallization was induced, and the metamorphism was complete ?

The district, considered as more or less metamorphosed, exhibits two sets of phenomena, one where the ferruginous walls and lines are the principal feature, and the other where they are rare or absent. In the first case the plutonic action appears to have extended to great distances from its place of full operation, and in fact to have had no limit but the length of fissures and planes of discontinuity in the superincumbent rocks. On the other hand it has often been extremely weak, hardly extending beyond the walls, and leaving the adjoining rock unaltered. In the second case the plutonic action appears to have from the first pervaded the whole mass of rock, as far as it operated, and to have produced the following succession of changes, examples of all which occur :—1st. The original rock is indurated in different degrees ; 2nd, a subcrystalline texture is induced, but without any perfect crystals being generated ; 3rd, crystals of particular minerals begin to appear in this basis ; 4th, the same process is continued and various forms of crystalline rock are produced, the granitic being that where the conditions of crystallization were most favourable, either from the longer continuance or greater intensity of the plutonic action. The process when preceded by the formation of ferruginous walls is the same in effect, as these merely tend to determine the manner in which the iron is diffused, the lines where crystallization begins, and particular minerals are generated, the structure in the mass of the regenerated rock, and the distance to which the plutonic action, in its various degrees and with its various results, extends.

It is evident that fusion did not precede crystallization in the progress of change up to the formation of the highly crystalline porphyries of Pulo Nanas, South Point, and many places in the eastern islands, and in the greenstones and other highly hornblendic rocks which are undoubtedly the common dark shale of the District disguised, since the graduation from the one to the other can be traced. On the other hand there is no direct evidence that the granitic form was preceded by fusion, while all the facts we have been considering, particularly the generation of other crystalline forms without fusion and the passage of these forms into the granitic, tend to show that it may be produced without fusion. The truth probably is that powerful electro-chemical action can produce all the effects of crystal-

* In the course of the crystallization of the rock they were probably converted into ores of iron where the iron was in excess. In some walls and parts of walls, and in the lateral radiating zones, where the iron was too thinly diffused to generate proper ores, it would produce a predominance of hornblende, black mica, and other ferruginous minerals.

lization from fusion *, and that the whole series of nether-formed rocks, volcanic and plutonic, may be produced in both modes. It is probable also that wherever an electro-chemical action is so intense as to generate the granitic and other highly crystalline forms, it emanates from a lower region where the heat is so great as to cause fluidity†.

Remarks on the accompanying Geological Map‡.

The basis is the Chart made from the Surveys of Mr. Thomson and Captain Congalton, published in 1846. The rivers, creeks, and alluvial tracts of Singapore I have added from later and unpublished surveys by Mr. Thomson, which he has kindly allowed me to use; and from the same source I took the great majority of the additions and corrections of the names of localities. The insulation of Tanjong Surat and the rough outline of the estuary of the river Johōr above it are merely from my own observations, and not from a survey. The coast of Batam has not been surveyed. A general outline with some blanks appears on the published chart. These blanks and the names of localities § I have filled in from my own observation, as well as the names of islands to the west of it from P. Blakang Padang to P. Kapal. The Silat Batu Haji I copied roughly from a Dutch chart. The orthography of the Malayan names, which has hitherto varied greatly, has been corrected in accordance with the system which I have lately adopted in the 'Journal of the Indian Archipelago,' as the most simple, uniform, and accurate which occurred to me, after giving much consideration to the subject.

The geological colouring is entirely from my own observations, with the exception of Pulo Pisang||, Gunong Pulái¶, the islets to the N.W. of the Krimuns (Carimons), a few places on the coast of the mainland in the Silat Tambrāu, the south coast of the S.E. part of Johōr from Tanjong Stapah to Point Romania, the islets and rocks off that Point, and the coast of Bintang with the islets and rocks off it, all which are from information, and partly from specimens, given me by

* In the crystallization of a molten rock the same electro-chemical action may operate. The process must be greatly similar in both cases. The electro-chemical force segregates minerals and gives a symmetrical structure in spite of the solid form of the rock in the one case, and in spite of its fluid form in the other. The solid form tends to keep each particle of the matter of the rock in the place where it has been deposited, the fluid form tends to keep them all mixed up indiscriminately. The electro-chemical agencies of plutonic regeneration perform their office unshackled by either kind of resistance.

† Fusion is perhaps not necessary to account for plutonic intumescence and the consequent elevation, rupture, and inclination of superincumbent strata. If granitic crystallization is always attended with expansion (see the "Rocks of P. Ubin," *l. c.* p. 38), the amount in a vast plutonic mass like that of the Hindu-Chinese and Malayan Peninsulas, Sumatra, and Borneo, with all the inequalities of degree to which the process is evidently subject, might be sufficient to account for the condition in which the sedimentary remnants are now found resting on it.

‡ The lithographed Map, Pl. XVIII., is reduced from the original map coloured by the Author.

§ Few names except those referred to in the paper appear in the reduced map.

|| See p. 318, *note*. Not included in the accompanying Map.

¶ G. Pulai (2152 feet high) lies N.W. by W. of S. Sakodai, beyond the limits of the map. It forms part of a tract of "Crystalline rocks" running N.W.—S.E.

Mr. Thomson. Long Island to the south of Barn Island I have not visited. It is described in an account of a voyage round Singapore (by Mr. Crawford, I think; long ago published in the 'Singapore Chronicle') as sandstone and clay-slate.

The alluvial valleys, running into Singapore Plain from the N.W., have been laid down with much care; and, as Mr. Thomson is well acquainted with the ground, they may be relied on as accurate. Those at the back of the island are also correct, but the exact outlines of the bases of the hills have not been laid down, no land-survey having yet been made of that part of the island, and dense mangrove and other jungles generally concealing the outlines.

Coral-reefs fringe the south-western shores of Singapore Island and the shores of the islands to the south, the northern coasts of Batam, with the adjacent islands, and a group of small islands in Tilo' Subong. Isolated patches of coral occur also on the eastern coasts of Singapore.

Explanation of Geographical Affixes used in the Memoir and Map.

Suŋge or Suŋgěi, <i>river, stream.</i>	Tanjong, <i>promontory.</i>
Silat, <i>strait.</i>	Pulāu (Pulo), <i>island.</i>
Tilo', <i>bay.</i>	Malang, <i>rock above water.</i>
Tana, <i>land.</i>	Batu, <i>stone, rock.</i>
Gunong, <i>mountain.</i>	Padang, <i>plain.</i>
Bukit, <i>hill.</i>	Kampong, <i>a collection of houses.</i>
Pĕrmatang, <i>a long elevation.</i>	Ayĕr, <i>water, stream.</i>
Busong, <i>a spit of sand.</i>	Paya, <i>marsh.</i>

[A few small sections of the rocks of Singapore and the neighbouring islands are given in Journ. Ind. Archip. vol. i., in illustration of Col. Low's "Notes on the Geological Features of Singapore."]

GEOLOGICAL MAP
OF THE DISTRICT OF THE
STRAITS OF SINGAPORE
and the
JOHŌR ARCHIPELAGO

By J. R. Logan, Esq. F. G. S.