dangerous. It has, at times, an irritant effect on the kidney, setting up an acute toxic nephritis in an organ so often the seat of chronic parenchymatous changes. This effect was not common, but the accident was so serious as to warrant our return to thymol as a much safer vermicide. Thymol on rare occasions may irritate the kidney, but its effect is much less severe. It may produce, infrequently, more or less severe collapse. The chief objection to its use was its irritating effect on the bowel and several times, we believe that enterocolitis has been initiated by its administration.

Eucalyptol is now being studied by the present commission, but their report has not yet been made. Both thymol and betanaphthol were administered in the same manner except that one-half as much betanaphthol was used at a dose as thymol. Betanaphthol in larger doses did not seem to give proportional results.

The patient was required to abstain from solid food during the day and at night 25 grams of sodium sulphate was given. The next morning, providing free purgation from the salts was secured, after a glass of hot milk or on a fasting stomach, one-half of the anthelmintic was given and repeated in two hours, followed, about noon, by another purge which may be omitted if the bowels are moving freely. This procedure was repeated each week until ova disappeared from the feces.

We have given thousands of doses of thymol without having known of a case of poisoning from it, notwithstanding that at times, we knew that, contrary to our orders, alcohol in some form had been taken to relieve the dizziness and other disagreeable effects sometimes produced by thymol in large doses.

In choosing the anthelmintic we have to be governed somewhat by the circumstances of the case, condition of the patient, and so forth, but, all things being equal, we decidedly prefer thymol for general use. When it is not possible to use it, betanaphthol is an excellent substitute. The taking of either is unpleasant and, naturally, we want to use that one which the least number of doses need be taken to obtain the greatest result.

Our thanks are due to Drs. Carroll, Russell and Gray of the laboratory of the Army Medical Museum and Dr. Rosenau of the Hygienic Laboratory, Washington, D. C., for many courtesies extended us in our work on the morbid anatomy of uncinariasis and acknowledgment is made of painstaking services rendered by First Class Sergeants Neat and Britton, Hospital Corps, U. S. Army.

EXPERIMENTAL ANEMIA.*

C. H. BUNTING, M.D.

Professor of Pathology, University of Virginia.
CHARLOTTESVILLE, VA.

When one considers how incomplete is our knowledge of the anatomy and physiology of the hematopoietic system it is perhaps not remarkable that the diseases of that system should have remained so long shrouded in obscurity. Although the clinical pictures of the anemias and leukemias are sharply defined, their pathology still offers many problems. When in his classical description of pernicious anemia Addison noted that it occurred "without any discoverable cause," he stated the first and greatest problem of that disease and one which a half-century of investigation has not been able to solve, al-

though by clinical, pathologic and experimental studies it has been able to narrow the field of search somewhat. However, this, the etiology of the disease, still remains its great problem.

The second problem arose at a time when the clinical picture, especially the blood picture of this anemia described by Addison, had not been sharply defined, and it arose from the description by Biermer and many others of progressive and pernicious anemias following many well-defined causes, causes almost too numerous to catalogue, and causes which in general are followed by an anemia of the so-called secondary type. This difficulty was partially cleared up by the clinicians who showed that the Addison anemia had a distinctive blood picture. It has left still unsettled, however, the question as to whether or not at some stage secondary anemias of varied causation may not present the blood picture of pernicious anemia.

So the second problem may be stated as the relation of the primary and secondary anemias. To express it more fully, if pernicious anemia be not always due to the same cause, but to many causes, and among them to those that at times produce anemias of the secondary type, what is the factor which determines the type of blood picture?

The third problem is one that arose from the pathologic investigations of the disease. Cohnheim discovered the characteristic bone-marrow lesion of the disease, and it was later thoroughly studied by Ehrlich and others, who emphasized its greatly increased megaloblastic content as pathognomonic of the disease. As a result of these studies many have considered the bone-marrow changes as the primary lesion in the disease and accountable for the blood changes and other lesions.

Exclusive of the bone-marrow changes the pathologic study has revealed in many cases no lesions that could not be considered as secondary to the anemia, with the exception of the lesion emphasized chiefly by Quincke and Hunter. This lesion is the excessive deposition of iron-containing pigment in the liver and kidneys especially, though occurring also in the spleen and other organs. These deposits were interpreted as the result of the action of a hemolytic substance on the red blood cells and formed the basis of the theory accepted by many that the disease is essentially a hemolytic process and that hemolysis occurring in the circulation is responsible for the other lesions of the disease.

This theory received additional support from the facts that infection with the intestinal parasite, *Bothriocephalus latus*, gives the typical picture of pernicious anemia and that an extract of the worm has been shown experimentally to produce hemolysis in dogs. Yet it became evident that hemolysis alone is not adequate to explain the disease, for hemolysis with deposition of blood pigment is a factor in secondary anemias not caused by hemorrhage.

The third problem then concerns the pathology and the pathogenesis of the disease, the value of the claims as to the primary lesion. Is there a primary bone-marrow lesion or is there an original hemolysis with subsequent changes in the marrow?

Briefly, then, the problems of pernicious anemia concern its entity as a disease and the establishment of the blood picture, its relation to other types of anemia, its pathogenesis and its etiology. All of these are not subject to experimental investigation. The establishment of the blood picture and of the disease as an entity has been done by clinicians and pathologic anatomists. A

* Read in the Section on Practice of Medicine of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

definite blood picture characteristic of the disease has been made out, and it has been determined that there is a disease corresponding to Addison's description with this definite blood picture and with pathologically a characteristic bone-marrow lesion, signs of hemolysis and no lesion otherwise, apparently adequate to explain the anemia.

For the etiology one must look to further clinical and pathologic study. On the other hand, the relation to secondary anemia, and the pathogenesis of the disease lend themselves to experimental investigation. The work of Quincke, Hunter and others in establishing the fact that hemolysis is a large factor in the production of the anemia furnished the lead along which profitable experimental work has been done. In this work the extensive study of hemolytic processes, stimulated by the Ehrlich hypothesis, has been of importance.

In the first place, it has been shown conclusively that the blood picture of pernicious anemia can be produced experimentally in such animals as the dog and the rabbit, the animals used most extensively.¹ This picture simulates the picture in human blood as closely as any pathologic picture produced in animals simulates the picture in man produced by the same etiologic agent. The blood picture in rabbits shows the nucleated red cells both megaloblastic and normoblastic, often occurring in showers. It shows further the characteristic changes in non-nucleated red cells: a variation in size, with the predominant cells above the average normal diameter; a certain amount of poikilocytosis, polychromatophilia and in many cases a granular basophilia and even an increased color index (Tallquist). Even the reduced white cell count with a relative increase of the large mononuclear variety may occur. In short, practically none of the features of the blood picture of pernicious anemia is lacking.

Furthermore, a condition of the marrow agreeing with that found in human cases may be produced—a marrow in which there is a predominance of the megaloblastic type of nucleated red cells and of cells of the large lymphocyte type. There seems to be no doubt, therefore, that one can thus produce the picture of pernicious anemia experimentally in animals.

As to the manner in which this picture can be produced it is of interest, as it bears somewhat on the possibility of a multiplicity of causes in the production of pernicious anemia in man, that it may be produced by the injection of a number of substances—but substances which have a common characteristic and that is, that they are hemolytic for the corpuscles of the experimental animal.

I have used artificially produced hemolytic sera, but chiefly ricin and saponin. Others have used pyridin, pyrogallol and other toxins. With large doses of ricin or saponin, injected intravenously and repeated at intervals, one may produce a progressive anemia with the blood picture described to the point at which the animal establishes a tolerance to the poison. The animals react with machine-like regularity to the dose of poison, with a sharp nucleated red cell reaction and a fall in the number of red cells, with a gradual assumption by the non-nucleated red cells of the abnormalities noted.

This reaction, the blood picture and the marrow picture are entirely different from those of a secondary

anemia produced in animals by hemorrhage. In these latter cases, with repeated hemorrhage and a constant fall in the red cells, there is no sharp nucleated red cell crisis; megaloblasts are not found in the circulation and normoblasts but occasionally, after the first reaction. The color-index falls rapidly, the corpuscles becoming extremely pale, and the marrow has the characteristic picture seen in cases of secondary anemia where the nucleated cells of the normoblastic type are the predominant cells. It is an entirely different picture. While this picture of pernicious anemia may be produced by several hemolytic substances it is only produced by them if they are injected into the circulation in relatively large doses, which is of importance in considering the pathogenesis of the disease.

If these hemolytic substances are injected, not intravenously but subcutaneously or intraperitoneally, the result is entirely different. The hemolysis goes on; deposits of blood pigment are found in the spleen; there is a reduction in the red cell count, but the blood picture and the marrow picture are different unless very large doses are used. The sharp nucleated red cell reaction in the circulating blood is not seen, the fall in the number of red cells is less sharp, and the marrow shows the picture of a secondary anemia with the great predominance of normoblastic cells so characteristic.

Briefly, by the injection of a large dose of hemolysin directly into the veins, one gets the primary anemia blood picture; by the injection of the same dose, or even larger doses subcutaneously, when absorption into the circulation is slow one gets the picture of a secondary anemia and a similar difference occurs in the marrow pictures.

The explanation of this difference lies, it seems to me, in the appearance of the marrow of animals dying after a recent dose of toxin. Where the toxin has been injected intravenously in sufficient doses it is not entirely saturated by red cells before it reaches the marrow and one finds as a result signs of direct injury to the marrow, in extensive necrosis of cells and in injury to the capillary endothelium resulting in minute hemorrhages. That there is hemolytic destruction of nucleated red cells is shown by naked pyknotic red cell nuclei found both in the marrow and in the circulating blood, and also larger cells in the marrow show pyknotic and fragmented nuclei. Where the injection has been subcutaneous, unless the dose of toxin has been extreme, the injury to the marrow is very slight and may even not appear.

Herein lies the explanation, I believe, of the relation of anemias of the pernicious type to those of the secondary type, due to causes other than hemorrhage. That is, in a disease like tuberculosis, in addition to the general malnutrition, there is a constant but slow absorption into the circulation of substances destructive to red blood cells. This absorption, however, is so gradual that the toxin is completely anchored by circulating red cells, causes their destruction or injury as shown by the splenic accumulation of blood pigment, but does not reach the marrow in an uncombined state. As a result the marrow escapes injury.

As far as the marrow is concerned, the lesion is the equivalent of a hemorrhage. It is uninjured and is able to repair the blood lesion in a normal way in as far as the general nutrition of the patient will permit. On the other hand, in pernicious anemia at some periods large quantities of a hemolytic poison gain entrance to the circulation, are not completely combined with circu-

1. The experiments forming the basis of this paper will be found in detail in the following references: *Etiology and Pathogenesis of Pernicious Anemia*, Johns Hopkins Hosp. Bull., 1905, xvi, 222; *Experimental Anemia in the Rabbit*, Jour. Exp. Med., 1906, viii, 625.

lating red cells and reach the marrow. There they cause destruction, not only of mature red cells, but of nucleated red cells and the function of the marrow is impaired, leading to a faulty response to the call made on it to repair the lesion—and the blood picture of pernicious anemia is the result.

It seems possible in this way to explain why in certain severe secondary anemias the picture of pernicious anemia is seen in the blood. One need only postulate that in a progressing anemia there comes a time when the pathologic process, the tuberculosis for example, also progresses more rapidly, more toxin is produced to be absorbed into the circulation, there are fewer red blood cells to neutralize it, and enough of it reaches the marrow uncombined to cause an injury to the marrow cells—and then the pernicious type of blood picture results.

Why one finds this blood picture as a result of marrow injury seems to depend largely on the anatomic grouping of the cells in the marrow. As I have emphasized in previous papers, the cells seem to be so arranged in the marrow that by the multiplication of the parent cells in the erythrogenetic and leukogenetic centers the young cells as they mature are gradually pushed toward the periphery of the group and thus the mature differentiated cells, the polymorphonuclear leucocytes or the non-nucleated erythrocytes are found on the periphery close to the marrow capillaries. These mature cells are the reserve stock from which the circulating blood is supplied to fill in deficiencies caused in the normal wear and tear of the circulation, and normally the marrow cells proliferate only rapidly enough to keep this supply at par.

Let there be an unusual demand, as for leucocytes in an acute inflammation, and one finds that at first the marrow is cleared of all its mature polymorphonuclear cells, and myelocytes form practically the entire white cell count in the marrow. Mitosis and maturation of the products of division, however, soon produce an abundance, usually a superabundance, of polymorphous cells, and the demand is supplied. Occasionally, as in acute lobar pneumonia, the great demand for leucocytes may cause the marrow to react with immature white cells, myelocytes—which are often found in small percentage just before the crisis in that disease.

A similar story is that of the red cells. In a hemorrhage of moderate severity the marrow responds with its mature red cells and proliferation of nucleated cells takes place to supply the demand. Commonly the demand exceeds the supply of mature red cells and normoblasts are contributed to the circulating blood, but seldom in great numbers. Megaloblasts only extremely rarely are found in the stream.

This is the reaction in a hemorrhage or secondary anemia, and if the cause be withdrawn the blood count gradually returns to the normal as the result of this reaction of an unimpaired marrow. In man one misses a hypererythrocytosis corresponding to the leucocytosis which expresses the pathologic reaction in excess. This probably is due to the fact that serum withdrawals from the tissue keep the red cells diluted to the normal number, even though they be produced in excess. In an animal like the rabbit, with dry tissues, a reaction in excess may be obtained, with high red cell counts in the circulation, and then a gradual return to the normal.

Now in the cases with marrow injury we have an entirely different reaction, for the reason, it appears, that the toxin reaches first these mature red blood cells and

the normoblasts and causes their destruction in part. In its first response the marrow throws into the circulating blood the most peripheral and most mature cells left, and these are often normoblasts and intermediates which appear in the circulation in a shower often aggregating thousands per cubic millimeter.

If the injury be more severe, intermediate nucleated red cells and even megaloblasts appear in the circulation as the other cells have been destroyed. This, however, is ruinous to the proliferative power of the marrow; cells capable of mitosis have been lost to it. As a result, with this continued demand for red cells, there is a hasty transition from megaloblast to erythrocyte—without the orderly progression, through intermediate and normoblastic stages. The resulting cell is thus larger than normal, bears more hemoglobin and possibly is an attempt at compensation—an attempt to make quickly a cell which shall be the equivalent of more than a single normal cell. The distorted forms are expression also of this faulty hematopoiesis.

The nucleated red blood cell crises or showers so widely considered as signs of regeneration on the part of the marrow can not be so considered from the results of experimental anemias. They do not occur during recovery from hemorrhage or injury, but only immediately following direct injury to the marrow by toxins, and it would appear must be interpreted as the result of that injury. The marrow deprived of its mature red cells responds with the nearest substitute it has—the immature nucleated red cells. The greater the injury to the marrow, so much less the maturity of the cells which it can offer to supply the deficiency, so after severe or repeated injury the number of megaloblasts which are found in the circulation is increased and they are in a way an index of the gravity of the lesion.

On the evidence then of experimental work and of pathologic study it seems justifiable to conclude as to the pathogenesis of pernicious anemia that it is due to the absorption into the circulation of a substance or substances of hemolytic power toward human red blood cells, and in such quantities that it produces not only destruction of circulating red blood cells, but also destruction of, and injury to, cells of the erythrogenetic groups in the bone-marrow. This results in a defective and atypical blood formation, some phases of which may be considered as an attempt at compensation, others as a sign of injury.

One may conclude further that in secondary anemias there is a loss of red blood cells or a destruction of red blood cells by toxins. In either case the marrow is unaffected except in being stimulated to respond in a normal manner with the orderly production of non-nucleated red blood cells, and that its proliferative power is not impaired. At some point in the disease, however, the amount of toxin may be so increased that the marrow becomes injured and the blood picture becomes of the pernicious anemia type.

As to the source and nature of this poison in pernicious anemia, there is little to add. In one type only, the *Bothriocephalus* infections, has it been determined. In those cases the source is the intestine, the poison a hemolytic substance derived from the worm. Such cases may recover, with a normal blood picture, after removal of the parasite.

In the great majority of cases no *Bothriocephalus* infection exists. In these one must assume another cause. There is a tendency, however, on the basis of clinical and pathologic findings, to regard the gastrointestinal

tract as the source of intoxication. From the clinical side gastrointestinal symptoms in pernicious anemia patients are frequent and often severe, the persistent diarrhea with the general weakness furnishing the reason that brings the patient to the doctor.

From a pathologic standpoint a chronic gastritis is a frequent lesion found in the patients, and Hunter has called attention to the frequency with which oral sepsis and in particular a chronic glossitis are found in these cases. He considers that the infection of the mouth extending to lower portions of the gastrointestinal tract is the cause of the disease.

There seems to me much in this theory of Hunter's, and while I can quote no large series of personal investigations of the oral cavity, I would like to record that in two recent cases of pernicious anemia in which the oral cavity was examined there were found extensive dental caries and pyorrhea alveolaris, and also at the base of the tongue a chronic inflammatory process, in one of the cases, with shallow superficial ulceration. In the other case there was no ulceration but a degeneration and numerical atrophy of the tongue muscle and a deep sclerosis with round cell infiltration and hyperplasia of the mucous glands.

In both cases there was also a chronic gastritis. There were no other lesions found to which the anemia might be attributed. I hesitate to lay much stress on the importance of this chronic glossitis, as I am unable to furnish control statistics as to the microscopic condition of the tongue in cases dying of other diseases than anemia. I have, however, failed to observe the glossitis grossly.

If the lesion be due to infection of the gastrointestinal tract the clinical history of partial or almost complete recovery of the patient with subsequent relapse, or repeated alternation of these states, would suggest that the infecting organism was one against which a lasting immunity was not obtained. It may be one which possibly leaves the patient even more susceptible—the common history in infections by the organisms of the streptococcus, pneumococcus and gonococcus groups. There is no absolute evidence that the organism does belong to that group, and yet there appears to be no evidence that it does not. As to etiology, we must still say with Addison as to the great number of cases, that it is a disease without any discoverable cause whatever.

[THE DISCUSSION ON THE PAPERS IN THE SYMPOSIUM ON ANEMIA (ARTICLES BY DRs. ASHFORD AND KING, BUNTING, CABOT, AND MELTZER) WILL APPEAR WITH THE TWO ARTICLES LAST MENTIONED.]

Certified Milk.—This term, which has been copyrighted to prevent its misuse and to limit it to products marketed under the direction and sanction of medical milk commissions, had its origin with the Medical Milk Commission of Essex County, New Jersey, which was organized in 1893 to secure a supply of milk which should always conform to a definite standard of purity. There are now 22 such commissions in the United States, chiefly in the eastern states. The original commission has just made its twenty-first report. This commission deals with only one dairyman, who employs in the production of the milk 1,000 acres of land, 515 cows and produces daily 2,900 quarts of certified milk. A new concrete stable has been constructed on the most modern plans. A bacteriologic laboratory is established. It appears from the reports that the objects of the commission are being satisfactorily attained. The report gives a very instructive view of what can be done by private medical supervision to secure the sanitary purity of an important food product.

THE PRINCIPLES OF VACCINE THERAPY.

BACTERIAL INOCULATION UNDER THE GUIDANCE OF THE
OPSONIC INDEX.*

SIR ALAIROTH E. WRIGHT, M.D.
LONDON, ENGLAND.

INTRODUCTION.

I have undertaken to outline to you to-night—so far as I can do so within the limits of a single lecture—the principles of vaccine therapy; i. e., of the treatment of bacterial disease by the inoculation of the corresponding vaccines. Let me preface what I have to say on this subject by reviewing briefly the methods—other than vaccine therapy—which we have to-day at disposal for the treatment of bacterial disease. The following are, I think, the only methods which here come into consideration:

- (1) Treatment by chemical antiseptics.
- (2) Treatment by the extirpation of the obtrusive focus of infection.
- (3) Treatment by the determination of lymph to the focus of infection.
- (4) Serotherapy.
- (5) Expectant treatment.

TREATMENT BY CHEMICAL ANTISEPTICS.

Antiseptics have found in medicine a threefold application. They have been administered internally with a view to checking microbial growth in the blood, or in regions which can be reached only by the channel of the blood. They have been applied locally with a view to holding in check and extinguishing localized bacterial infections. They have been used for the purpose of checking putrefaction in discharges and devitalized tissues. Neither the first nor the third of these applications calls for any discussion. For some time past it has been all but universally recognized that it is futile to attempt to check bacterial growth in the interior of the organism by our present antiseptics, which have a greater affinity for the constituent elements of the body than they have for any bacteria¹. Again, there has never been any doubt as to the possibility of suppressing putrefactive changes by antiseptic irrigations. Attention, therefore, may be concentrated here on the issue as to whether or not the antiseptic applications are effective in holding in check and extinguishing localized bacterial infections.

It is, of course, currently believed that this method of treatment is effective. It is in this faith that the surgeon introduces antiseptics into septic wounds, or, when he happens to be so minded, into abscess cavities. It is in this faith that the physician resorts in the case of pulmonary infections to antiseptic inhalations. And it is in this faith that the dermatologist, gynecologist, laryngologist, aurist, and genitourinary specialist are strenuous in the application of antiseptics each to the particular province of the body which he takes under his special care. It will be profitable for us to collate the facts and to consider whether or not there is in reality any trustworthy basis for the belief that inspires all this practice.

Significant in this connection appears to me the fact that antiseptics are now by general consent abandoned in the treatment of ordinary surgical wounds. Signifi-

*EDITOR'S NOTE: This represents a lecture delivered before the Harvey Society, New York, Oct. 20, 1906. The manuscript, however, was only received July 22, 1907, so it is fair to assume that it embodies the author's latest ideas on the subject.

1. We have a general formula for such antiseptics when we say in technical terms, which perhaps improve a little on those of Ehrlich, that they are more "histotropic" than "parasitotropic."