hyperthyroidism with intestinal manifestations. An adrenalin test gave a positive result on two occasions. The adenoma was removed, and six weeks later the patient had gained 25 pounds, had had no return of the diarrhea or constipation, was sleeping well, and was less nervous; microscopically the adenoma showed abundant mitochondria. The case illustrates the claim of Goetsch that between adenoma of the thyroid and certain types of hyperthyroidism there exists a relation, depending not on the size of the adenoma but upon its activity, which may be measured by the response to the adrenalin test and by the presence of mitochondria in the tumor cells.

McCarrison, R. EFFECT OF FOOD DEFICIENCY AND EXCESSIVE THYROID GLAND. [Ind. J. of Med. Research, January, 1920. J. A. M. A.]

McCarrison's studies have shown that dietaries deficient in vitamins lead to a reduction in size and weight of the thyroid gland and render it susceptible to the noxious action of intestinal bacteria, or of their products, with resultant atrophic and necrotic changes. A scorbutic diet of crushed oats and autoclaved milk may cause in guinea pigs considerable enlargement of the thyroid, in the main, the result of congestion and hemorrhagic infiltration of its tissues. Dietaries containing adequate provision of vitamins, but excessively rich in proteins and fats, induce in the thyroid of pigeons in confinement marked degrees of hyperplasia, the extent of the hyperplasia being largely dependent on the duration of the organs' exposure to the goitrogenous influences induced by the excessive protein and fat content of the food. The addition of onions to a dietary excessively rich in protein and fats, while containing at the same time an abundance of vitamins, markedly retards the development of thyroid hyperplasia, and the tendency to acinar budding in pigeons living in confinement. The beneficial influence of the onions is held to be due in part at least to their action in restraining the growth of putrefactive types of bacteria in the gastro-intestinal tract and in retarding the absorption of their products. It is suggested that succus alii might prove of benefit in restraining the thyroid hyperplasia of exophthalmic goiter. The changes in the parathyroids induced by a diet deficient in vitamins and excessively rich in starch and fat appear to be related in their origin to intestinal anaerobes, the noxious action of which is greatly favored by the defective diet.

Fasano, M. RARE SYMPTOMS IN THYROIDITIS. [Il policlinico, April 26, 1920.]

The case of a married woman of thirty years of age is here reported. During an attack of suppurative thyroiditis tachycardia developed, tremors of the upper limbs, slight exophthalmos, and Moebius's sign, but neither Stellwag's or von Graefe's sign. Rapid disappearance of the symptoms followed evacuation of the abscess.