Original Articles.

A FEW REMARKS ON BLOOD PRESSURE.

BY JAMES MARSH JACKSON, M.D., BOSTON,

Physician to Out-patients, Massachusetts General Hospital.

It is about three years since I first began to experiment with Gaertner's blood-pressure machine, and the results have been so satisfactory that I am led to say a few words on a subject which is just now exciting much interest.

I was very fortunate in getting a tonometer that was fairly accurate, though it has been necessary to test it from time to time with a mercury manometer (also Gaertner's) and to rearrange the dial. Although these instruments (I refer to the spring manometers) are very convenient, I should hardly care to recommend them for general use, as they vary greatly in accuracy, as must all spring manometers. If, however, a tonometer can be found that will stand the test with a mercury manometer, I do not see why it should not be accepted as a standard machine. It is certainly much more convenient and portable.

Whether or not this method of Professor Gaertner is an exact index of the actual blood pressure I am not prepared to say, but I believe that it is a sufficiently accurate measure of tension for all practical purposes, or at least gives us fairly accurate measures for comparison. It does not seem to me that the objection usually raised, namely, that the reaction looked for in the finger is largely dependent on the tension in the capillaries and that this tension is a very variable one, is of much moment. As I have noted it, the blush in the finger tip is almost instantaneous and takes place the instant the pressure on the arteries of the finger is sufficiently relaxed to permit the blood to flow by the obstruction. I do not see why this method is not the same as the Riva Rocci, the former utilizing the smaller, the latter the larger, arteries of the arm.

I have recently been using the Riva Rocci apparatus which Dr. Locke very kindly made for me, and have been comparing the results obtained by these two machines. As a rule the readings have been about the same, but occasionally there has been a wide divergence for some unknown reason, and I am not able to say which has been the more accurate. This is a matter which must be worked out carefully.

There have been several cases of marked arteriosclerosis in which the blood pressure with the Riva Rocci was very much higher than with the Gaertner, and I believe that the latter was nearer right, as the brachials in these cases were so rigid that it took great force to constrict them, whereas the smaller vessels of the finger seem to be less affected by the disease, so that in all cases of arteriosclerosis I shall continue to use the Gaertner.

Just here I should like to suggest that all records of blood pressure should be accompanied by the name of the instrument used, as otherwise some of the high readings in arteriosclerosis with the Riva Rocci might be very misleading.

Last week an interesting case of angina brachialis appeared at the clinic, and an attempt was made to take the blood pressure. As there were no brach-

ial or ulnar arteries to be felt, and as the radial was so feeble that it could scarcely be perceived and disappeared entirely during the attack of angina, we were obliged to give up the Riva Rocci and use the Gaertner. It was interesting to note that the blood pressure was lower during the angina, which is the reverse of what happened in Dr. Walton's case of intermittent claudication published last year.

Every one who intends to take blood pressures should be equipped with a mercury manometer of one form or another to start with. Then he should have the Riva Rocci armlet and the Gaertner finger ring, both of which can be used with the same machine. The advantage of this will be seen in what was said of the arteriosclerosis cases, and the same applies in patients with very fat arms or in whom the radial pulses for some reason are hard to find.

In order to have the readings taken by different observers fairly uniform, the following rules should be observed: With the Riva Rocci method the blood pressure should be taken with the patient in recumbent position; with the Gaertner method the patient may either lie down or sit up, but the finger should always be on a level with the heart. The patient should be under no mental excitement at the time of the reading, as otherwise an elevation of the blood pressure will be found. Two readings at least should always be taken and if they agree this may be considered accurate, but if there is a marked difference then there has been some mistake in the experiment, and other readings will have to be taken.

I should like to speak of a few observations which I have made in the course of my clinical work. In a series of cases of exophthalmic goiter which I have followed for over two years, and in which I used at Dr. F. C. Shattuck's suggestion bromide of quinine, I have found no variation in the blood pressure of any account. These cases (all women) have nearly all had a pressure ranging from 120 to 160 (Gaertner), that is, they were above the normal, but in none of the cases have I found a marked diminution of the pressure with improvement in the symptoms. None of these cases are absolutely well, but even those without symptoms still have a rise of blood pressure.

In one case of small cystic goiter the pressure has ranged about 160, whereas in a case of large cystic goiter which reached to the second rib the pressure was below normal.

It might also be interesting to note the range of pressure in the cases of chronic nephritis in which the operation of decapsulation of both kidneys was done. In 5 out of 6 cases which I had the opportunity of following there was a rise of pressure even with marked improvement in the symptoms. In one the pressure rose from 125 at time of operation to 210 at the end of two weeks. At this point digitalis was given on account of a failing pulse, and the tension soon fell to 180, which was contrary to my expectation, as I thought that the digitalis would maintain the high tension. As improvement set in at the end of another month the pressure again rose to 220. Three months after the operation the tension was 190, and the patient was in very good condition. The albumen had gradually diminished in amount from $\frac{1}{2}\%$ to a trace. This was a case of chronic diffuse nephritis.

In a case of chronic glomerulo-nephritis the tension rose from 100 at time of operation to 150 in a month, although there had been a great improvement of the symptoms and the albumen had gone from $\frac{1}{2}$ of 1% to $\frac{1}{10}$ of 1% in this time. One showed a drop in the pressure and that of only 20 mm., or from 170 to 150 in four weeks. These observations were all made with the Gaertner.

As an aid in diagnosis the blood pressure is fast taking an important place. We know, for instance, in arteriosclerosis the blood pressure is high; in chronic nephritis, especially of the interstitial type, the pressure runs very high, and in a way the amount of tension is a good index to the extent of the disease.

I remember one case in this connection which is of interest. A young girl came to my office complaining of headache and some dimness of vision, and on examination I found considerable enlargement of the heart and beginning retinitis, although she was still at work and made but little complaint. No edema was present, and examination of the urine showed no trace of albumen and the sediment was also normal. The blood pressure was 250, or as high as could be measured by the tonometer. I sent her at once to the hospital, as I could not explain her condition. Nothing more was found until a few days later, when a slight hyperemia was apparent in the urine. She died suddenly after a week in the hospital, and it was found that she had a tuberculous kidney with occluded ureter on one side and a normal but somewhat hypertrophied kidney on the other. Although this patient did not appear very ill until the last day or two, the blood pressure was a very significant sign.

Thus far the best use to which I have put the instrument has been in the demonstration of tension to the students. This has always been a hard subject to teach, but with a direct measure before them they have been much keener in their interest to perfect themselves, and the results have been very satisfactory. I have always taught them that the blood pressure in a young healthy man ranges from 100 to 130 mm. of mercury, but I have frequently found it as high as 150 in young men who, so far as I could ascertain, were perfectly sound. These same cases, however, are liable to considerable variation, and I think this ought always to be taken into consideration in all our experiments.

In young women the pressure is decidedly lower, and ranges from 90 to 110, so that while a pressure of 140 to 150 may be normal for a man it is abnormal for a woman. Elderly people almost always, so far as I have observed, have an elevated pressure, due for the most part to changes in the arteries, so one should not be disturbed in finding in a person of fifty years a pressure of 175.

Low pressures are uncommon, except in cases of shock and collapse, and it is with high pressures that we have to deal with principally. Pressures of 200 are not infrequently found, and should always be considered more or less dangerous, while pressures of 250 are very dangerous and are only seen in grave cases. One hears of much higher readings, but I do not believe that they are always accurate except, perhaps, in some bad head injuries. In the last year five of my patients with pressures of over 190 have died of apoplexy, and I now make it a rule

to warn the family of a patient with a tension above 190.

I regret very much that I did not have an instrument of high registering power in a case of apoplexy which occurred last year in my clinic, for I am sure the pressure must have been very high. The patient, a woman of about sixty, with marked arteriosclerosis and interstitial nephritis, had been coming to the clinic for some time, and I had made several readings of the blood pressure, which ranged about 210 Gaertner. One day she came in complaining of severe headache, and as I was taking her pressure I found it to be higher than my machine would register (250), and while I was making a third attempt to take it, to be sure that I was correct, she fell to the floor unconscious, and died within a few hours of cerebral hemorrhage.

In closing, I wish to urge on all who are doing clinical work, or who are teaching, the importance of this new measure of tension. There is a great deal to be learned, and there is no question but that it will be of great diagnostic value as soon as we arrive at more definite conclusions.

INTESTINAL OBSTRUCTION BELOW THE ILEO-CECAL JUNCTION.¹

BY THOMAS H. MANLEY, PH.D., M.D., NEW YORK CITY.

VARIOUS types of intestinal obstruction call for active surgical interference after tentative therapy fails.

During the past twenty-five years remarkable progress has been made in the direction of enlarging our knowledge of the diverse pathological characters of disease processes situated in the large intestine, from its joint with the ileum to the point where it passes externally at the anus. Moreover, aggressive and ingenious surgery of the intestine—comparatively a modern creation—has wrought a radical revolution in therapy here, and in this area of the alimentary canal it promises to accomplish more, in what was formerly a group of most distressing maladies.

SURGICAL EXPERIMENTS IN PHYSIOLOGICAL PROBLEMS.

Modern surgery points to the necessity of a modified conception of function in digestion, a revision of physiological teaching. It has recently been demonstrated that the complete excision of the colon or the stomach is not incompatible with full nutrition of the body; those who survived the loss of their stomachs even gained in weight and vigor. The colon is a long, sacculated funnel for the reception of the residue of digestion; it was formerly supposed to serve only as a reservoir for the *débris* and bye-products of alimentation.

From early infancy until late in life the terminal coil is more frequently the seat of functional derangement and organic disease than all other areas of the alimentary canal, from the pylorus down, and yet, until a recent period, only its rectal end, when diseased, was considered as worthy of exhaustive study by pathologists, and was but indifferently treated by practitioners.

¹ Read before the Hartford Medical Society, Hartford, Conn., Dec. 22, 1902.