Schreiben des Herrn Prof. Dr. C. H. F. Peters, Directors der Sternwarte zu Clinton,
an den Herausgeber.

Wie durch transatlantischen Telegraphen nach Europa berichtet worden, wurde am 24. Mai ein kleiner Planet aufgefunden, dessen Grösse ich auf etwa 11.3

schätze. Die folgenden Beobachtungen sind bis dahin gelungen, wobei die Vergleichsterne aus *Argelander's* südlichen Zonen entnommen sind.

1873	Ham. Coll. M. Zt.	α (131)	δ (131)	$\log(p''\Delta)$	Anz. d. Vgl.	Vgl. Stern
Mai 24	15h 17m 56s	16h 16m 10s.90	— 21° 17' 49" 8	0.720 0 859	2	Oe. Arg. 15557—60
25	13 13 5	15 14.03	18 1.4	0.345 0.898	10	"
27	13 55 44	13 6.97	18 23.4	0.566 0.885	6	"
"	14 30 41	13 5.06	18 25.9	0.658 0.873	3	Oe. Arg. 15490—1
30	13 26 33	10 1.46	18 49.1	0.518 0.890	10	"
31	13 17 30	9 0.27	18 56.6	0.502 0.891	6	"
"	13 43 7	8 59.01	18 56.4	0.587 0.883	6	Oe. Arg. 15432.5.6
Juni 1	12 50 6	16 8 0.36	— 21 19 1.7	0.407 0.896	10	"

Der periodische Comet von Tempel wurde ohne Schwierigkeit aufgefunden, und beobachtet in der ersten heiteren Nacht nach Empfang einer von Herrn *Hind* gütigst übersandten Ephemeride. Wegen Mangel an

passenden Vergleichsternen habe ich indess blos die folgenden zwei Beobachtungen gemacht, die erste mit dem Ringmikrometer, die zweite mit erleuchteten Fäden am Fadenmicrometer.

1873	M. Zt.	α //	δ //	Anz. d. Vgl.	Vgl. Stern
Mai 23	13h 25m 35s	16h 29m 18s.06	— 16° 26' 28" 0	8	Oe. Arg. 15776—7
"	13 25 52	a — 30.99	a + 34.1	16	a
30	14 43 18	16 24 52.58	— 17 39 20.5	6	Oe. Arg. 15661

Der genäherte Ort des Vergleichsterns a 11. Grösse ist für 1873.0:

16h 29m 47s.3; — 16° 26' 53".

Die Declinationsdifferenz gestattete blos einseitige Vergleichung mit Oe. Arg. 15776—7 (Comet Süd,

Stern Nord, äusserer und innerer Ring); dagegen sind die Vergleichungen mit Stern a sowohl Süd als Nord.

Litchfield Obs. of Hamilton College,

Clinton, N.-Y., 1873, Juni 3.

C. H. F. Peters.

2

Positions of the principal Stars derived from observations made at the U. S. Naval
Observatory at Washington, by Prof. M. Yarnall in the years 1853 to 1860.

a. The second appendix to the Washington observations of 1871 contains the results of the observations with the Transit instrument and Mural Circle in the years 1853 to 1860 inclusive. In these observations the right ascensions given in the British Nautical Almanac for 1860 were assumed as the standard of reference. The correction and rate of the clock were determined for each night by combining the observations of clock stars into groups, and the right ascensions are therefore strictly differential. From 200 observations of stars near the equator I find the probable error of a single right ascension to be $\pm 0s.034$. It was thought however not worth while to compute the probable errors of the right ascensions of the principal stars, since these errors would have but little significance. In the following table the second column gives the observed right ascen-

sion of the star, the third column the number of observations, and the fourth the difference from the Berliner Jahrbuch 1860.

b. The declinations of the stars were referred directly to observations of the nadir point and are absolute determinations, the latitude being assumed as $+38^{\circ}53'39''25$. The errors of graduation of the Mural Circle have never been completely investigated, but from partial examinations in 1845 and 1846 it appeared that errors arising from defects in the form of the Circle and from errors of graduation are nearly eliminated from the mean of six microscopes. The position of the Circle was changed but once during these observations, viz: at the beginning of 1858. From 912 observations the average probable error of a single declination is $\pm 0''49$. In transferring the declinations from 1850 to 1860 the yearly variations