The Nature of Matter—I.*

This is Not a Universe of Dead Atoms, But of Active Energy

By J. C. Whitehorn, Doane College, Crete, Neb.

FROM the time when men began to wonder at the world and ponder its problems they have puzzled over the nature of the universe. Sensing an external world of objects and feeling the vital necessity of dealing with these objects, they have sought for an understanding of the universe which would give them a mastery over it. And in their more contemplative moments they have tried to attain to a more comprehensive view which would not only be serviceable in handling objects but would also give a true presentation of the reality of the universe. Through the whole range of solutions offered one can find an ever present attempt to conceive the universe in some unitary and comprehensive manner. Some have looked upon it as a chance collection of dead atoms propelled by blind force; others as pure idea or thought. Between these two extremes there have been all sorts of gradations and compromises.

The unreflective attitude of the average man, which may be called the common sense view, is that there is a certain substance or matter back of the phenomena of nature. This primary matter has certain properties such as color, odor, taste, etc. The phenomena of nature consist of the activities of this matter. Common sense, then, postulates three planes or kinds of reality, (1) a substratum of matter or substance, (2) the properties inherent in this substance, and (3) the activities of this substance.

This naïve solution is, however, quite unsatisfactory. In the first place, it is practically impossible to draw a fast line between activities and properties. For example, luminosity was formerly thought to be a property of matter, now it is considered an activity of matter. Activities and properties merge into one another until we can find no satisfactory line of demarcation.

In the second place, no one ever saw, heard, felt, tasted or smelled "matter." All of our knowledge of objects is a knowledge of properties, not of a "matter" or "substance." An object, as known to us, is simply a self-subsistent organization of properties. The idea of a hard and changeless block of reality which serves as a support to these properties is a purely metaphysical assumption.

And yet, even though there is no logical necessity, we feel a psychological need for the idea of a matter or substance to which we can *fasten* the qualities, as it were. The basis for this psychological need can probably be found in those universal and stable properties, extension and mass. We see about us objects undergoing various transformations of color, odor, density, etc. Most of the properties seem to be variable and changing, and sometimes absent altogether. But there are certain properties which are universally present in all objects, under all conditions and through all changes. These universal properties are mass and extension. Their persistency and universality indicate a "something" which endures and is constant through the changes. Hence, there has unconsciously grown up in our thought a concept of a stable and universal substance or matter which lies back of and gives support to the varying properties.

If, however, we think soberly and consistently upon the matter we shall come to the conclusion that this is a fault of unreflective thought, which has seized upon the more stable and enduring properties as a support for the variable and changing. Extension and mass are, in fact, merely those properties which are more permanent and universal. There is no logically necessary implication or assumption of any material entity, as ordinarily understood.

Let no one think, however, that by this statement we deny the existence of external and objective reality. We are not trying to reduce the world to a phantasmagoria of subjective ideas. We are attempting to get rid of a troublesome metaphysical assumption as to the nature of this objective reality. Our purpose is to ascertain as accurately as possible the true nature of reality, and more particularly of that phase of objective reality commonly known as matter.

We shall strive to conduct this investigation in the genuine spirit of a seeker after truth and therefore we shall consider of primary importance what science has to say upon the subject. And science has much to say. Modern physical research has shed great light upon this problem. The strange and erratic behavior of cathode rays and of the emanations of radio-active substances has greatly modified our conceptions of matter.

We shall take up first a discussion of those phenomena

*Republished from *Popular Astronomy*.

which occur when electricity is conducted through rare gases. If an electric current of high potential be passed through an air-tight glass tube containing gas at atmospheric pressure, an ordinary spark will be produced, just as in the atmosphere. If, however, the gas be somewhat exhausted, so as to create a partial vacuum, the current, which need not now be of high potential, will pass readily through the tube. Its passage will be marked by curious phenomena. The negative electrode (cathode) will be covered by a luminous layer. Next to this is a dark space, the Crookes dark space. Beyond this is a luminous part called the negative glow, and beyond this glow another dark space, the Faraday dark space. Between this and the anode is the positive column of alternately dark and light spaces called striae.

If the gas be still further exhausted to a pressure of about .01 mm. the above phenomena disappear and a bright phosphorescence appears on the sides of the tube. This phosphorescence is produced by radiations of very minute particles issuing in straight lines from the cathode. If a solid body be placed inside the tube so as to intercept the rays, it will cast a well-defined shadow. If the rays be focused upon a body, it may be heated even to incandescence. But what is of much more significance, these particles can be deflected from their normally straight path by means of a magnet. This indicates that the particles are either themselves magnetic or are electrically charged so that their motion sets up lines of magnetic force.

Perrin and later J. J. Thomson have shown that the deflection by a magnet is due to a negative charge of electricity carried by the particles. If the end of a conductor, carefully insulated and protected, be placed near a pencil of cathode rays, an electrometer connected to this conductor will show very little effect. But if the rays be deflected so that the particles fall directly upon the conductor, the electrometer shows a considerable negative charge. Furthermore, if a pencil of cathode rays be passed between two parallel plates which are oppositely charged at a great difference of potential, the beam of rays will be deflected towards the positive plate. These experiments indicate that the cathode particles carry a negative charge of electricity.

This gives us a clue as to the method of finding the velocity of these rays. Measurements have shown that a moving charged particle produces a magnetic field which is equal to the field produced, per unit of its length, by a linear conductor carrying a current equal to eV , where e is the charge on the moving particle and V is its speed. Hence if the speed of a cathode particle is V and its charge e , its magnetic field is eV . If a magnet be brought to bear with field intensity H upon the cathode particle, the particle will be deflected from its path by a force HeV (the product of the two magnetic forces) and will follow a curved path.

Suppose now that two parallel plates be placed on opposite sides of the pencil of rays and a high difference of potential established between them in such a way as to pull the charged particles in the opposite direction from that in which the magnet deflects them. The electric force acting on the particle will then be the product of the charge on the particle times the strength of the electric field. If the latter be X , the force will be eX . Now the magnetic and electric fields, which counteract each other, can be so adjusted that there will be no deflection, at which time the opposing forces are equal. That is, eX equals HeV . Then V equals X/H . Now X and H can be measured and hence V can be calculated. By this method Thomson has found the velocity of cathode particles to be 2.8×10^9 cm. per sec., which is about one-tenth the velocity of light (29.99×10^9). The value for V varies slightly with the potential in the tube, but is fairly constant.

If we remove the electric field and measure the deflection caused by the magnet, we can calculate the radius of the curve of deflection. The force producing this deflection is from the above discussion HeV . The centrifugal force of the particle whose mass we shall call m , moving with speed V in a circle of radius r , is, by mechanics, mV^2/r . Action and reaction are equal, hence $HeV = mV^2/r$. $\therefore V/Hr = e/m$.

Now H can be measured, r can be calculated and V can be determined as shown above. Hence the ratio e/m can be found. Thomson, by this method, obtained a value for e/m of 1.7×10^7 .

The values of V and e/m have thus been shown to be

¹Duff-Carman, p. 387.

²Duff-McClung, p. 495.

interdependent. This indicates an intimate relationship between the velocity, the charge and the mass. Since V , e and m have been practically constant in the above mentioned experiments, we have been unable to determine the nature of their inter-relationships. We must search for a case in which these values are variable in order to find their mutual relations.

Such a case has been found in the radium emanation. The phosphorescent action of the cathode rays led scientists to investigate phosphorescent substances. Investigation of uranium salts and pitch blende ores led to the discovery by Mme. Curie of the element radium. This substance seems to defy the laws of physics. It constantly gives off an enormous amount of energy without seeming to lose weight or be consumed. This energy is given off in the form of radiations. Ernest Rutherford has found that these radiations are composed of three distinct types of rays, called the alpha, beta and gamma rays. Becquerel showed that the beta rays behave in every respect like the cathode rays. They are composed of negatively charged particles having the same relations between mass, speed and charge as do the cathode particles. Becquerel, by practically the same method as Thomson used for cathode particles, determined the velocity and the ratio e/m for the beta rays. The value which he found for e/m does not differ much from the value of e/m found by Thomson for the cathode particles (1.7×10^7). He observed, however, that the velocities varied from 6×10^9 to 2.8×10^{10} cm. per sec.

This variation in velocity led Kaufmann to make more accurate determination of e/m to find whether the value of this ratio varied with the velocity. He made the very important discovery that e/m decreases when the speed increases.³ If we assume that the charge e is constant, then the mass of the beta particle appears to increase with increase of velocity. This is not the old law of mechanics that the momentum varies with the speed, but a new discovery that the mass itself varies with the speed. Now it seems to be quite well established at least for bodies of appreciable size, that the mass of an uncharged particle does not vary with the velocity. This renders it probable that the variation of the mass of a beta particle with the velocity is due to the electric charge.

Indeed, if we stop to reflect, we can find other considerations which would lead us to the conclusion that an electric charge on a moving body will affect its mass. Suppose that a certain force F is applied to an uncharged particle, imparting to it a certain acceleration a . If now the same particle be charged, the force F will be unable to impart to it the same acceleration, because the force not only has to overcome the same inertia as before, but also, by moving the charged particle, sets up lines of magnetic force. By the law of the conservation of energy, this force cannot be created from nothing. Additional force is required to set up these magnetic lines. Hence more force will be required to produce the same acceleration. Now by Newton's second law F equals ma . Then m equals F/a . Hence in the case of the uncharged particle considered above, m equals F/a ; but in the case of the charged particle m equals $(F+f)/a$. In other words, the mass of the particle has been increased by the electric charge which it carries. The increase of mass depends upon the magnetic field set up and this depends, in turn, upon the amount of electric charge and the speed of the moving particle. Therefore the mass of a particle carrying a constant charge is a function of the velocity. It must furthermore be remembered that, when magnetic lines have once been set up, they tend to keep the particle in motion. Hence in either case, whether of negative or positive acceleration, an electric charge on a particle increases its mass.

These theoretical considerations (which, it must be remembered, deal only with moving particles) bear out the empirical findings of Kaufmann. We can therefore conclude that the mass of the beta particle is due, in part at least, to the moving charge. And if in part, then why not altogether? There is scientific material which seems to indicate that the whole mass is due to the charge and its velocity.

J. J. Thomson has worked out mathematical formulae for the mass of a moving charged particle. He has shown that the part of the mass due to the moving charge (the "electrical" mass, as it were,) is $2ae^2/3a$, where e is the charge, a the radius of the particle and a a variable depending upon the velocity. This variable a is taken as unity when V is very small compared to the velocity of light, and becomes larger as V approaches the

³Duff-McClung, p. 507.

velocity of light. The variable α is the ratio of the mass at any given speed to the mass at a certain slow speed (taken as unity). Thomson has calculated the value of α at different velocities. The results are given in the following table, where they are compared with certain experimental results of Kaufmann. The latter has worked out experimentally the values of e/m at different velocities. With these results and taking the mass at a certain slow speed as unity, he has calculated the ratios of the mass at different speeds to this unit mass. His experimental results for the whole mass agree with Thomson's theoretical calculations for the electrical mass within the limits of experimental error, as shown by the following table:⁴

Velocity in cm per second	KAUFMANN	THOMPSON
	Ratio of whole mass at given velocity to whole mass at slow velocity (taken as 1)	α the ratio of elec. mass at given velocity to elec. mass at slow velocity (taken as 1)
2.85×10^{10}	3.09	3.1
2.72×10^{10}	2.43	2.42
2.59×10^{10}	2.04	2.0
2.48×10^{10}	1.83	1.66
2.36×10^{10}	1.65	1.5

This close agreement leads us to conclude that the electrical charge and its motion are sufficient to account for all the mass of the particle. Mass is wholly a product of electrical charge and motion.

The objection will instantly be raised that such a conclusion is untenable, for all are familiar with the fact that uncharged bodies have mass. But an uncharged body is a rather difficult thing to define. It is, in the first place, merely a relative distinction. When we speak of a body as "uncharged" we mean that there is no difference in electrical potential between it and the earth. In the second place the term "uncharged," when applied to a body, means simply that the body, as a system, is at the same potential as the earth. It does not mean that there is no electricity in the body, but that what charges of electricity there may be are balanced so that the system does not show them. Thus copper sulphate may be dissolved and dissociated in water. Under such conditions it most certainly shows evidences of the presence of electrical charges. But these charges are of opposite kinds and are present in such amounts as to exactly balance so that when the CuSO_4 is precipitated the one kind of electricity exactly balances the other and the precipitate gives no evidence of their presence. We say that the precipitate is uncharged, whereas in fact the particles of which the precipitate is composed are fairly alive with charges which will reappear if the copper sulphate is again dissociated. Thus we see that a body might be entirely made up, in its inner constitution, of disembodied charges moving about with great velocity, while the body as a whole showed no signs of this electricity and hence would be "uncharged." If these disembodied charges were continually moving at uniform speed, their masses and consequently the aggregate mass of the whole body would be constant, and would not be affected by movements of the body as a system. Thus the fact that uncharged bodies have mass does not invalidate the hypothesis that mass is due to electric charges in motion, for the mass of such an "uncharged" body may be due to the rapid movement of internal charges which are equally balanced and hence do not give any charge to the body as a system.

From the consideration brought forward above, showing that mass varies with velocity, that an electric charge in motion can produce mass, and that these factors of charge and motion are sufficient to account for all the mass, we are led to the conclusion that mass is not a primary but a secondary quality and is derived from an electric charge in motion. We are now in a position to make the hypothesis that matter is composed altogether of electrical charges, which, by their motion, give to bodies their mass.

This hypothesis has been worked out quite in detail by Thomson, Rutherford and others, and it serves so well to explain the phenomena of physics and chemistry that it is coming to be generally accepted. It is called the corpuscular or electron theory.

I cannot hope within the limits of this paper to undertake a comprehensive discussion. I will have to leave untouched a great mass of interesting facts and a whole body of new theory concerning the bearing of this new idea upon such phenomena as heat, polarization, electrical and thermal conduction, incandescence, etc. I will attempt only to indicate the general outlines of the theory and to point out its application to a few phenomena.

The beta particles and the cathode particles, which are identical in their properties, are taken to be the unit of electrical charge and are called electrons or corpuscles. The atom is thought to be made up of corpuscles. But corpuscles are negative charges and hence repel each

other. There must therefore be a positive charge which holds them together in the atom. The form in which this positive charge occurs in the atom is at present a matter about which we have very little information.

Thomson, in his calculations, considered that the positive charge was spread with uniform density through a sphere of quite large radius, within which the corpuscles were arranged.⁵ Rutherford has shown, however, that the distribution of positive electricity assumed in the Thomson atom was far too diffuse to account for many facts which have later been discovered. According to this view of Rutherford's, the positive electricity is concentrated in a very minute nucleus which is thought to have a radius of $1/1800$ of that of the negative electron.⁶ But this minor disagreement need not divert us from the main consideration, or the difference between these two views is merely a refined and technical distinction. Rutherford's view is more in accordance with observed facts but in order to present Thomson's work as originally developed we will adhere to his theory of a sphere of positive electrification of uniform density.

The positive electricity attracts the corpuscles toward the center of this positive sphere, while their mutual repulsion drives them away from it; when in equilibrium they will be distributed in such a way that the attraction to the positive electrification is just balanced by the mutual repulsion of the corpuscles. When there is only one corpuscle it will of course go to the center of positive electrification; two corpuscles will be in equilibrium on opposite sides of and at equal distances from this center; three, at the corners of an equilateral triangle; four, at the corners of a regular tetrahedron; and six, at the corners of a regular octahedron, the center of each of these configurations being at the center of the positive sphere.

Thomson has been unable to work out the arrangement of more than six corpuscles in three dimensions. He has, however, solved the case where the corpuscles are confined to a plane passing through the center of the positive sphere. According to his calculations, stable equilibrium is assured when the corpuscles are arranged in concentric rings as follows: for numbers of corpuscles from 2 to 5, one ring; from 6 to 16, two rings, 5 to 11 in the outer ring; from 17 to 31, three rings, 11 to 15 in the outer ring; from 32 to 48, four rings, 15 to 17 in the outer ring; from 49 to 69, five rings, 17 to 21 in the outer ring; from 70 to 93, six rings, 21 to 24 in the outer ring. A complete table is given on pages 109-110 of Thomson's "The Corpuscular Theory of Matter" showing the number of rings and the number of corpuscles in each ring for groups of corpuscles ranging in number from 1 to 100.

Let us take for example the arrangement of corpuscles in those configurations which have 20 in the outer ring. The following table gives the number in each ring:⁷

Ring	A	B	C	D	E	F	G	H	I
5th (outer)	20	20	20	20	20	20	20	20	20
4th	16	16	16	17	17	17	17	17	17
3rd	13	13	13	13	13	13	14	14	15
2nd	8	8	9	9	10	10	10	10	10
1st (inner)	2	3	3	3	3	4	4	5	5
Totals	59	60	61	62	63	64	65	66	67

The group of 59 corpuscles shown in Column A has 20 in the outer ring, 16 in the next, 13 in the next, 8 in the next and 2 in the inner ring. Suppose that the charges in this group balance, i.e., that the positive charge equals the sum of the negative charges of the corpuscles. If an additional corpuscle were shot into the group, the 60 corpuscles would then arrange themselves with 20 in the outer ring, 16 in the next, 13 in the next, 8 in the next and 3 in the inner, as in Column B. Since the positive and negative charges at first balanced, this new group in Column B has one excess negative charge. If two corpuscles were shot in, the group would then have two excess negative charges, and the 61 corpuscles would be arranged as shown in Column C. If three were shot in, the group would have three excess negative charges and the 62 corpuscles would be arranged as shown in Column D. Similarly, if 4, 5, 6, 7 or 8 corpuscles were shot in, the groups would have 4, 5, 6, 7, 8 excess negative charges and the 63, 64, 65, 66 or 67 corpuscles would arrange themselves as in Columns E, F, G, H and I. These arrangements would all be stable. If, however, more than 8 corpuscles were shot in, the rearrangement would put 21 in the outer ring, and so would pass outside the group we are considering. Moreover, it would be leading a disproportionate number of corpuscles upon the comparatively small positive charge, which it must be remembered is the same as in Column A. These arrangements then would tend to eject corpuscles and come back to one of the forms with 20 in the outer ring. The stable groups then have 59, 60, 61, 62, 63, 64, 65,

⁴Thomson, p. 103.

⁵Rutherford, pp. 129-130.

⁷Thomson, p. 109.

66 and 67 corpuscles and have respectively 0, 1, 2, 3, 4, 5, 6, 7 and 8 excess negative charges.

If, however, instead of starting with an evenly balanced group of 59 corpuscles, we had started with the group of 67 corpuscles with a positive charge sufficient to balance them and, instead of shooting corpuscles in, had taken them away, the resulting groups would have had 66, 65, 64, 63, 62, 61, 60 and 59 corpuscles, respectively, and would have had excess positive charges of 0, 1, 2, 3, 4, 5, 6, 7, 8.

The group having an excess positive charge of, say, 2 would attract to itself either two groups with an excess negative charge of one each or one group with an excess negative charge of two. That is, the number of excess positive charges measures the attraction for negative groups, and hence may be considered the electro-negative valency. Similarly the number of excess negative charges may be considered the electro-positive valency. We can now give a table showing these results:

Group Letter	A	B	C	D	E	F	G	H	I
No. of Corpuscles	59	60	61	62	63	64	65	66	67
Electro-positive valency obtained by shooting electrons into 59 (A)	0	1	2	3	4	5	6	7	(8)
Electro-negative valency obtained by taking electrons from 67 (I)	(8)	7	6	5	4	3	2	1	0

These groups are arranged according to the number of corpuscles they contain. The valency is seen to be a function of the number of corpuscles. Now, since we have shown that mass is due to electric charges, if there actually exist atoms analogous to these groups, the number of corpuscles of which they are composed would determine their atomic weights. Hence the valency, being a function of the number of electrons, would be a function of the atomic weight. The corpuscular theory then, as worked out by Thomson, leads to the conclusion that valency is a function of atomic weight.

If we were to take up a consideration of those groups of corpuscles having 21 in the outer ring we could in like manner show that the valencies of this group would run from 1 to 8, with an additional group with valency 0 (signifying an inert group) at each end. This would indicate that valency is not only a function of the atomic weight but also a periodic function.

This theoretical deduction from the corpuscular hypothesis is amply confirmed by empirical science. The periodic tables of the elements arranged by Mendeleeff and Meyer show clearly that not only valency but many other properties as well are periodic functions of the atomic weights. The following short table illustrates this fact, with reference to valency:

EL.	He	Li	Be	B	C	N	O	F	Ne	Na	Mg	Al	Si	P	S	Cl	Arg.
At. Wt.	4	7	9.1	11	12	14	16	19	20	23	24.4	27	28	31	32	35.5	39.9
Val.	0	1	2	3	4	5	6	7	0	1	2	3	4	5	6	7	0
Val.	0	7	6	5	4	3	2	1	0	7	6	5	4	3	2	1	0

One can ask for no more beautiful confirmation of any hypothesis than this confirmation of the corpuscular theory by the Periodic Table of the elements.

This confirmation is still more remarkable when we realize that Thomson's calculations have all been based upon the arrangement of corpuscles in a plane. It is to be hoped that three-dimensional arrangements will soon be calculated. If this can be accomplished it will probably give us a still clearer insight into the inner constitution of the atom.

If we return to a consideration of Thomson's groups we shall find that, according to his hypothesis, the atom of one element could be changed into the atom of another element by the loss or gain of one or more corpuscles. This would be equivalent to a transformation of matter. A few years ago such a theory would have been scoffed at and ridiculed. But here again empirical investigation has supported the theory. Ernest Rutherford, through a study of radio-active substances, has found that such transformations are actually taking place.⁸

Radium, as mentioned earlier in this paper, is continually giving off radiations. These radiations are of three kinds, alpha, beta and gamma. The first kind of radiations, the alpha rays, are composed of alpha particles. The alpha particle is very heavy, having four times the atomic weight of hydrogen. It carries two unit positive charges. It is thought to be an atom of helium, carrying twice the ionic charge of hydrogen. The alpha particles furnish the positive electrification for the atom. The beta rays are composed of beta particles, which are simply corpuscles or electrons. The sudden emission of the beta particle or electron from the atom produces an electro-magnetic disturbance in the ether, just as the sudden stopping of a cathode particle produces an X-ray. These electro-magnetic pulses produced by the emission of beta particles are called gamma rays and behave exactly as X-rays.

[TO BE CONTINUED]

⁸Duff-McClung, pp. 506-510.

⁴Thomson, p. 33.