disadvantage without an instrument of this kind to do the tying quickly.

Where the surgeon is operating to remove an impacted gall stone in the cystic or common duct and it is found necessary to split the duct over the site of the stone, the tyer will aid him very much in tightening the sutures when he comes to closing the opening made in this canal.

In the small capillary bleeding hæmorrhoids where the hæmorrhage may be so great as to endanger the life of the patient, the quickest and surest procedure is to ligate the small raspberry like tumor; the tyer here is of much service, making a tedious procedure very quickly and easily performed. This variety of hæmorrhoidal tumors is usually situated high up in the rectum. Allingham has well said, "The secret of the well being of your patient (in ligating internal hæmorrhoids) depends greatly on this tying— a part

of the operation by no means easy as all practical men know, to effect."

Of 29 cases of splenectomy 14 died directly from hæmorrhage and 3 from shock. The difficulty in tying the vessels and the great *shock* produced by dragging on the pedicle can be avoided by the use of the tyer. The application of the instrument has a wide range.

The tyer as the reader will see by the cut has the same arrangement governing the movements of the blades as is used in the Knott's determine dilator. The carriers have automatic threading ends, this permits the ligatures to be quickly placed in position to be tightened as soon as the handles are closed. Between

the blades (see cut) is the loop carrier, which carries the loop down to the exact place where the surgeon desires the knot to be tightened, this carrier holds the loop or knot stationary while the blades are being separated and the knot tightened. The ends of the ligatures are held in the left hand of the surgeon. As soon as the knot is tightened down on the vessel or the suture is so secure that it will hold its place the carrier automatically retracts. The sutures may be tightened even with more force than it is possible for the hands to stand without the ligature cutting the fingers.

These are a few only of its many applications. The tyer is manufactured by Wm. H. Armstrong & Co., Indianapolis, Ind.

Kansas City, Mo., 1024 Walnut St.

SKIN GRAFTING UPON THE CRANIUM.

Presented before the Chicago Medical Society.

BY FREDERICK C. SCHAEFER, M.D.,

OF CHICAGO.

PROFESSOR OF CLINICAL SURGERY, NORTHWESTERN UNIVERSITY MEDICAL COLLEGE; PROFESSOR OF SURGERY, WOMAN'S MEDICAL COLLEGE; SURGEON TO WESLEY HOSPITAL.

The patient I present this evening has been cured for five years. I exhibited her before the American Medical Association in the spring of 1887, when the new scalp covered about three-fourths of the skull. In September, 1887, the case was reported in the International Medical Congress in Washington. At this time there remained an open spot upon the head corresponding to the dimensions of a silver half dollar; a month later this was also covered. The scalp is complete and has been in this condition ever since November, 1887. It may now be appropriately entered upon record as a cure.

Since the subject of skin grafting was first introduced by M. M. Reverdin and Frank Hamilton, innumerable and varied trials have been made to close wounds by this procedure, with variable degrees of success, all over the civilized world, but an extensive search of the medical literature gives very meager information concerning skin grafting upon the cranium.

This young lady, twenty-four years of age, was at work in a laundry under a revolving shaft, July 9, 1886. The shaft was between four and five cm. below a heavy beam, and about twenty-five dm. above the floor; it was making 225 revolutions a minute, as computed by the engineer. The girl was said to have reached up for something on the wall, when the wind blew her hair toward the shaft; instantly she was drawn against the beam and shaft, and fell back to the floor without her scalp.

The narrow space between the beam and shaft doubtless saved her life. She then walked up stairs (the accident occurred in a basement), where I found her sitting in a chair moaning piteously, presenting a most ghastly, heartrending appearance. The entire skull was laid bare, from the margin of the eyelids to the neck. The nasal bones were uncovered and broken, exposing the superior nasal meatuses. The skin of the eyelids was removed from within three mm. of their edges. The border of the wound was traceable from the lower portion of the left external angular process of the frontal bone and margin of the left upper eyelid, downward and backward below the left ear, which was entirely removed; thence the

wound extended across the neck five cm. below the superior curved line of the occipital bone, and could be traced forward through the lower third of the right auricle, to the right external angular process of the frontal bone and margin of the right upper eyelid, across the right lid, nose, and the left lid to the point of commencement. The superior border of the left zygomatic process was broken, exposing the cavity within the process. Every vessel and nerve supplying the scalp was torn off, and the pericranium was lost in three places; one spot, corresponding to

the center of the forehead, was oval in contour and measured 5x7 cm.

The other exposed surfaces of the bones were located on either side of the calvarium, on the corresponding parietal and temporal bones; were elliptical, eight cm. long, six cm. wide. The neck flap of the wound fell away from the muscular structures beneath it, exposing the trapezius muscle almost halfway to the shoulder-blade. The trachelo and sterno-mastoid muscles were also uncovered, the latter to the extent of seven cm. The right ear was

torn through its lower third, and hung at the side of the neck by an isthmus of skin less than five mm. in width. The exposed surface of the wound measured forty-five cm. in length from before backward, and thirty-nine cm. in width near its central portion. The sutures of the cranium were distinctly seen at several places. There were only a few muscular fibres left upon the head—a portion of the temporal muscles on either side, and the levator palpebræ muscles in the orbits. Hæmorrhage was profuse; the tem-

poral, occipital and posterior auricular arteries had to be tied.

I came to the patient within three-fourths of an hour after accident occurred. Controlled hæmorrhage, examined the scalp, wrapped the head up in carbolized gauze. The scalp was thoroughly mangled and covered with fragments of hair on its inner surface, which had to be removed. In order to save time, I sent for my friend and neighbor, Dr. J. D. Skeer, to assist in the immediate treatment of the case. Dipped the scalp in warm carbolized water,

picked off the broken hairs from its inner surface, and cut off the hair from the outer surface, while Dr. Skeer prepared the head for the reception of the scalp. Together we freshened the edges of the wound, stitched the scalp back by continuous suture, and sewed the right ear in its position. Applied iodoform over the entire suture, and coiled small narrow sacks filled with warm sand about the head to keep the scalp warm.

After three days, uncovered the head and cut

through the scalp in several places to let out the pus which had accumulated beneath it. On the seventh day removed all dressings. The scalp was now permeated with gases of decomposition, and baked firmly against the calvarium in many places. Removed it. The right ear was warm and joined at the sutures. During the first three weeks we had little hopes for the patient's life. Her pulse varied from 90 to 130 per minute, while the temperature fluctuated between 100° and 140° F. Was delirious about

three weeks, having lucid intervals; her head had to be propped up with pillows. After the second week her neck became stiff, the muscles were rigid, neck turning to one side. To overcome this tendency, had the bed turned so that it became necessary for the patient to move her neck in the opposite direction. Also had her exercise the neck by rotating and bending it. The optic nerves were extremely sensitive. Vision weak at $\frac{1}{4}$. She complained of noises in

lation tissue, the others near the edge of the wound upon the eyelids, where there was the least suppuration. The head at this time was covered with granulations from the pericranium, excepting in places where the periosteum was lost; these were gradually covering in also, from their margins. The grafts were held in place with fish skin, and the bandages were permitted to remain in position forty-eight hours, when the four grafts upon the vertex presented

both ears. Hearing distance, right ear, $\frac{6}{40}$; left ear, watch C. During the fifth week she left her bed and walked about the room.

Six weeks from the day the accident occurred, I concluded that skin grafting might be most promising of satisfactory results. Suppuration was very

a pink color, and five of the grafts over the eyelids were washed away by the pus. I then attached eighteen grafts, eight on top and ten near the margin, as before. After another forty-eight hours all on the top of the head were pink, the four previously planted were spreading. Those near the margin of the wound remained also, but did not look as good.

free at this time, and the patient was too feeble to permit of any treatment requiring extensive operative procedure. I therefore, on the 15th day of August, took thirteen epidermal grafts from the flexor side of her sister's arm and placed them upon the patient's eyelids four mm. apart. By way of experiment, four of them were placed upon the center of the cranium in direct contact with the granu-

The constant accumulation of pus interfered with the growth of the marginal grafts. I was therefore lured on to place the greater portion of them near the vertex in semicircles, coming forward to meet those which were growing over the lids. October 12, my notes say there is a piece of transparent skin, thirteen cm. long and seven cm. wide, extending from the eyelids backward, besides twenty pieces, varying in size from a ten cent piece to a quarter of a dollar,

noon the sides. On the 15th of October, noticed, that the new skin began to liquefy in spots. October 17, 2 o'clock A.M., a copious hæmorrhage occurred from one of the venous channels between the tables of the right parietal bone; the blood ran forward and to the left side, beneath the front half of the new skin, which was thus destroyed. A few days later the other half was perforated by many small ulcers. In ten days more it also disappeared. Over two months' work was destroyed in as many weeks. The amount of tissue lost would be represented in one piece by an oval measuring fifteen cm. by 11 cm., to produce which 915 grafts were required. Although somewhat discouraged by this disaster, we were not dismayed, but endeavored to retrieve our loss by vigorous grafting.

Having noticed that the skin derived from the epidermis, chiefly, was very thin, immovable and short lived, I resorted to *dermal* grafting, picking up each time, also, a few fibers of subcutaneous connective tissue. Suppuration having greatly diminished, all of them were placed about the circumference of the wound. About sixty-five per cent. lived. During this month, November, two irregular pieces of bone measuring 3x2 cm., were removed from the outer table, where the pericranium was gone.

December 1st, there was an irregular band of skin around the entire head, six cm. wide in front, and four cm. wide at the back and sides. The last two series of grafts did not thrive. Up to this time heterodermal grafts only were brought into requisition from her sister and three brothers. As the patient was now able to walk out of doors, she gained in strength, and the thought occurred to us—Dr. Skeer, who called to see her at this time, and myself—that tissue taken from her own person might do better than the heterodermal material. On the 24th of December, commenced taking grafts from her left arm. The rapidity of the growth of her own grafts was simply marvelous. For example: One day thirty were placed in a double row, set three mm. apart. In thirty-six hours they put on a pink color. Three days later they were joined in one piece. The connecting parts were slightly depressed, and of course quite thin. By the eighteenth day a few hair like vessels were detected in the new tissue. Eight times after our first misfortune, numerous ulcers started at different places. At one time there were eighteen of them. They were evidently caused by the contraction of the subcutaneous tissue, whereby the fine vessels were broken.

Many of the ulcers healed of themselves, but the larger ones were assisted by autoplasmic grafts; 4,500 grafts were planted, of which the patient supplied 1,800.

May 1, 10th month, there remained uncovered four spots, each a trifle larger than a silver dollar. I expected to complete the scalp in four weeks, when suddenly her strength gave out. She took a severe cold, suppurative inflammation of the left middle ear followed, the mastoid process became very tender, and a dozen new ulcers appeared; two large ones, each about twelve cm. long and from one to three cm. wide. The combined area of all might have been represented by a circle sixteen cm. in diameter. The patient was almost disheartened; she slept only three hours of a night for three weeks; her head ached, sight grew dim, and I feared leptomeningitis was about to occur. Gave her opiates and

tonics. After three week's depression she reacted, when grafting was resumed. Occasional discouragements have been met with since, but now she feels good; says the head and eyes seem as strong as before the accident occurred. Her hearing has greatly improved also.

From the beginning, the entire wound was treated antiseptically with carbolic acid, iodoform, and corrosive sublimate. It was dressed with strips of carbolyzed gauze prepared with 2½ per cent. solution. The strips were 6 cm. wide, 30 to 35 cm. long. A layer of them, which came in direct contact with the granulations, was anointed with vaseline, so that it could be readily removed, then a series of strips was superimposed upon each other; next a piece of lint covered on one side with vaseline was placed over these and external to the lint a roller bandage made of cheese cloth was wrapped about the head. This formed convenient and serviceable dressing. The lint and strips were easily removed. The interstices of the gauze, by capillary attraction, drew the secretion from the wound's surface; the lint prevented too rapid evaporation whereby the dressings were kept from baking against the wound. The outer bandage retained the other parts in place. Thinking the lint heated the head, by way of experiment it was left off; but each time the dressings were stuck against the cranium, so that they were removed with difficulty, therefore continued using it.

At first the grafts were held in place by means of a piece of fish skin, which after thirty-six hours began to liquefy, and could be removed with a sponge. Repeated experiments caused me to discard this. The fish skin would break down, commingle with the secretion and help to macerate the grafts. Next tried McIntosh dressing. This heated the tissue too much, and the grafts were often found floating in the fluid beneath it. Having observed that the secretion always clung to the gauze dressing, that as a rule the thicker the layer of gauze used, the less discharge remained upon the wound's surface, I concluded to place a pad of oiled carbolyzed gauze upon the grafts instead of the rubber or fish skin. This worked remarkably well. Six or eight thicknesses of gauze, extending two or three cm. beyond the grafts in all directions, were applied. This had the desired effect of absorbing the secretion, reducing the heat to a minimum, while the grafts generally remained in place. The grafts usually assumed a pinkish appearance in forty-eight hours. Occasionally a series of the autoplasmic grafts when the patient felt exceptionally good, would be almost pink in twenty-four hours. Many of them, however, gave no evidence of being alive until after seventy-two hours. Several times grafts which hung to the dressings, covered with serum after forty-eight hours, being replaced, took hold and lived.

The progress of the growth of the new skin was similar to the description of M. Riverdin. A narrow, one mm. zone of hyaline tissue, somewhat depressed, surrounded the grafts in ninety-six hours; this grew with variable degrees of rapidity until the grafts were joined, then it grew thicker. After three weeks, sometimes a few days earlier or later, fine vessels were traceable in this new tissue; these increased in size until the skin was literally a network of vessels. As they continued developing it was noticed that the larger ones were directed upward. Later some of them assumed very nearly the position of the

normal arteries, like the temporal, occipital, etc. Now many of them have a bluish color, being evidently veins. The grafts were derived from her sister, three brothers, and herself, aged respectively seventeen, nineteen, twenty-four, twenty-seven and thirty years. They varied in size averaging about four by three mm., the central portion consisted of the entire *derma* with subcutaneous tissue, the marginal portion comprised epidermis only. Simply a fold of the skin was lifted with the sclerotica forceps and hastily snipped with iris scissors. When removed their under surfaces curved inward. They were then spread out and laid in direct contact with the granulations. The peculiar form of the grafts seemed to exert a favorable influence upon them. Their inward curvature caused them to cling closely to the granulations and an immediate suction action took place. The patient said she could feel them draw as soon as they rested upon the head. I have no doubt that the empty vessels in the grafts expedited this process.

Many times grafts six by seven cm. were tried without success. Tissue with hair was also applied; but the hair bulbs atrophied. Ten to one hundred grafts were planted at a sitting. Most of the time twenty to thirty only were used. Usually two days intervened between the graftings. Frequently, however, several weeks' rest had to be given the patient, on account of her weak condition.

All of the skin from the margin of the superior eyelids upward presents the characteristics of the normal skin. It is pliable, soft and movable. The lower half can be pinched up in folds. It imparts a velvety sensation to the fingers when touched. Furrows with pores opening into them are plainly visible with the hand glass. Follicles are abundantly present.

Her head sweats freely all over, and the new skin feels to the patient like the skin in other portions of her body. The eyelids are somewhat retracted, less, however, than they were four months ago. The younger portion of the tissue appears quite red, myriads of small vessels can be seen in it. As the skin grows older, the smaller vessels gradually disappear from view; others appear larger; the integument assumes a more normal appearance as to color and consistency. It also becomes more movable owing to the development of subcutaneous connective tissue. One cannot see where the new skin begins upon the eyelids. The lids close perfectly so that the lady sleeps with closed eyes.

Our works on surgery give no statistics, although at different times such accidents have occurred in the manufacturing States of our country. After having written many letters to physicians for information, concerning cases which came under their observation or treatment, I received responses from two only. I have imperfect reports to give of six cases:

Case 1.—A girl, nineteen years of age, lost her scalp fourteen years ago, in the Elgin Watch Factory, Elgin, Ill. Dr. E. Powell, of Chicago, informed me that the wound extended across the forehead above the eyebrows, and that her ears were not touched. Skin grafting was tried repeatedly in this case, with no results. Was told by a surgeon who saw her recently that one of her eyes was lost since by exposure occasioned by retraction of the eyelids.

Case 2.—Gross' "Surgery," last edition, makes mention of a young lady who lost her scalp in a factory in Philadelphia, 1869. He simply says the wound cicatrized over.

Case 3.—Reported in the "Surgical History of the War,"

Part I, p. 315. A conductor on the Union Pacific Railroad, was scalped by the Sioux Indians, near Cheyenne, in 1869. Wound as indicated by the illustration given, was elliptical, measured about 10x8 cm. A portion of the outer table of the cranium was removed. The wound healed over.

Case 4.—Reported by her employer, Mr. E. F. Smith. Miss Hattie Thomas, of Union City, Conn., March 29, 1871, in a button factory, placed her head under a swift running shaft, to pick up a button when her hair was caught by the shaft and her scalp taken off from the nape of her neck to her eyebrows. Her physician was Dr. S. C. Bartlett, now deceased. He cleaned the scalp and placed it upon the head about two hours after it was torn off, but it did not stay long. Then commenced skin planting. Had the head almost covered twice. Each time it was lost. A third effort proved successful in healing the wound, so that she was able to work after two years.

Case 5.—Miss Newman, Bristol, Conn., had her scalp torn off by her hair catching in the machinery in a mill where she was at work during the past winter. The entire scalp was gone, an eyelid and an ear slightly torn. She was treated by Drs. Wilson and W. E. Way, of Bristol. Result of treatment not given. "The Hartford Times" of last July, contained an advertisement for volunteers, to furnish skin for grafting.

Case 6.—Reported by Dr. W. W. Horton, of Unionville, Conn., June 12, 1885. Rosa Flynn had her entire scalp torn off by her hair catching in the shafting of a paper mill, Unionville, Conn. The scalp was torn from the back of the neck around, just above ear to the nose, taking the eyebrows. There was also flap torn down the right side of the face nearly to the mouth. I treated her for one month, then sent her to the Hartford Hospital, where skin grafting was tried. Both human and rabbit skin was used. It was a failure. Lids are retracted. The wound is still open. Hemorrhage occurs from the capillaries when her head goes to a lower level than her body. Expect the wound will cicatrize over in another year. It will be noticed that in none of these cases was the wound as extensive as in the one under consideration.

Deductions from experience in this case.

1. When grafts are set within five mm. of each other, a stronger skin is produced than when they are placed further apart.

2. A graft will grow to many times its dimensions, so that a circular graft three mm. in diameter will form a piece of skin eighteen mm. wide, but the skin will not be as strong or of as good a quality as when the grafts are closer together.

3. *Dermal* grafts with a few fibres of connective tissue form a smooth, soft, pliable, movable skin, presenting the characteristics of normal integument.

4. Epidermis placed on the cranium produces only a thin film, which ulcerates readily.

5. Auto-dermal grafts are more reliable than hetero-dermal grafts, providing the subject is healthy.

6. Grafts from a person near the patient's own age do better than from one much older or younger.

7. With *dermal* grafts the follicles are preserved, whereby the patient is made more comfortable, the skin produced is more natural, hence more durable.

8. The most reliable skin is formed by grafting about the circumference of the wound, say one or two rows, producing a narrow strip one cm. wide, and waiting until the blood vessels are well developed in it, then plant another series close to the last ones. By this plan, contraction in the marginal tissues takes place in advance of the subsequent graftings, diminishing the liability for the ulcerative process to occur by reason of such contraction.

9. The new skin is fed by the tissue upon which it rests, probably by osmosis, but depends for its permanent sustenance upon the vessels growing into it from the circumferential tissue. This fact is thoroughly established in my mind, as it was frequently demonstrated by the death of the patches of new skin removed from the marginal tissue. Time and ma-

terial can therefore be saved by constantly grafting from the circumference in large wounds.

10. All things considered, grafts will grow better in a new wound than in an old one; and the more recent the wound the more certain are they to thrive. In the present case, during the last three months, notwithstanding the improved physical condition of the patient, only twenty per cent. of the auto-dermal grafts took, while six months ago eighty-five per cent. could be depended upon.

582 Washington Boulevard.

DOMESTIC CORRESPONDENCE.

PHILADELPHIA LETTER.

The fortnight of Siberian winter which prevailed during the early part of January, has been succeeded by a genial thaw, which has filled the streets with indescribable compost, and the air with germs and moisture. Diphtheria and scarlatina are still with us; but are on the decline. No cases of small-pox or of typhus have appeared in this city for many months, but typhoid fever and phthisis claim entirely too much prominence in the weekly death returns. The City Board of Health is not as fully sustained by public sentiment as it should be, and in fact has been criticised for the continuance of diphtheria, of which there have been from ten to twenty cases daily. It probably should have done much more; but it has had difficulty in securing sufficient funds for properly carrying on its work. The method of dealing with contagious disease in the city is as follows:

All practicing physicians are required to report every case of contagious disease (not including consumption) within twenty-four hours, under a penalty of \$50 for neglect.

So soon as a case of diphtheria or other contagious disease is reported, no matter what the circumstances of the family may be, a medical inspector is promptly sent to the place to make an inspection. If he finds that it is a case which cannot be properly cared for at home, he sends it to the Municipal Hospital, and institutes proper measures for disinfection, and holds the place under observation until the period of incubation of the disease has passed. If the disease is one which does not require such rigid measures, he holds the case under supervision during its entire continuance. It is his business to supply families with circulars, giving plain instructions as to the management of the disease from beginning to end, and after convalescence; or after death, should it occur.

A medical inspector is sent to coöperate with the medical attendant in all necessary precautionary measures, and, if disinfection cannot be properly carried out by the family, a disinfectant is placed in charge of the work, the disinfectants, chloride of lime, liquid carbolic acid and sulphur, being furnished gratuitously. If there are children in a family where contagious disease exists, it is the practice to notify the principal of the school where the children attend, to that effect. A patient is always isolated and the family cautioned to prevent intercourse with the house. When this advice is not taken, and where there is a lack of confidence of the ability of a family to carry out sanitary measures, it is the practice to place watchmen before the premises to see that the instructions of the Board are complied with.

The Board, also offers freely to the poor the use of disinfecting chambers and cremation furnaces at the grounds of the Municipal Hospital, for disinfecting such articles as cannot be done at home. The rules governing the burial of bodies are strictly enforced. The medical inspectors, of

whom there are 15, are giving the greater part of their time to the supervision of diphtheria cases, while five disinfectors are employed to purify houses where diphtheria is prevalent.

The Health Officer, Major Veale, has to contend with doctors who are loth to report contagious diseases until the cases terminate fatally, and with others who hold to the theory that diphtheria and scarlet fever are not contagious. He thinks that there ought to be a building, or series of buildings, where persons who are liable to contagion could be taken and kept under observation, their homes in the meanwhile being thoroughly disinfected. The city at present, generally speaking, is in a fair sanitary condition.

There is a movement now on foot to establish a special hospital for the treatment of diphtheria, or of infectious diseases other than small-pox, for which a Municipal Hospital already exists. Provost Pepper of the University, has come out with a strong appeal in the *Public Ledger* from which we take the following paragraphs:

"When will the people come to be willing to pay for the inestimable blessings of preventive medicine? Do they not know that modern methods of isolation and disinfection enable us to prevent the spread of infectious diseases, if the patients are placed under proper conditions? The germs of these infectious diseases retain their vitality and their virulence for a long time. They attach themselves to furniture and clothing. They are carried by physicians, by nurses, by relatives. Infected linen is sent to the laundries, and a fresh centre for the spread of the disease is created. Cases in families, after arriving at a hotel, develop. There is no suitable place to send the patient. The occurrence is concealed as far as possible. Isolation is impossible. Disinfection is imperfectly carried out. The germs of the disease remain in the apartment and may also be diffused widely. So long as there is no attractive, properly constructed and well administered special hospital for contagious diseases, with isolation wards for each of the chief affections, it is impossible for the community to escape the grave consequences of their neglect.

"The Municipal Hospital answers an excellent purpose, but there is needed in addition a special hospital, more centrally situated, with accommodations adapted to the various classes of patients, and with a staff of attending physicians of large experience in the treatment of these diseases. There need be no objection to the location of such a hospital near the centre of large cities. A sufficient space should be secured, but it would not require an extensive tract. Proper construction and careful administration would remove any chance of the disease spreading in the neighborhood.

"The prompt removal of each case of contagious disease from hotels, public institutions and as far as possible from private residences to such a hospital would effect a vast saving in life and in the risk of the disease spreading. As a matter of fact, the well-to-do classes will not allow one of their family to go to the Municipal Hospital save under the constraint of law, in the case of small-pox. As a further matter of fact, it would be essential that all respectable physicians in the community must have the privilege of attending patients whom they may send to such a special hospital as is here urged. Otherwise it would be idle to hope for the coöperation of the profession or the support of the community."

This proposed hospital has met with general favor among physicians and Dr. Robert B. Cruice has made the suggestion that, as the Government has no further use for Fort Mifflin, this island in the Delaware would provide an admirable location for a hospital for infectious diseases. Thus far, however, no steps have been taken towards carrying out these important suggestions.

While discussing hospitals, it should be noticed that the friends of the late Prof. D. Hayes Agnew have concluded, very opportunely, to erect a memorial ward at the hospital of the University of Pennsylvania, devoted particularly to the treatment of surgical diseases and injuries of children. This memorial seems especially appropriate, since Dr. Agnew was known to be opposed to the post-mortem