diet the urea continued to increase throughout the day.

During the experiment the usual bodily and mental work was performed. He noticed, however, during the restricted diet that he experienced a feeling of faintness and some increase in his appetite. Before the experiment was commenced he noticed that his urine was loaded with alkalies. During the restricted diet his urine was clear and the quantity passed was larger than when animal food was being consumed. This shows that when an increased amount of urea was excreted a less amount of urine was passed. Had the urine been increased with the increased amount of urea excreted, it might be argued that the increased amount of urea was due to the increased amount of urine passed.

In the second experiment of two days, on a very restricted diet, the urea fell from a range of 21 to 25 grains per hour to 8.87 grains per hour. At the end of two days, on the very restricted diet, nitrogenous food was taken in the form of egg and milk beaten together so absorption would take place more easily and quickly. Half an hour later an ordinary meal of cold meat was taken. During the three hours succeeding the first ingestion of nitrogenous matter, the urea excreted stood 12.43 grains per hour against 8.87 grains per hour, the mean amount for the eight hours previously.

Now in Bright's disease of the kidneys we have certain conditions to be met and ameliorated in order to benefit the patient so afflicted. It must be admitted that when the kidneys are diseased they are embarrassed by reason of having too much labor to perform and owing to their enfeebled condition, being unable to perform the necessary functions which is shown usually by a dropsical condition of the system. Reason would then dictate that the kidneys should be relieved of a part of their duties. Now from the foregoing observations we have learned that in order to diminish their labor it is necessary to prescribe for our patient so afflicted, such food as will produce the least amount of or excrete the least amount of urea. It has been shown that urea is most excreted when nitrogenous diet is ingested, and least excreted when non-nitrogenous food is ingested. It has also been shown that the bulk of the nitrogen in nitrogenous food is excreted by the kidneys in the shape of urea. The food necessary to be ingested then by a person suffering with Bright's disease of the kidneys should be principally non-nitrogenous or approximating it.

It has been shown, also, by the above observations that when non-nitrogenous diet is ingested, urea is very much decreased in quantity, and that the urine passed is clear and increased in quantity. Then it will be seen that non-nitrogenous food not only diminishes the urea excreted in the urine, but leads to an increased amount of that fluid. This being true, it must be admitted that the patient suffering from Bright's disease is doubly benefited by a non-nitrogenous diet ingested. Namely, rest to the kidneys and an increased flow of urine.

The patient with Bright's disease is also benefited by the carbo-hydrates for the reason that the cast off material from the ingestion of them is excreted by the bowels and skin and otherwise. This would also add to our armamentarium in the treatment of a case of Bright's disease. The bowels are kept more soluble, throwing off an amount of liquid that might

not otherwise be excreted and they also act as a diaphoretic, passing through the skin an additional amount of fluid.

DIPHTHERIA; OUR MODERN VIEWS—ITS ETIOLOGY AND PATHOLOGY, WITH REMARKS ON THE EARLY HISTORY OF THIS MALADY.

Read in the Section on Laryngology and Otology, at the Forty-fifth Annual Meeting of the American Medical Association, held in San Francisco, June 5-8, 1894.

BY J. MOUNT-BLEYER, M.D.

FELLOW OF THE ROYAL ACADEMY OF MEDICINE, NAPLES, ITALY; SURGEON TO THE NEW YORK THROAT AND NOSE HOSPITAL; LARYNGOLOGIST TO GERMAN WEST SIDE CLINIC, ETC., NEW YORK.

In the work of Hippocrates on dentition we find the following: "*Quibus cito in tonsillis ulcera serpentina considunt, febrilis ac tussi per manebitibus, periculum est rursus esse generanda ulcera.*"

This is the first record of testimony regarding the recognition of the disease we now classify as diphtheria. Whether Hippocrates had in mind the simple aphthous inflammation of the mouth or the well-defined disease itself is an open question. Suffice it to say, the above quotation is the first literature we possess upon ulcerative and exudative disease of the pharynx, tonsil, uvula, larynx, etc.

Aretæus gives us a more satisfactory clinical picture and intimates that Egyptians were quite familiar with diphtheria. Ætius, who lived some two centuries later, left a manuscript giving clear evidence that he understood diphtheria. And so we pass on rapidly to the end of the sixteenth century. It was the distinguished French physician, Baillou, who it may be truly said gives us the first accurate description of diphtheria. He speaks of a false membrane which we can scarcely mistake for anything save fibrinous exudate of diphtheria.

Ever since the day of Baillou, diphtheria has prevailed in the various quarters of the world.¹ We have the history of an epidemic of malignant angina in Paris toward the close of the sixteenth century; again we find it in Spain—as we learn from Villa Real, who studied an epidemic which raged in Spain in the early years of the seventeenth century, which he portrayed as *garotello* or more specifically *morbi suffocantis*, and which possesses without drawing the broad line of differentiation between diphtheria and the scarlatinal exudates.

Coming down to more recent times, we find such eminent observers as Dr. Patrick Blair, Schobinger, Ghisi, John Star, Huxham, Francis House, Trouseau, Samuel Bard and others engaged in the study of diphtheria. The first exact record comes to us in the form of two treatises on angina diphtheritis which were submitted to the French Academy of Medicine by Bretonneau in 1821. In these brochures he made the broad claim that an inflammation without an exudation is never diphtheria, while no inflammation with an exudation is diphtheritis, when it does not spread by contagion.

He believed that inoculation was the only means of conveying the disease from one to another, and held firmly to the opinion that the atmosphere did not act as a conveyor of the contagion. We have since modified our views regarding the spread of the contagion of diphtheria, and unfortunately the violent epidemics in France, Holland, Germany and our own country, where it raged from time to time, have

given us ample opportunity and material for the study of this feature of the disease.

It was Virchow who established the pathologico-anatomic basis upon which we have since based our investigation of the diphtheritic process. He taught us to direct our attention to the exudation in the substance of the mucous membrane, and the death of this membrane in consequence of the interference with its nutritive supply, in contra-distinction to croupous inflammation where the exudate does not penetrate the structure of the membrane but simply lies upon its surface.

There are still many who hold that diphtheria and croup are essentially similar conditions pathologically considered. Their numbers, however, are rapidly diminishing, and there seems to be little question that within the next decade we will all agree that the views of Wagner will have to give way to the more recent bacteriologic and pathologic discoveries.

Bahl considered diphtheria as a general infectious disease, entirely independent of any previously existing local disease, whose distinctive characteristic was a nuclear or cystoid growth of the submucous connective tissues which caused the death of the membrane by compressing the vessels of its vascular supply. To him the manifestation in the throat and tonsils was simply an indication, or symptom as it were, of the general infection just as the eruptive basis chose the skin as the most favorable seat of characteristic indication.

These are the views about which the profession at large are still at variance. The older practitioner, whose good judgment is based upon a large experience, is not yet ready to resign the theories in vogue during the longer period of his professional career, but there can be but one correct solution of the problem concerning the origin and character of a disease which even to-day is an almost unconquerable foe, and the weight of research prompts to the fact that we must accept and carry on the work laid out by Oertel, Huebner and Welsh if we hope to be able to cope intelligently with diphtheria. The bacillus, described by Löffler and Klebs, coupled with the study of the toxic ferment it ushers into existence, and which finds its way into the circulation, together with the pathology as Oertel has taught it to us, it seems accounts in a satisfactory way for the clinical manifestation of the disease. In other words, if I may be permitted to say it, we have only now begun to know something about the nature of diphtheria.

The etiology of diphtheria, in a great measure, is the practical lesson we have learned from the researches of Klebs, Löffler and their followers. Before the time of these observers we were apt to confound the exudative diseases of the pharynx, tonsils and upper respiratory tract. To day we know, with almost absolute certainty, that there are two forms of this class of disease, which to the eye of the clinical observer present similar changes in the mucous membrane, inasmuch as both are characterized by the appearance of a pseudo-membrane, at times dirty white, often tinted greenish or yellow white in color. Viewed at the bedside alone, in both you will observe alike, prostration, a febrile condition, swell-

ing of the submaxillary glands, disturbance of respiration, often slight it is true, so that a differential diagnosis, clinically, is a qualification to which few of us can lay claim. Notwithstanding this, the two are totally dissimilar. Examined bacteriologically, the exudate of one will reveal nests of bacilli which are found only in the exudate of true diphtheria, while in the other there are no bacilli, but instead the so-called staphylo- and streptococci.

The first variety is highly fatal, statistics showing a mortality of almost 50 per cent., while the second, so far as danger to life is concerned, is comparatively harmless. Thus it becomes of great importance, whenever we have a case of exudative disease of these parts to deal with, to make, if possible, a positive diagnosis of its true character. The New York Board of Health has instituted a most practical method by which every practitioner can ascertain beyond a doubt the disease under his care, and thus avoid mistakes. Where this mode, as aforesaid, can not be made use of, I will explain, when discussing the subject of bacteriology, a very simple manner in a comparatively short time, so that the clinical observer can avail himself of this means of confirming his diagnosis, thus saving much time and trouble.

The causation of diphtheria leaves us much in doubt. We know that it prevails at all seasons of the year, most extensively in the damp chilly weather of the late fall and early winter. We find it in all latitudes and longitudes, although quite recently I was assured by Dr. Angel Garvino, Professor of Bacteriology at the University of Mexico, who visited New York, that the City of Mexico, which lies in a very high altitude, some 1,500 meters above the level of the sea, was comparatively free from diphtheria; in fact so much so that the experience of a case of true diphtheria is regarded with rare interest by the medical men of that city. The disease, too, is not limited to classes. We find it among the rich and poor alike; in well and badly ventilated and drained houses; in the thickly and sparsely populated districts.

Essentially, diphtheria is a disease of early childhood, although no age is exempt from it. Sometimes we find it raging in epidemics, again it is met with in isolated cases in which the question of origin by contagion can have but little weight. In short, these are the conditions under which we find diphtheria.

Many valuable statistics have been collected on the subject, but the limits of a chapter like this do not permit of such detail. Then, again, profitable as statistics are to the specialist and the doctor at large, they make the driest of dry reading. For such as desire more in this regard, the late brochures of Huebner, Baginsky, Oertel, von Brieger, Sidney Martin, William Welsh and others may be consulted.

Diphtheria presents itself under two series of symptoms—the local, and constitutional manifestations, to which may be added a third form best considered, however, as the sequelæ of the disease. Thus in speaking of diphtheria we should bear in mind these conditions, although they do not always manifest themselves together in each individual case. The local disease makes its appearance in an inflammatory process upon certain mucous membranes and denuded parts of the skin, the most common seat of affection being the membranes of the throat and fauces, those most directly in contact with the air, and the inflammation leads to formation of a membranous exudate which is usually the first indication that causes us

¹ I have purposely avoided more extensive reference to the bibliography of the history of diphtheria which may be found in nearly all the works on diphtheria, particularly that of Oertel, Mackenzie and others, in order to devote my limited space to the consideration of the disease in its more modern aspect.

to suspect the existence of diphtheria. Upon superficial examination the clinical observer meets with grayish or dirty white false membrane deposits upon the mucous membrane of the mouth, the pharynx, tonsil, nose; in fact, anywhere along the upper air passages. It is under the same conditions found upon the conjunctiva, the rectum or vagina, or upon any of the mucous membranes of the body that are directly exposed to the air.

When this false membrane invades the deeper mucous membranes, such as that of the intestinal canal, etc., we are inclined to regard it as secondary infection, notwithstanding that we lean to the opinion that the diphtheritic process is usually circumscribed locally, and spreading by way of the general circulation all through the body. Thus we account for the evidence of diphtheria as we find it in the various organs as a primary condition. This, by the way, is the most modern view taken of the disease and the one which to us, after careful consideration, is the theory most in accord with clinical manifestations.

Diphtheria is an infectious disease, and may be classified as somewhere between simple excitement of the circulatory system, and the most aggravated form of typhoid fever and pyemic poisoning. The sequelæ, in my opinion, will depend more or less on the extent of the general invasion from the original local disease, and largely upon the quantity and nature of the septic ferments that find their way into the circulation, involving those organs that offer the least resistance. The latest investigations point to the necessity of turning our attention to the form of true diphtheritis as much as has been hitherto done to the local manifestations of the disease.

In the study of the etiology of diphtheria the first consideration is to explain, if we can, the process of the disease itself. Roux and Yersin, in substantiation of their theory that the systemic effect of diphtheria is not due to the bacillus described by Löffler and Klebs, but rather is caused by the ptomaines or poisonous substances produced by it, proved it by separating the bacillus from the ptomaines and subjected it to chemic analysis and physiologic test. They described the disease as a toxemia, differing from others in the respect that the toxic acts as a ferment which when injected into living tissue, even in the most minute quantities, ushers into existence certain well defined albuminous substances or bodies.

Sidney Martin succeeded in isolating these poisonous albuminoids from the organs of persons who died from the disease. This substance was secured from the spleen and blood of children, was subjected to repeated alcohol falling and resolved itself into a proto- and deatro-albumose which he injected into the skin of rabbits.

In doses of 0.117 per kilo it produced a febrile condition and edema in the animal; slightly increasing the amount to 0.112 per kilo, rapid death of the animal followed. Intravenous introduction of this albumose in the same animals produced fever, progressive paralysis and degeneration of the heart, nerves and muscles, precisely analogous to those conditions we find resulting from diphtheria.

Löffler's theory of the local origin of diphtheria, when contrasted with the clinical manifestations of the disease, does not seem as plausible as when viewed from its bacteriologic and pathologic standpoint alone. True, he has given us a most careful

and competent description of the bacillus, since honored with his name, and the majority of clinicians agree that upon the presence of this bacillus alone can the diagnosis of true diphtheritis be made beyond peradventure. We are not all bacteriologists, and we propose to lay the greatest stress on the clinical view of diphtheria. I shall refer to its bacteriologic significance only so far as it can be practically made use of at the bedside and aid us in diagnosis, treatment and prophylaxis.

Ab intra diphtheria appears to be a disease which draws into sympathy certain organs, preferably the heart, muscles, kidneys and nerves, and thus it becomes at first glance difficult for us to understand how it is possible that such conditions can be brought into play by a microorganism that has its habitat near the surfaces, and it is even more difficult to harmonize with the insidious progress of the disease and the toxemic symptoms we observe. Once, however, we establish the presence and origin of the diffusible ferments, all these characteristic manifestations become apparent. Bear in mind that the bacteriologic investigations of recent years do not stand alone; we can carry them one by one to the bedside and study them side by side, and reason logically from cause to effect.

One thing judged by the weight of careful research appears true, that it is essentially certain we can no longer hazard a positive opinion upon a case of diphtheria without having ascertained the presence or absence of the bacillus. Many of us appreciate the force of this broad statement from practical lessons learned at the bedside where without any indication that would direct our attention to it, we have seen certain cases of diphtheria resolve themselves in from one to three days into simple exudative tonsillitis or angina lacunaris.

Skigo reports several such cases which he met with during an epidemic of diphtheria, where the tonsils were covered with disseminated spots of exudate the size of a pea; and he tells us that there was absolutely no clinical indication by which he could differentiate the cases which ran the violent course of true diphtheria, and those which took the milder course.

The value of the bacteriologic character of the exudate in this connection received much attention from Morton, who examined 200 cases of diphtheria that had been diagnosed by that eminent clinician, Dr. Simon, of Paris. In only 137 of these cases did he find the bacillus; 43 of the remaining 63 cases proved to be tonsillitis; 13 proved to be croup, of which 12 got well without operation; 7 cases proved to be diphtheria with croup, 6 of which developed bacilli after the operation of tracheotomy had been done, and which Morton says was clearly due to hospital infection.

It is necessary in the very first instance to differentiate between exudative tonsillitis and true diphtheria. This, if we are obliged to follow the procedure of Koch, which requires a bacteriologic experience which the average clinician does not possess. The cultivation of the bacillus in glycerin agar requires time and patience, to say nothing of careful work. The bacillus must be isolated from the membranous exudate. The cultivation of the bacillus in blood serum requires at least from twelve to sixteen hours, and a day or a day and a half, at least, in making the diagnosis. With a case of true diphthe-

ria to deal with, so much time can not be sacrificed, and hence we must look to other more rapid means of gaining the object of which we are in search. In hospitals, with all the facilities at hand, we can allow ourselves privileges that we can not think of in private practice.

But we have a simple means at hand of making a hurried and certain examination of the exudate, where one can not have the opportunities of a city board of health. The method is the one devised by Roux and Yersin, modified by Huebner. A bit of the yellow covering of the membrane from the tonsil is removed with a sterilized spatula, and rubbed over the surface of a cover-glass until the film is very thin and almost dry. The glass is then passed through a flame and stained with methyl blue. The specimen, it will be found, readily takes the stain, and inside of a very short time it is ready for the field of the microscope. An ordinary objective magnifying glass of 400 or 500 diameters answers the purpose, and with a little practice, between the detritus, mucus, pus cells and the various bacteria heaps, you will observe nests or pockets of mixed up bacilli about the size of the tubercle bacillus, most of them slightly bent, some club shaped, often irregularly seamed.

With the presence of these nests established you may be reasonably certain, in fact we almost feel like saying absolutely certain, that the case is one of true diphtheritis.

On the other hand, if you find only the cocci—the strepto- and diplococci, with here and there, short, thick, weakly stained bacilli not arranged in nests, you will, in all probability, in fact you may be certain the case you have to deal with is not a true diphtheria. The statistics of Huebner covering 113 cases examined by him during a period of six years after Löffler described the specific bacillus, go a great way toward supporting the above statement.

In thirty-six cases, Huebner found cocci, many solitary ones, with some side by side, but no nests of bacilli with stained ends. Ten of these cases were scarlet fever, and thirteen presented the clinical picture of follicular tonsillitis or croup. The exudate of the other seventy-seven cases revealed bacilli arranged in nests; of these the nests of bacilli alone presented, while in fifty-three there were nests besides cocci. Of these seventy-seven cases, fifty-three died with all the manifestations of true diphtheria. The mortality of the cases where the bacilli predominated in very large quantities was 66 per cent., and where the nests and cocci both existed 59 per cent. died. Of the thirty-two recovered cases, seventeen presented all the clinical appearances and were of the most violent type, four were tracheotomized and fifteen were mild cases without any of the marked characteristics of true diphtheria.

From this, alone, the high value of a bacteriologic examination becomes evident, and to say the least, it is beyond question the most reasonable diagnostic factor we have had in years and of undeniable practical value, for the reason that in each case the cover-glass method before described alone was sufficient to make a satisfactory examination.

"It is possible," says Huebner, "that a recovered case may show bacilli some days after the disappearance of the membrane." I can add my testimony to this, for in four cases similar, I also discovered sparsely filled nests, two days after all visible signs of the membrane had disappeared.

It must not be forgotten that you will find cocci in angina lacunaris to which Fraenkel called our attention in 1886. And lately the claim has been made that a small coccus is always found in the pseudo-membranous disease, which is pathognomonic of the disease. This still remains to be proved, although it appears to us to be a fact ready for acceptance, yet it will require more careful investigation before we can put ourselves on record with an unmistakable picture of this microorganism. Martin, of Paris, first described the organism which he pronounced to be pathognomonic of the pseudo-membrane, and I am inclined to agree with him. The significance and value of these recent researches will bear much upon our treatment of this malady.

With Laycock in 1858, we have the first intimation of the possible parasitic origin of diphtheria, the parasite being the *oidium albicans*. Later, Letzerich ascribed it as due to the *zygodemus*, and Tuscos and Joffe to the *leptothrix buccalis*.

In 1868, Oertel carefully studied the habits of micrococcus which he isolated from the exudate, membranous exudate. It occurred as a minute, dark, point-like microbe; some round, others oval, occurring singly or in masses—*zöoglea*; it was always accompanied by the rod bacterium. In 1883, Klebs examined the exudates obtained during an epidemic in hemorrhagic diphtheria in Prague, and claimed to have discovered the specific germ of true diphtheria—the *microsporon diphtheriticum*, a small micrococcus, arranged in round balls covered by a thin layer of gelatinous matter. These subsequently developed into motive bacilli and finally into tufts of mycilium.

In another epidemic occurring in Zurich, and characterized by a rapid extent of the membrane into the larynx, followed by interstitial inflammatory process in the internal organs, Klebs came across a microorganism that differed totally from the one he first observed. It was lobular and essentially belonging to the bacillus group. He naturally concluded that there were two forms of diphtheria, *i. e.*, diphtheria *microsporon* and diphtheria *bacillaris*. Klebs undoubtedly was on the right track, but he lost the scent, which also escaped Emmerich-Solomen, until Dr. Frederick Löffler took it up and accurately pictured to us the bacillus which we now accept as characteristic of true diphtheria. It is essentially the one observed long before by Klebs whose weak point seems to have been insufficient accuracy and evenness in conducting his investigations.

To Löffler we owe the streptococcus and the bacillus; the bacillus of which I desire to speak at length here. It is non-mobile, its form is difficult to describe in words that will give you an adequate idea of its appearance to the eye, for it is neither straight nor bent; in short, it is about the size of a tubercle bacillus, somewhat thicker. It is not found in the intestinal organs, and micrococci morphologically identical with those found by Löffler are observed also in various diseases accompanied by inflammation of the mucous membrane, like variola, typhoid and puerperal fever, but their presence there is regarded as purely accidental.

Löffler found his micrococci not only in the membranes, but also in the lymphatics, from which point they penetrated every part of the body, causing a tissue necrosis.

The micrococci are not always present. Inocula-

tion by injection of streptococci do not produce true diphtheria. This is the great evidence that they are not specific of the disease. Not so with the bacillus.

The bacteria of diphtheria is aerobic; that is, it requires oxygen for its life and growth.

The pathology of diphtheria.—As in the preceding pages on the etiology and bacteriology, I will not be able to go into details, but rather confine myself to the pathology, as it is most nearly associated with changes that are manifested by ordinary clinical observation. The most complete studies on the pathology of the disease have been made by Oertel, both as to the structure of the membrane and the changes in the tissues of the body, as we find them affected by diphtheria.

Let us take up the membrane first. In color, it is grayish or white, varying in thickness from one to three millimeters. It is tenacious, thick and closely adherent to the parts, and for this reason, on removing any of it for examination it is necessary to scratch with some vigor and not scrape lightly over the surface with a spatula. In the main it consists of degenerated fibrinous epithelium and faser stoff exudate. This with the coagulation of the fibrin produces a hyaline substance made up largely of degenerated cells and tissue elements. This occurs not only on the surface of the membrane, but deep down in the sub-epithelial tissue. The inflammation and fibrinous exudate is a secondary process.

The classification the modern observers are inclined to make, summed up is this: Diphtheritic pseudo-membrane is true diphtheria. Fibrinous pseudo-membrane is not true diphtheria. The two, as they appear clinically, are absolutely identical. There is absolutely no way of differentiating when viewed in gross. Practically speaking they are alike, and it is in this connection that the bacteriologic examination alone can determine the diagnosis. Both are a coagulation of fibrinous exudate.

The degenerative process in the spleen, in Peyer's patches, in the lymphatics, the solution and degeneration of the cells into coagulable hyaline substance are precisely similar to that which produces the pseudo-membrane in the larynx, trachea, uvula and pharynx.

The histologic structure of the membrane varies. The epithelium, according to von Wagner, is a fibrinous metamorphosis. Between the epithelium there are leucocytes which (*necrotistisch zu grunde gegangen sind*) have become resolved into hyaline substance, the masses of (*fibrinogener flüssigkeit*), which transudated assumes in the epithelial structure the form of a network or fibrillated structure.

Masses of larger round cells are also intermixed, some broken down, others still retaining their apparent integrity, with disintegrated similar masses. The first change occurs in the nuclei of the cells.

The major part of the membranous masses of the pseudo-membrane is of secondary origin. It originates from the mucous tissue and the elements which form it have collected and originated in it. They were simply evolved from the depths of the membrane to the surfaces.

There is a fibrinous infiltration of the connective tissue observable in the lowest strata of the mucous membrane, also hyaline degeneration of the membrane; larger and similar round cells with the detritus of the breaking up of these cells, around which in the deeper layers we find healthy cells sickened, or

so affected that they have gathered in heaps ready to undergo further change.

These, in brief, are the pathologic products and changes we find in the structure of the membrane and its continuance indicates that the disease is advancing.

It is claimed that pathologic changes, such as have been described, may be produced by injections, in the membranes and the tissues directly underlying, of powerful irritants, and thus the possibility of producing diphtheria by mechanical means is claimed to be possible.

460 Lexington Avenue, New York City.

DUTY—LOYALTY—SUCCESS.

Read before the Joint Meeting of the Golden Belt and Northwestern Kansas Medical Societies, held at Ellsworth, Kan., Oct. 4, 1894.

BY WILLIAM B. DEWEES, A.M., M.D.

SALINA, KAN.

Fain would I call the attention of not only those within the sound of my voice, but of every member of the two Societies here represented, as well as of every reputable practitioner in America, to the too much neglected duty of contributing to and studying the proper current medical literature. By the right endeavor in this direction the great distinctive service of American practitioners of medicine may justly become effectually manifested by its counter-acting influence in removing the pedantry shown in some of our medical colleges and periodicals, and encourage the expression of original thought and observation in young physicians. The distinction of young physicians is made because it is from this rank that the future advance in medicine will be made, and also from the cognizance that most of the older members of our guild have never learned to write nor to observe or study, while as they become burdened with the cares of a large practice or the habit of indifference, it becomes impossible for them to acquire the art of writing. Thus, whatever knowledge they may possess of positive value to the profession, dies with them.

Advance in medical education follows the earnest endeavor of the thoughtful and observing doctor to learn something of profit and interest, and then place this before the profession in the most attractive manner. When a young practitioner has begun to realize that he can talk to the medical profession throughout the State, the nation, and even the world, life and study take on an entirely new aspect. To the day of small things will now be added the day of an enlarged and enlarging physicianhood. He who fully realizes this truth will never write a poor article for publication. The poor articles come from quite a different sort of doctors. To every young practitioner who would make the most of his powers I would say: Think and observe for yourself, and write for the best periodicals constantly. Though it be only one or two articles a year and much time be required to prepare the same, be sure to acquire and keep up the habit of correct thinking and observing, constant study and perfect writing.

With regard to studying the proper current medical literature, it is but necessary to call attention to the fact that no practitioner can be considered as progressive who has not provided himself, first of all, with the true representative periodicals of the profession of his State and nation. Therefore every