deals with such questions as other evidences of unconscious activities, other instances of the mechanisms of identification, condensation, displacement of affect, abreaction, and the process of overdetermination, other indications of the significance of morbid anxiety and compulsion (Zwang), and so on. He illustrates all these matters with a rich series of examples taken from daily life as well as from various writers, and expounds them with remarkable lucidity. More important still, perhaps, is his penetrating discussion of the fundamental psychological principles involved in Freud's work. Unfortunately, this part of the volume does not lend itself to summarizing, and in any case it certainly should be read in the original.

In his concluding resumé Bleuler states that in his opinion "the essential part of the Freudian theory rests in a logical manner on assured facts, so that one has to accept it as correct. . . . The other part of the Freudian psychology is not nonsense, but disputable hypothesis, which may prove to be very fruitful. That in the minor work of the whole school many details are problematical, too hastily generalized, or directly untrue, should not appear strange. It would be curious if in this freshly explored field and in the endless complication of the mind false conclusions were not reached as well as in every other sphere."

From a man of Professor Bleuler's authoritative standing in psychiatry, and from a man who has spent as many years as he has in the study of this particular branch of the subject, this criticism must carry especial weight. Additional value, and one might say charm, is obtained by the equanimity and moderation of its style, which might with advantage be taken as a model by many overheated writers on both sides of the controversy. To those who are desirous of attaining an impartial decision on an involved subject, and of removing various misconceptions that are bound to arise when one's acquaintance with it is imperfect, to those, in short, who prefer the investigation of truth to merely proving themselves right or others wrong, this volume can be warmly recommended.

ERNEST JONES.

THE FUNCTION OF SLEEP (*La Fonction du Sommeil*). By
Albert Salmon. Paris, 1910. Pp. 235.

In this monograph, Salmon returns to his theory of the relation between sleep and the function of the pituitary body, which he had originally promulgated in 1905. He starts with an account of the mechanism of plants by which they close their leaves

and petals in darkness, and open them either in natural or in artificial light. This he considers sleep, a view which appears to us to be erroneous, since plants do not possess a nervous system, and therefore literally cannot be said to sleep. This opening and closing of leaves and petals in plants is probably a mere chemical mechanism, allied to a tropism and not sleep at all. Our daily sleep is a biochemical phenomenon, whose function is the nutrition of the central nerve elements, and consequently those tissues which depend upon the activity of the nervous system rest likewise during sleep. Sleep is therefore really a biological reaction of defence against starvation. In the loss of coenesthetic sensations which precede sleep there is a diminution of muscular tonus, and in sleep itself we give to the body its least muscular effort. In the semi-drowsy states there is also some loss of muscular tonus, but here it is not complete. In this semi-drowsy state, also, hallucinations very frequently arise, due, undoubtedly, to the activity of the peripheral sensory neurones. He then discusses the conditions which resemble sleep, such as stupor, hypnosis, electrical sleep, conditions of muscular inhibition in animals and narcosis.

The primary psychic condition for sleep he believes with Claparede to be disinterest and inattention. On the contrary, some conditions, such as hypnosis, may outwardly resemble sleep yet are not identical with it. For instance, ordinary stimuli if sufficiently strong will awaken a sleeper, but not a hypnotized subject. Hypnosis, as a rule, is rapidly induced and does not assume the characteristic course of sleep. In sleep there is a certain degree of muscular tonus. The inhibition of muscular movements in animals, when they are suddenly placed in an abnormal position, is not due to fright, because there is no loss of sphincteric control and no symptoms of fear except a slight tremor of the rigid limbs. Sleep is a negative state, a cessation of brain activity, an instinct. This instinct consists of a periodic relaxation of muscular tonus, which becomes automatic by habit. Lack of interest and diminished attention cause sleep, because these are unaccompanied by muscular tonus. When we sleep we take the position in which there is the least muscular tension.

All secretions are regulated by psychic stimuli, particularly the salivary, gastric, and pancreatic juice as shown by Pawlow. The desire to sleep is brought on by a gland or secretory apparatus, which has a periodic activity. The principal factor in sleep as in secretions is the periodicity. Both depend on habit and psychic stimuli. The various points of analogy between sleep and the function of secretion is one of the strongest arguments for the secretory

theory of sleep. The prime motive in sleep as in secretion is psychic,—the desire to sleep. The sensation or the desire for sleep is called the appetite for sleep, analogous to normal appetite and hunger. Both the appetite for sleep and for food are rhythmic mechanisms. Through habit we can change the hour for hunger as well as of sleep, and therefore both are due to psychic stimuli. The need of sleeping and of eating are organic needs, and the curve of the gastric secretion bears a striking analogy to the curve of the depth of sleep. Both are more intense during the first or second hour, and then gradually descend and both are increased or diminished by the same circumstances, such as sea air, excessive fatigue, cold, etc. Drugs which diminish gastric secretion, such as atropine, cause insomnia, those which increase it, such as alcohol, produce drowsiness.

Disorders of sleep are very frequent in those diseases which interfere with the internal secretion. For instance, the drowsiness of myxedema, which is due to a diminished secretion of the thyroid gland, diminishes under thyroid treatment, while in goitre, in which there is an increased secretion of the gland, insomnia is frequent. Somnolence also occurs in adiposity, which condition, according to modern researches, may be due to an alteration in the secretion of hypophysis. Somnolence also occurs in diabetes, a disorder of glycogenic secretion, and it is interesting to note, that in hypophyseal tumors, with or without acromegaly, there is frequently associated a glycosuria.

The very marked relation pointed out between disorders of sleep and hypophyseal tumors, forms one of the most interesting and important parts of the book. Somnolence or even prolonged sleep or stupor are very frequent in hypophyseal tumors. In most pituitary tumors of any lobe, there is either a prolonged state resembling sleep or frequent attacks of drowsiness if there is a hypersecretion, or insomnia, if a diminished secretion. He also points out that excluding pressure symptoms, somnolence is very rare in brain tumors unless they invade the hypophysis or press upon it, and thus interfere with its secretion. In the histology of the gland there are not only secretory cells, but also bipolar nerve cells. Somnolence has also been reported in hypophyseal tumors, where the tumor was too small to exert pressure or interfere with cerebral circulation, and therefore the sleepy attacks occurring in hypophyseal tumors cannot be due to either of these two factors. Therefore, in these hypophyseal tumors, the somnolence must be due to some change in interference with the internal secretion of the gland, and not to mere mechanical pressure, because the somnolence only occurs in tumors which interfere with the glandular

cells and not with the bipolar nerve cells. Sometimes in pituitary tumors (hypersecretion) insomnia takes place, and these insomnias have been known to yield to the administration of pituitary extract. These insomnias may take place in pituitary tumors in spite of marked pressure symptoms, a proof that somnolence under like conditions cannot be due to pressure. Insomnia is most likely to take place if there is hyposecretion, while somnolence occurs where the gland is over active (hypersecretion).

He then goes on and points out the analogy between the winter sleep of animals and our daily sleep. The winter sleep of animals is also due to diminished secretory activity of the tissues, or to a special tissue which he terms "*la glande hibernale*," an important organ which is found almost exclusively among hibernating animals. This gland is located near the thymus, and is small in summer when the animal is active, grows larger and becomes very vascular at the approach of lethargy, while during sleep, it gradually diminishes with the exhaustion of the activity of the gland. Awakening takes place with an increase in the activity of the gland.

According to this secretory theory of sleep,—“sleep is a vegetative function, a function of secretion, intimately related to the functional activity of certain special organs of internal secretion. Deep sleep is an absolute repose of the brain, in which there follows a slowing of the metabolic process of the entire body.” It is the nerve cell which furnishes the key to sleep. The nerve cell, particularly by means of its Nissl bodies, nucleus and neurofibrils, has an important secretory function, characterized by an elaboration of the chromatic substance. This chromatic substance accumulates during repose and disappears during activity, particularly under conditions of fatigue. It is logical to infer, therefore, that during sleep there ought to be a large accumulation of this chromatic substance in the nerve cell. By means of the Nissl method, he examined the brains of chickens which he suddenly killed during sleep, and he found that the cortical nerve cells showed a marked affinity for basic dyes, due, he thinks, to an increase of the chromatophilic elements in the protoplasm. In the experiments of others it was also shown that there was an increase of the chromatic elements in the cells of the spinal ganglia during sleep and a diminution during activity. Therefore the condition most favorable to an accumulation of the chromatic substance is a state of repose of the nerve cell, exactly the condition which is found in sleep. As the chromatic substance accumulates in the nerve cells during this functional inactivity, so sleep is most easily obtained in a state of functional repose, when the sensory stimuli

which bring the nerve cells into activity are most feeble, that is, disinterest for the present situation, and therefore repose of the psychical centers. The chemical modification which makes the nerve cell lose its activity during sleep is probably a process of dehydration and therefore a decrease in its electric conductivity. The waking from sleep is due to an increase of osmotic pressure, and therefore hydration. This chromatic material is colloidal in nature and colloids are the physical basis of living matter. This accumulation of chromatic elements in the nerve cell forms an obstacle for the transmission of psychic stimuli. There follows consequently a loss of attention, then unconsciousness, then sleep, which is a negative state, a cessation of nerve function. Sleep, which bears remarkable analogy to the hibernation lethargy of animals, is also dependent upon an organ of internal secretion. The hypophysis is the nervous center for sleep, but is not a specific organ of sleep, any more than it is a specific organ of adiposity and osteogenesis.

Sleep is essentially a mechanism for the repair of the neural elements, which cease their activity during sleep. Waking activity is the cause of sleep in the same way that starvation and abstinence from food prepare the way for the digestive function. In sleep the chromatic substance increases, and this increase produces a diminution of the osmotic pressure of the nerve cells. Then the formation of the chromatic substance gradually decreases, finally reaching a point where it permits the passage of psychic stimuli. Thus a preparation is made for awakening and sleep ceases when this activity is exhausted.

In discussing the origin of sleep, he points out that sleep is necessary for the preservation of the organism, because waking activity if prolonged would have a necessarily fatal effect. According to Salmon, sleep is a defence of the organism against intoxication, and has, therefore, a positive, reparative action upon the nerve centers. The best explanation of sleep is therefore biochemical, which considers sleep, not a vegetative passive state, but a positive condition, a function.

I. H. CORIAT.

PHYSIOLOGY OF THE SPECIAL SENSES. By *M. Greenwood, Jr.*,
London. Edward Arnold. 1910. Pp. vi, 239.

THE book before me purports to be of aid to the psychologist and also to a general class of readers who would like to know something about the special physiology of the sense organs. I shall consider it only from the psychological side.