it can perforate a healthy intestine is still under discussion, though a case recently reported⁵ seems to show that it may. Askanazy attempted to show the same relationship true between ascaris and appendicitis as between trichocephalus and appendicitis. He found in the intestinal epithelium of the ascaris a substance which reacted to the test for iron as did that in the trichocephalus. He felt, therefore, that this iron came from the blood of the host and that it was fair to assume that the ascaris obtained it by sucking. To suck the blood the mucous membrane must necessarily be wounded and entrance given to bacteria.

When we come to oxyuris vermicularis the proof of its etiological value as a cause of appendicitis other than through its presence as a foreign body is lacking. The pin worm is generally considered an inhabitant of the rectum and lower bowel. That this is not true is shown by the frequency of its occurrence in the appendix in a series of 200 autopsies done by Still⁶ on children under twelve years of age. He found pin worms in the intestine in 38 cases, in 25 of which they were in the appendix. In 6 they were in the appendix and nowhere else in the large intestine or cecum. Sometimes they were in the extreme tip of the appendix, as in the case of a boy who died of pleurisy and pericarditis. Before death he complained of pain in the right iliac fossa. At the autopsy the appendix was found thickened and in a condition of catarrhal inflammation. In the distal 2½ inches 111 worms were found. No others were found anywhere in the intestine. Some have thought that oxyuris might wound the mucous membrane of the appendix by its wriggling and thus by furnishing a place of entrance to bacteria cause an attack of appendicitis. This theory as far as I know has not been substantiated by any definite proof and remains therefore simply as an attempt to carry the relation that exists between the other two parasites and appendicitis to oxyuris vermicularis.

Conclusions. — An intestinal parasite causes appendicitis through its presence as a foreign body; and by its struggles may bring about a perforation and peritonitis where otherwise none would have occurred.

Trichocephalus dispar has been proved to cause an inflammation of the appendix by injuring the mucous membrane while sucking the blood of the host.

Ascaris lumbricoides has been found with a material similar to that in trichocephalus in its intestinal epithelium and therefore is supposed to attach itself to the mucosa of the host and

thus start the processes which result in appendicitis.

Oxyuris vermicularis. No proof has been found that this worm causes an appendicitis except in its rôle of foreign body.

Clinical Department.

A CASE OF SARCOMA OF THE LOWER JAW.¹

BY F. B. LUND, M.D., BOSTON.

The accompanying case is reported in order to illustrate the slight amount of deformity which results from subperiosteal excision of the body of the lower jaw, from the lateral incisor tooth to the angle. The patient, a young girl of thirteen, was operated upon on the 16th of July, 1903, for a sarcoma of the lower jaw, which began in the medulla, had caused a fusiform swelling of the lower jaw, had forced its way upward through the alveolar process and grown up into the mouth

¹ Contributed to the fourteenth series of Medical and Surgical Reports of the Boston City Hospital.

Case of Sarcoma of the Lower Jaw.

⁵ Agrer; J. Am. Med. Assoc. Feb. 28, 1903.

⁶ Brit. Med. J. 1899, Vol. i, 898.

Case of Sarcoma of the Lower Jaw.

as a papillary growth which had been considered an epulis. The anterior molars and rear bicuspids had been pushed inward and loosened by the growth. The patient was etherized, and the portion of the tumor projecting into the mouth removed with scissors, and examined by the microscope, proving to be a giant-celled sarcoma. An external incision was then made along the body and angle of the jaw, the facial artery divided, the periosteum peeled from the entire body of the jaw and the jaw divided by a saw and cutting forceps just anterior to the socket of the canine tooth in front and at a point on the ramus of the jaw level with the alveolar process behind; the submaxillary gland was removed; the lingual and hypoglossal nerves were exposed during the operation and the latter, having been partially divided, was sutured with fine silk; all suspicious bits of tissue were cut away, divided muscles and deep tissues sewn with catgut, and the skin incisions with a subcutaneous horsehair suture; the mucous membrane was sewn inside the mouth with catgut. The patient made an excellent recovery from the operation, but the wound suppurred, un-

doubtedly owing to the infection from the mouth, and had to be drained. It finally healed well, however, and the periosteum has evidently replaced the lost bone to a very considerable extent, as a firm, bony bar can be felt in the place of the bone which was removed. The photograph shows that the deformity on front view is very slight; the scar on lateral view is rather deep and conspicuous, but will fade out considerably in time, after which the deformity on side will be much less marked.

CHRONIC CYANOSIS FROM ACETANILID POISONING.

BY J. N. HALL, M.D., AND H. R. MCGRAW, M.D.,
DENVER, COL.

IN view of the cases of "Chronic Cyanotic Polycythemia" reported by Hall in *American Medicine*, June 27, and of Chronic Cyanosis with Polycythemia and Enlarged Spleen," by Osler, in the August number of the *American Journal of the Medical Sciences*, these cases of cyanosis of drug origin are of especial interest. Dr. Osler has informed me that he has recently seen such a case from the use of bromoseltzer, well known to contain acetanilid, which was mistaken for the newly described disease.

The patient was a professional man from a western state, fifty years of age. He stated that he had had bleeding piles for thirty years, during which time three ischio-rectal abscesses had been opened. He had been generally constipated. Although decidedly sallow at times he has had no real jaundice.

He stated that his present trouble dated from the use of large doses of acetanilid for headache, begun six months or more ago. It is probably safe to estimate that he had taken 30 to 40 gr. daily for this period. There was no history of alcoholism. His weight had fallen from 160 to 140 pounds. He felt weak, and suffered from dyspnea, and from palpitation on exertion. Digestion was poor, with occasional vomiting of mucus. His face was slaty in color, and much darker than the cyanosis of advanced emphysema. The mucous membranes were similarly discolored, and the skin of the body to a lesser extent. There was no increased pigmentation over the joints and bony prominences, as in Addison's disease. The color lessened upon pressure, but immediately returned. The temperature rose to about 102° daily. The physical examination was negative so far as the chest and abdomen were concerned. Pulse moderately increased in frequency, and of fair quality. The pupils were somewhat contracted. Several bleeding hemorrhoids and a blind rectal fistula were found by Dr. Sherman T. Brown.

The blood was very dark, with 3,038,000 reds and 18,000 whites. Moderate poikilocytosis existed. The increase in the whites was in the polymorphonuclear leucocytes, doubtless explained by the rectal trouble. The reds were pale, and had a "brittle" appearance. The test with Fleischl's hemoglobinometer gave 30% of coloring matter, though this was chiefly methemoglobin, probably. No spectroscopic examination was made.

The urine had a specific gravity of 1.022, no sugar nor albumen. Its color was dark, and the patient stated