marsupials. Like the platypus, they resemble reptiles in some important skeletal characters and in being oviparous, the eggs being hatched, as in birds, by the warmth of the mother's body. The Monotremata are confined to Australia, Tasmania and New Guinea, and probably had the same distributional history as the marsupials, although up to the present time no palæontological evidence of their former existence in South America has been discovered.

Mr. Pocock having been duly thanked by the President for his most interesting and instructive demonstration, the party dispersed to other parts of the Gardens.

REPORT OF AN EXCURSION TO THE LIZARD, CORNWALL.

MARCH 20TH TO 26TH, EASTER, 1913.

BY J. S. FLETT, M.A., LL.D., D.Sc., F.R.S., F.G.S., AND
J. B. HILL, R.N., F.G.S., *Directors of the Excursion.*

PART I.—BY DR. FLETT.

MARCH 20TH.

THE party left Paddington by the Cornish Riviera Express at 10.30 a.m. for Helston, and from there drove to Coverack, where headquarters were established at the Coverack Headlands Hotel.

An unusually large number of members attended, the official party numbering 82 on several occasions, and many local visitors joined on one or more days.

Miss G. M. Bauer acted as Excursion Secretary during the Excursion.

MARCH 21ST.

The fine rock scenery in front of the Headlands Hotel first attracted our attention, the serpentine weathering out into jagged rocks more or less masked by growth of thorn and Cornish heath. The serpentine here contains bastite in fairly large crystals, which stand out as prominent nodules on the rusty brown surfaces of weathered slabs. Above the hotel the flat-topped Pliocene platform extends for several miles in excellent preservation. As low tide was early in the forenoon we descended by the path to Perprean Cove, where a small boss of dark-coloured troctolite (very rich in olivine but with large spotted feldspars sometimes nearly idiomorphic) occurs in the

serpentine. This troctolite is quite massive and obviously intrusive into the serpentine, while it is in turn cut by veins of gabbro. We next visited the harbour on the north side of the village of Coverack to examine the famous locality for troctolite on the shore below the post office. Here there are two varieties, one having red spots of weathered olivine in a matrix of felspar, while in the other the olivine spots are green, the colour depending on the state of oxidation of the iron in the rock. The sections in this cove are most instructive, for they show four kinds of rock, viz.: serpentine, troctolite, gabbro (including olivine gabbro) and olivine dolerite, and it is quite clear that they were intruded in the order named. The "black dykes" of quite massive dolerite are very fresh, and show no foliation; they are fine-grained with chilled edges against the other rocks, proving that the whole complex had cooled down before their intrusion.

The gabbro, on the other hand, is often exceedingly coarse-grained, with crystals of diallage and felspar three or four inches in length; and in many places this rock shows a well-developed foliation, and passes into a true gabbro-schist. A specially fine section is seen in the rocky beach near the west corner of the cove, where the serpentine is cut by a broad vein of highly-foliated gabbro across which runs a dyke of nearly massive gabbro-pegmatite, while a massive dolerite dyke cuts the whole suite. This proves that the foliation of the gabbro-schist was already complete before the gabbro-pegmatite and dolerite were intruded; in other words the foliation arose either at the time at which the dyke was injected, or very shortly thereafter. The sections of raised-beach platform covered by gravels and sands and topped by stony Head claimed attention, and also the manner in which the serpentine weathers, showing a network of furrows engraved on its surface (see Plate 45.B). We next moved on to the North Corner, where a halt was called for lunch. At that locality there are blocks of olivine-gabbro which are nearly coal-black in colour, and so little affected either by decomposition or by foliation that they are as fresh and as massive as if they were of Tertiary age.

The wet state of the ground after the heavy rains had made it undesirable to proceed to Lowlands Point as intended, and we now followed the road leading to Crousa Common. On each side the dry stone walls are built of gabbro blocks, most of them belonging to the type known as the "flaser-gabbro," and the fields are dotted with large, rough, lichen-covered stones which are the undecomposed masses left behind where the softer parts of the weathered rock have been washed away by the rain. Near the crest of the hill the down becomes smooth and regular, and this change marks the limit of the well-known gravel beds that are supposed to be of Pliocene age. Proceeding along the Helston road we visited several pits in these gravels, and found

the pebbles to be mostly vein-quartz with occasionally quartz-schist rock and sometimes a bit of rotten flint. No fossils have been obtained from these pits, but from the position of the gravels on the summit of the Lizard platform, and the analogy with the gravels of Polcrebo and the St. Erth Beds, the inference that they are early Pliocene seems fairly strong.

Half a mile farther west is situated the old cromlech or dolmen known as the "Three Brothers of Grugwith," consisting of three gabbro blocks, one of which shows cup-markings. This provided the subject of some very interesting remarks by Mr. Reginald Smith. The downs around the dolmen are cumbered with great blocks of gabbro and afford a typical example of the scenery that characterises this formation where the land has not been taken under cultivation.

Our homeward route lay entirely on the smooth Lizard platform, here exceedingly well preserved, and about half a mile north of Penhallick we paused to gather specimens of the massive black gabbro which are easily obtained by the roadside.

MARCH 22ND.

In heavy rain a start was made for Porthallow, and it was fortunate that covered omnibuses had been secured for the journey. The cove was reached by 11 o'clock, and to take advantage of the tide it was decided first to visit the section of Veryan rocks under the north cliffs. This is the best section of this group to be obtained on the mainland in this part of Cornwall, though the pillow-structure in the lavas is by no means so well seen as on Mullion Island. The Lizard boundary fault is exposed in section east of the inn; it dips south at 45° , and there is a broad zone of crushed breccia between the Lizard rocks on the south and the Palæozoic rocks to the north.

The rock on the north of the breccia is crushed Veryan "killas." Then comes the sand and gravel of the beach. Beyond this we found several small quarries in white Veryan quartzite, much veined and broken, though completely re-cemented and firmly compacted. The section in the north cliff then begins with shattered dark-grey and black killas, followed by quartzite and a large mass of pale-grey soda-granite-porphry, an intrusive member of the "spilitic suite" of rocks. Beyond this comes more quartzite, some killas, and finally a high cliff of green, much-sheared pillow-lava. To get into Nelly's Cove it is necessary to scramble over the blocks strewn on the shore and the much-eroded platform of the raised beach. Just before entering the cove we meet with grey killas containing large lenticles of white Veryan limestone, which weather in a honeycombed manner; this limestone, though for the most part highly crystalline, has yielded Ordovician fossils. In the centre of the

cove the radiolarian cherts are seen as thin black bands in the dark shales, and the gravel of the beach is full of pebbles of this rock. The north side of the cove shows a magnificent section of raised-beach gravels covered with head, (see *ante*, Plate 35.A, p. 148). As we returned to the inn we paused to note the remarkably fine development of shear-structures in killas, limestone and pillow-lava, which marks the south side of the cove. The rocks are torn up into lenticles by planes of movement which dip to south or south-east.

The weather by this time had much improved, and after lunch we proceeded to the south side of the cove to examine the section of the Lizard rocks. Of these a great variety may be found within a comparatively short distance, but the sequence is broken by faults, and little certainty can be attained regarding the geological structure. South of the boundary fault we first met with brown and greenish mica-schist, in which occurs a sill of augen-gneiss, great fallen blocks of which lie on the beach. Next comes thin-banded hornblende-schist with fine yellowish epidosite; then we pass a great landslip, which cloaks the face of the cliff with débris, and is extensively used as a source of building-stone. In this occurs gabbro, serpentine and coarser hornblende-schist. The gabbro is evidently a vein in the serpentine, but the other two rocks are folded together in intricate fashion, and sometimes occur interbanded. The serpentine is all fine-grained, with a rather resinous lustre—the true marginal type, which was originally a fine olivine-rock or “dunite.” Several good folds, involving hornblende-schist and serpentine, were seen in the rock-platform of the raised beach. (See *ante*, Plate 35.A, p. 148.)

Thereafter taking the road from Porthallow to St. Keverne we passed over a long outcrop of the Old Lizard Head Series, and roadside pits were inspected, containing garnet-biotite-gneiss (paragneiss), fine grey mica-schist, silky fine hornblende-schist, and, near Trenance Farm, quartz-hornblende-granulite, showing good bedding, though highly metamorphic. Footpaths were then followed to Treleague farmyard, where the best exposure of the Treleague Quartzite was seen, a much less metamorphic rock than the quartz-granulites of the Old Lizard Head Series, though, like them, containing hornblende. From this point the road led over the gabbro by St. Keverne into Coverack.

MARCH 23RD.

In the morning some of the party inspected the bands of chromite-serpentine in the bastite-serpentine in Perprean Cove, and then proceeded to the shore on the south side of the Headlands Hotel, where many interesting structures are to be observed.

The raised-beach deposits and Head are both in evidence, resting on serpentine, which is cut by many dykes, large and small, of gabbro and olivine-gabbro. One of these veins terminates by splitting up into a number of radiating branches, like the ribs of a fan, and another is penetrated by a fine-grained black dyke which runs for some distance in the centre of the gabbro dyke.

Later in the day a large party walked across the downs to Poldowrian to see the great dyke of gabbro that forms the promontory of Carrick Luz. Descending at the south end of the mass we saw the intrusive junction with the serpentine on the east side, where the older rock is filled with gabbro veins, and then returning to Sperric Cove on the north-west side those who were in search of specimens filled their bags with augen-gabbro (Plate 45.A), gabbro-schist, and fine examples of the white gabbro, which contains nearly colourless crystals of chrome-diopside in a matrix of white labradorite.

The special feature of Carrick Luz, however, is the manner in which the great dyke shows injection foliation, for the gabbro is far more highly foliated than the serpentine into which it was intruded, and the augen structure and "rodding" are parallel to the length of the dyke, and cut across the same structures in the serpentine.

PART II.—BY MR. J. B. HILL.

MARCH 24TH.

The party drove from Coverack to Lestowder, where a large exposure of Veryan quartzite was examined, as well as the Veryan killas associated with it on the coast section below Penare. Along that section the killas is in an advanced condition of metamorphism, the argillaceous members being represented by lustrous greenish phyllite showing incipient spotting, and the siliceous members by granulites. Crush breccias are also much in evidence.

The coast section between Nare Point and Flushing Cove was then examined, consisting mainly of the Manaccan series (Lower Devonian), the various members, from coarse conglomerates to fine clay slates, being admirably exposed. At Polnare Cove the Veryan series is seen overlain by the Devonian. On the beach due north of Lestowder a dyke of mica-trap cuts the conglomerate. On Men-aver-Beach there is a capital exposure of the Pleistocene beach a few feet above high-water mark and capped by an extensive platform of Head.

The party left the coast at Flushing and walked to the head of Gillan Creek, visiting en route the exposure of Manaccan

hornblende-schist at Tregithey. At the head of Gillan Creek the hornblende-schist, largely in the condition of a crush breccia, is well exposed, and the mica-schist on its flanks was examined. At Carn a good exposure of Veryan sandstone was seen, thoroughly brecciated, but the rock is nevertheless sufficiently solid to be quarried for macadam. Along the valley leading from Manaccan to Gillan Creek the Veryan phyllite, finely spotted with biotite, was examined, and on the roadside immediately south of Manaccan a small exposure of granophyre was seen lying in crushed killas of the Veryan series.

After tea at Manaccan the party drove back to Coverack.

PART III.—BY DR. FLETT.

MARCH 25TH.

A fine morning saw us on the way to Poldowrian, where, passing to the west of the farmhouse, we descended the cliff at the second cove to view a beautiful section, showing black dykes and gabbro veins in the serpentine. This cove has been figured in the Lizard Memoir. The fluxion-banding of the serpentine is cut by veins of gabbro, some highly foliated, while others are nearly massive. Across these come two black dykes, one rather schistose, the other almost a normal dolerite. The local character of the foliation in these dykes is very noticeable. Still more important is the fact that the upper of the two black dykes contains a number of veins of red granite, thus indicating a transition to the composite Kennack gneisses.

From this point we followed the cliff-path to the east side of Kennack Sands, where again we descended to the beach. At this locality broad bands of chromite-serpentine are visible in the bastite-serpentine. A large dyke or inclined sill of epidiorite occurs just at the east end of the sands, and contains huge inclusions of serpentine. A few yards farther east we meet with veins of talc and of pseudophite (Plate 42.B).

In the rocky beach below the cliff there is a small boss of red granite-gneiss, well foliated parallel to its junction with the serpentine. At its margins it passes into a darker rock, sometimes banded, and it seems to give off veins of dark epidiorite, with strings and veinlets of red granite. This little boss may be regarded as a type of the larger granite-gneisses of the Lizard, which emit sills and dykes of composite, veined and banded gneiss.

Crossing the eastern sands we found in the centre of the bay a serpentine headland, carrying on its forehead a thin patch of raised-beach gravels, and showing black dykes and fluxion-bands of chromite-serpentine.

Kennack Corner, on the west side of the sands, is one of the most famous geological sites in the Lizard. A sketch map of this locality is given in the Lizard Memoir. The principal feature of this section is the banded gneiss, which forms a group of picturesque stacks and ledges on the foreshore. The characters of this group are nowhere better developed than in this place. A dark component, which may be described as a hornblende-schist or epidiorite, forms streaks and bands enveloped in or alternating with a pale-red or grey acid gneiss. Sometimes the basic material appears as blocks in the acid matrix, but these inclusions are seldom well defined with angular outlines; they are practically always rounded or lenticular, as if they had been drawn out or moulded by the flow of the hot gneiss. In colour they range from black or very dark green to paler shades, which indicate permeation by the acid magma. These blocks are veined by the gneiss in such a manner as makes it certain that they were solid before incorporation in the complex. In one or two places the basic rock contains large phenocrysts of felspar. Another type shows nearly parallel fluxion-bands of dark and light-coloured rock; in this case the dark ingredient is not so basic as in the rounded blocks, but must be regarded as a mixture of the original basic and acid materials. The junction of the gneiss with the serpentine was well exposed owing to the recent storms having stripped the beach exceptionally bare of sand, and it was seen that the flow-banding of the composite gneiss follows the line of junction with the older rock. Several inclusions of coarse bastite-serpentine (from 2 to 6 ft. in diameter) were pointed out, and it was remarked that the banding of the gneiss wrapped round the surface of the inclusions. A few yards south of the gate at the west corner of the sands a dyke of banded gneiss was pointed out, cutting the serpentine and also intersecting a dyke of coarse pegmatitic gabbro. In this dyke the flow-banding is very obvious, and is parallel with the cheeks of the fissure which the dyke has filled, though its course is at right angles to the banding in the main body of the gneiss from which it can be shown to have originated. In the same way, at a point about seventy-five yards south of the gate, the junction of gneiss with serpentine changes its direction from south to west, and the banding of the gneiss swings round in conformity with it. These facts establish that the banding is really due to fluxion of a composite (basic and acid) magma intrusive into the serpentine.

Among the other interesting features of this remarkable shore section were the black dykes (epidiorite), one of which cuts across a gabbro vein that is three times faulted within the space of a few yards, while another black dyke contains a mixture of red granitic material that indicates an affinity to the composite Kennack Gneisses. In the banded gneiss there are veins of coarsely crystalline pegmatite consisting of quartz and felspar,

but passing, near their terminations, into veins of pure quartz. The serpentine contains veins of asbestos that have been known to Cornish geologists for over a hundred years. No part of the Lizard is more full of interest than the west side of Kennack Cove, and its rocks have been referred to so often in the literature of Lizard geology that merely to give a list of references would exceed the limits of space available in this account of the excursion.

From this point southwards we followed the beach below the cliff for about half a mile in order to examine the extraordinary variety in the banding and veining of the Kennack Gneisses. Ascending the grassy slope of Thorny Cliff we then directed our course to Polbrean Point, where a small party descended the steep path to see one of the most instructive sections of the Lizard intrusive rocks. The serpentine is there intersected by gabbro-veins which in turn are invaded by a black dyke. The last named can be seen to originate from a mass of fine, dark material (epidiorite), which on close inspection proves to contain threads of red granite, and, followed a little farther, merges into a typical intrusion of veined and banded gneiss. The sequence can accordingly be demonstrated in an excellent cliff section only a few yards long and perfectly accessible. It is (1) serpentine, (2) gabbro, (3) epidiorite, (4) composite gneiss. The south side of this little cove is defined by a splendid sill of Kennack Gneiss cutting obliquely across the flow-banding of the serpentine, and running up the cliff to its summit. This sill is figured on Plate 31A, p. 128. It contains many lumps of serpentine which are usually massive and coarse-grained at their centres, but have an outer zone of fine silvery talc-tremolite-schist.

After a short stop at the old serpentine works in Caerleon Cove we hastened on to Kildown, where there is a large caldron-shaped subsidence in the face of the cliff, produced by the collapse of the roof of a cave at some unknown date. Then we visited the famous quarry at Kildown about which there has been a good deal of discussion, but the section is by no means a convincing one. At the "Flagstaff Quarry" above Cadgwith for the first time we made acquaintance with the fine-grained tremolite-serpentine, while on the cliff below the banded epidotic hornblende-schists now make their appearance. In Cadgwith Cove these hornblende-schists are beautifully exposed, dipping in a north-west direction, but the tide was too high to allow us to do justice to this section. After tea a number of the party visited the famous Devil's Frying-pan, a large caldron entered by the sea through an arch on its eastern side. Then we met Mr. Howard Fox, who had walked over from the Lizard to join the excursion. At this locality the three main varieties of serpentine occur in close conjunction, for the bastite serpentine is seen in the farm-yard, the tremolite-serpentine in the interior of the Frying-pan,

[Photo by T. C. Hall.]

A.—THE RAISED BEACH, BELOW LOWLANDS POINT, AS SEEN FROM COVERACK.

[Photo by J. S. Flett.]

B.—KENNACK SANDS, WITH CARRICK LUZ AND THE BLACK HEAD IN THE DISTANCE. (See p. 318.)

and on the east face of the cliff, at the north side of the arch, there is a small outcrop of fine flinty "dunite-serpentine" in a curious rock which consists of hornblende-schist penetrated and veined by banded Kennack gneisses. After a full day's work in excellent weather we made our way by the field paths from Cadgwith to Lizard Town, to find comfortable quarters awaiting us at Hill's Hotel and Housel Bay Hotel.

MARCH 26TH.

A visit was first paid to Housel Bay, which is surrounded by rugged cliffs of dark hornblende-schist. At the steps leading down to the cove there is a slanting dyke of reddish felsite, much crushed and silicified. The Lion's Den on the west side of the bay is a caldron that originated through the collapse of the roof of a cave or "hugo" on the 19th of February, 1847. The Bumble Rock is a large pillar of hornblende-schist that guards the south-west corner of the bay, and just beside it we examined a little patch of raised-beach gravels lying in a corner of the face of the cliff about 25 ft. above high-water mark. It is the highest, and probably also the oldest, raised beach in the Lizard peninsula.

At Polpeor, the Lifeboat station, we descended to the beach to examine the fine section of crumpled garnetiferous mica-schist. Here occur porphyritic basic dykes which have been torn apart into pillow-shaped masses by the movement in the schist around them. Mixed with the brown mica-schist there is a good deal of fine silky green schist, containing fibrous hornblende. Thin beds of fine grey quartz-granulite occur among the schists, the whole presenting a very typical section of the members of the Old Lizard Head series, which are the oldest rocks in this quarter, and perhaps the oldest in England. The green schists are filled with bands and segregations of epidote, and are often intensely folded. On the beach we gathered pebbles of the hornblende-gneiss of the Man of War Rock and other skerries south of the Lizard Point, with its characteristic sinuous or corrugated foliation.

A few hundred yards farther west, in Pistil Ugo Cove, the brown mica-schist, green schist, and quartz-granulites are exceedingly well seen; their bedding is well preserved though sometimes much contorted by folding. It is remarkable to find the stratification so obvious in rocks that have been so completely recrystallised and have lost entirely their minute clastic structure. The next halt was made on Old Lizard Head, where a party descended the cliff to view the large sill of fine granulitic granite that represents the apophyses of the Man of War gneiss intruded into the sedimentary schists. In the small quarry at Venton Hill Point we gathered specimens of the garnetiferous mica-

schist containing andalusite, and saw also a large lenticle of uralitic ophitic diabase, probably a fragment of a dyke that had been severed by movement, like the greenstone pillows at Polpeor.

From this point we pushed northwards to see the boundary fault which brings in the serpentine at the south end of Pentreath Beach, and descending to the shore by the rock steps we found it to be a thrust hading to the north at an angle of 45 degrees. It is consequently a thrust of different character from that which we had seen on a previous day at Porthallow, which hades to the south. Presumably the thrusts at Pentreath Beach are of late Silurian age, while the Porthallow thrust is one of the late Carboniferous group. The serpentine and hornblende-schist are very much brecciated at their junction, and as we followed the beach northwards a number of minor thrusts, parallel to the main one, were seen in the cliffs. At the north end of Pentreath Beach intrusions of banded gneiss and black dykes were visible, one mass of red granite with dark-coloured margins being exposed on the path by which we ascended to the top of the cliff. Regaining the level of the Lizard platform we halted for a few minutes to view the great landslip or collapsed mass known as Holestrow, and after collecting specimens of chromite-serpentine (of yellow colour with black specks of chromite and picotite) at the Yellow Carn we descended into Kynance Cove (Pl. 43.A).

This cove itself, besides being famous as one of the best pieces of cliff scenery in Cornwall, possesses sufficient geological interest to occupy a geologist for more than one day, but as the tide was rising our stay had to be somewhat curtailed. Above the cove a streamlet cuts a deep narrow valley in the serpentine, with characteristic rocky slopes partly mantled with heather. The lower part of this valley has been submerged and flooded by the sea in comparatively recent times, and the continuation of the old land-valley is visible in the green slope which forms the east side of Asparagus Island. On the west the sea has pierced the walls of the valley, entering probably along lines of fault and intrusions of red granite and gneiss, which are comparatively easily cleared out by the waves. In this way the islands and rock pillars, such as the Steeple Rock, have originated. The Lion Rock on the east side is cut off from the Yellow Carn by a deep gully formed by the removal of the crushed rock that occupied the line of an important fault.

This fault passes up the cove and along the east side of the valley. It separates the bastite-serpentine on the east side of the cove from the tremolite-serpentine on the west. The latter is a very interesting rock, finely granular and usually rather schistose. Its banding is often faint, except on weathered surfaces. At Kynance many specimens of the tremolite-serpentine may be gathered on the beach, showing large elliptical augen of enstatite.

A.—KYNANCE, SEEN FROM PENTREATH. THE LION ROCK AND THE
BISHOP ARE THE ISLANDS. (See p. 322.)

B.—CONTORTED BANDED MYLOR BEDS, WITH QUARTZ VEINING, PORTHLEVEN.

A similar structure, but on a larger scale, is present in the bastite-serpentine at Pentreath Beach.

The principal features of geological interest at Kynance are the intrusions of granite and of banded gneiss near the Steeple Rock. One of these is seen as a rounded mass of red granite above the western cave; it has a narrow "aureole," in which the serpentine is altered to pale-green talc-tremolite-rock. In the centre of the cove other masses of red gneiss occur, and there are two small dykes of crushed gabbro. Remains of the raised-beach platform may be seen in several places, but no trace of the raised-beach gravels remains. On the east side of the cove there are several black dykes, and at one place a large white block of flint-like pseudophite lies on a ledge on the face of the cliff. Some of the party visited it and took specimens.

We left Kynance and ascended to the top of the downs by the cart-road, and on our way noted several intrusions of red and grey gneiss and of dark fine-grained rock which resembles the basic part of the Kennack Gneisses.

Above the bend of the road a halt was made to observe a beautiful branched dyke of dark hornblende-schist in the coarse bastite-serpentine. Opportunity was taken by Dr. Flett to contrast this basic dyke with those previously seen at Polpeor and Venton Hill Point. The latter were evidently tough resistant masses which had been dragged apart to form elliptical blocks, but still retained their original internal structure in some measure. The dyke at Kynance, on the other hand, had yielded readily, and had become schistose throughout. It was suggested, as the probable explanation of the different manner in which these dykes had behaved, that some of them were cold and hard when the pressures were initiated, while others were still in a warm and plastic condition, and consequently readily became foliated.

PART IV.—BY MR. J. B. HILL.

MARCH 27TH.

The members drove from the Lizard to Gunwalloe. The Manaccan series was examined along the coast at Jangye-ryn where the highly folded strata (Pl. 34. A., p. 138), and formerly described by Sedgwick, were inspected. At Baulk Head the beds yielding plant remains in the same series were examined; the best exposures, however, are now buried beneath the shingle, but a few members of the party succeeded in obtaining specimens with plant fragments. The coast section was examined as far as Porthleven, the Falmouth series extending from Chyanvounder to the Loe Bar, after which it is replaced by the Mylor series. Along

this foreshore relics of the Pleistocene beach are seen, capped occasionally by Head. The modern beach shingle is conspicuous by the large proportion of flint and chert pebbles; this shingle deposit fronting the Loe Estuary has formed a bar, converting what was originally a tidal estuary into a fresh-water lake. Approaching Porthleven the Mylor beds assume their characteristic banded appearance, and are, moreover, lying in the aureole of the Godolphin granite and are strongly knotted. The sills of greenstone so common in this part of the Mylor series were examined to the south of Porthleven Pier, after which the party crossed to the north side of Porthleven Harbour and inspected the enormous erratic on the foreshore known as the Giant's Rock and consisting of microcline gneiss of a type unknown in Britain (Pl. 35.B, p. 148).

The party then drove from Porthleven to Helston, where, having had tea, the return drive was made to the Lizard.

PART V.—BY DR. FLETT.

MARCH 28TH.

Although the set programme of the excursion was now completed, a considerable number of the party had decided to stay for a day or more, and Dr. Flett arranged to lead them along the west coast to Mullion, returning at night by motor-bus to the Lizard. The weather was showery at first, with a heavy sea coming in from the south-west, which gave us an excellent opportunity of seeing Kynance Cove in a storm. Great waves were breaking right across from the Lion Rock to the Bishop, and thick sheets of spray sweeping inland up the valley as we passed on our way to The Rill. At Lewarnick several dykes of granite were pointed out in the serpentine, and at The Rill we gathered specimens of the tremolite-serpentine, for which this is a well-known locality. At Gew-graze a visit was made to old workings in the "Soap Rock," which was at one time extracted for the manufacture of pottery. It proved to be a vein of pink granite that had been "steatitised" or impregnated with magnesia derived from the serpentine. In its typical state the "soap rock" is quite soft and plastic; this material is merely a variety of the pseudophite which we had already collected in several places.

We next proceeded by Vellan Head to Pol Cornick, but on the way one of the party was bitten by an adder, and arrangements were made to have him sent to the nearest farmhouse while a doctor was summoned from Mullion. It is sufficient to say that, though considerably inconvenienced for a day or two, he soon recovered and was none the worse for his experience.

At Potstone Point a halt was made for lunch, and the more active members of the party descended the grassy slope to see

[Photo by J. S. Flett.]

A.—EPIDIORITE DYKE CUTTING FAULTED GABBRO
DYKE IN SERPENTINE, KENNAK. (See p. 319.)

[Photo by J. S. Flett.]

B.—OLD LIZARD HEAD, SEEN FROM VENTON HILL
POINT. (See p. 321.)

the junction of the banded serpentine with the hornblende-schist, and the dykes of epidiorite and of "potstone" which traverse both of these rocks. Nice specimens of fine flinty-looking dunite-serpentine were collected here.

After passing for about six hundred yards along the cliffs of hornblende-schist we paused at Parc Bean to see masses of very fine-grained dunite-serpentine in the hornblende-schist, and by descending to the margin of the cliff a fine exposure was visited showing an anticline of serpentine wrapped round by the schist, both rocks possessing the same banding and foliation.

This cove is also the best locality in England for the mineral datolite. It occurs in the rocks below high-tide level, and some nice specimens both of the crystallised and the botryoidal variety were obtained by those of the party who searched for them.

On Predannack Head the rough gnarled foliation of the hornblende-schist attracted attention, and soon thereafter we came in sight of Mullion Island, but the day was so stormy that there was no hope of making a landing. The structure of the island was explained, but at that distance the pillow-jointing in the lava cannot be seen except with a good glass. A descent was made into Mullion Cove, where we visited the famous cave and inspected the boundary thrust, by which the serpentine has been forced north over the hornblende-schist. Passing round the harbour we ascended the path before the hotel, and marked in the cliffs a lenticle of fine serpentine in the hornblende-schist. A visit was then paid to the rocky promontory of Henscath, where a small boat-shaped infold of hornblende-schist is visible in fine dunite and tremolite-serpentine. After tea at the Mullion Cove Hotel we drove back to the Lizard.

MARCH 29TH.

Those who remained, about a dozen in number, set out for a walk along the eastern cliffs from Housel Bay to Cadgwith, and as the tides were getting late an early start was not desirable. The first point visited was Pen Olver, where Dr. Flett pointed out the lenticular masses of red mica-schist that occur in the epidiotic hornblende-schist, and called particular attention to one locality in which he had found sillimanite-mica-schist. Around Bass Point to Church Cove we followed the cliff-path, and noted in many places beautiful veins and bands of pale-yellow epidote and pyroxene in the dark hornblende-schist. North of Church Cove the serpentine is brought against the hornblende-schist by a thrust-plane that dips to the north, and is the continuation of that already seen on the west at Pentreath Beach.

In the well-known serpentine quarry of The Balk some time was spent in collecting specimens of many-coloured ornamental serpentine, some of them marked with blood-red spots, while

others were pale-yellow and green. A little distance to the north lies Pen Voose Cove, one of the most famous parts of the Lizard in geological literature. After descending to the beach we found a most interesting variety of rocks both in the cliffs and in the beach-pebbles. The south wall is serpentine, with veins of gabbro and a beautiful arched sill of banded gneiss. In the centre of the cove there is much banded gneiss, some of it filled with inclusions of coarse bastite-serpentine, some of which have assumed a curious pink colour. Large dark epidiorite blocks are scattered through this gneiss, and around them flows a banded matrix. A photograph of this is given on Plate 31. B. (see *ante*, p. 128). In the northern part of the cove an extraordinary jumble of rocks is seen. There is a good deal of serpentine, which is penetrated by veins and irregular masses of gabbro. Red gneiss occurs also, and seems to form dykes, cutting all the other rocks, but associated with it is a dark diorite or massive epidiorite, and there are rocks that have been called "mica-diorite," which seem to be mixtures of the two last named. This curious complex is part of the "mélange" which occurs in the cliffs for half a mile to the north of Pen Voose, and has presented an insolvable riddle to many geologists. It is very difficult, indeed, to explain how the rocks have become intermingled, and almost impossible to arrive at a correct conception of the sequence of intrusions. But the rock-specimens are excellent, and for the collector this is a happy hunting-ground.

As the tide was low we were able to get round the northern promontory and enter the cove below The Chair. The sections continue to be exceedingly intricate, but at one point there is a junction of the gabbro with the serpentine. The rocks are faulted against one another, and the remarkable fact is that the gabbro, which is massive at a distance of a foot or so from its margin, passes into gabbro-schist where it meets the serpentine. The most probable explanation of this is that this fault was in movement at a time when the gabbro was *still hot*, for in innumerable places in the Lizard district faults cut the gabbro but produce no change in its structure. The serpentine at this locality is converted for a short distance into pale green anthophyllite-schist, but the main mass is coarsely crystalline and shows no foliation. Many dykes of gabbro and of epidiorite were observed in the serpentine. Ascending the slope we took the cliff-path to Carn Barrow, and on the way Dr. Flett pointed out the sills of gabbro and of banded gneiss that were visible from above, but justice can be done to the coast sections only by visiting them with the help of a boat.

At Carn Barrow the quarries were visited, where hornblende-schist, tremolite-serpentine and dunite-serpentine are all seen together, the dunite being probably intrusive into the hornblende-schist.

[Photo by T. W. Reader.

A — AUGEN GABBRO, CARRICK LUZ. (See p. 317.)

[Photo by D. Leighton.

B.—SERPENTINE WEATHERING, COVERACK. (See p. 314.)

Thereafter we walked to Cadgwith, where a detailed examination was made of the interesting section on the east-side of the Devil's Frying-pan; its special feature is the intrusion of Kennack gneisses into hornblende-schist, and the presence of a small tongue of dunite-serpentine in the complex thus formed. After tea in Cadgwith we walked to Ruan Minor and took the opportunity of examining some sections of the weathered Ruan gneiss on which stands the village of St. Ruan.

REPORT OF AN EXCURSION TO BENTLEY, SUFFOLK.

SATURDAY, APRIL 5TH, 1913.

By P. G. H. BOSWELL, B.Sc., F.G.S., *Director of the Excursion.*

THE official party of the Association, travelling from Liverpool Street Station (G.E.R.), were met at 1.50 p.m. at Bentley by local members of the Association and members of the Ipswich and District Field Club, who had been invited to attend the excursion.

The whole party of fifty then proceeded eastward along the main road, halting for a few minutes in the Crag Pit at Danes Barn, south of the road, about 300 yards east of the station. Geological maps of the district on the scale of 3 inches to the mile (reduced from the Director's Geological Survey on the scale of 6 inches to the mile) were distributed, and the sequence of the beds in the immediate neighbourhood commented on. The London Clay was the lowest bed reached in the valleys, and this was succeeded by Red Crag (with an intervening bank of Coralline Crag at one place only—Tattingstone Hall), Glacial Sand, Gravel, Loam, and Chalky (non-Kimmeridgic) Boulder Clay. River loam and alluvium also occurred. In the pit (Danes Barn) some 30 ft. or more of unfossiliferous ferruginous sands were visible, these being the upper, decalcified portion of the Red Crag, which used to be seen in its ordinary shelly condition at the base of the section, now covered with talus. Members then proceeded to the small pit in the field on the opposite (north) side of the road. Here the complementary phenomenon to decalcification was pointed out. Percolating water, charged with carbonic acid gas derived in part from the atmosphere, but perhaps largely from the soil (which also provides organic acids), dissolves out the shells from the Crag (those made of aragonite disappearing long before the less abundant calcite forms, such as *Pecten*, *Anomia*, *Scalaria*, *Terebratula*, etc.), leaving decalcified sands. If, for any reason, the