from an intermittent heart action, nothing abnormal except the tumor in the right groin. This is a rounded, tense, pulsating swelling, four inches in diameter, extending from the pelvic bone upwards and outwards, filling the right iliac fossa. It is not tender, but pressure upon it causes pain down the anterior portion of the right thigh. On palpation of the tumor a thrill, synchronous with late systole and early diastole, is easily perceived. Auscultation reveals a loud, rough, blowing sound, also synchronous with cardiac systole. There is slight pulsation in the femoral artery of the right leg. There is an area of diminished sensation on the inner side of the right thigh. No tumor could be felt on rectal examination, but owing to the patient's resistance, examination was not satisfactory.

Dr. Balch operated July 27, 1908; Dr. Murphy assisted. Gas and ether were used as anesthetics. A seven-inch median incision was made, the upper end being one and one-half inches above the umbilicus, the rectus muscle split and the peritoneum opened. Exploration showed a large, tense, pulsating tumor, filling the hollow of the right iliac bone and extending from the posterior margin of the pelvis to Poupart's ligament, which it pushed up slightly. The external iliac artery seemed to enter the mass at its upper, inner border. A second incision was made in the abdominal wall at right angles to the first, joining it at the umbilicus and extending across the abdomen to the right flank. The deep epigastric vessels were tied and cut. The flap of the right lower quadrant was then turned down, giving a good exposure of the tumor. The right rectus was divided close to the pubes in order to better expose the lower end of the sac. A Crile clamp was applied to the right common iliac artery and an almost longitudinal incision about three and one-half inches long made in the tumor. Great masses of laminated partly organized clot were removed without hemorrhage until the bottom and lower end of the sac was reached, when there was a gush of blood from the external iliac. With the sac freely opened and dry an opening was seen in the posterior outer side of the external iliac within a half inch of Poupart's ligament. More careful examination now showed that this was the only communication between the sac and the external iliac. The artery was closely adherent to the sac on the inner side, thus appearing at first to enter it at the upper inner angle, but by loosening the clamp the pulsation could be demonstrated down to the opening behind Poupart's. There was some bleeding from this opening, probably from anastomosing vessels. This was closed with fine silk sutures drawn through vaseline and so placed that intima was approximated to intima. The Crile clamp was then loosened until slight pulsation appeared beyond it. There was no bleeding from the repaired artery wall. One side of the sac, which was about 3 mm. thick, had been driven against the ileum so long that the bone had been somewhat eroded and roughened. Into this roughened surface the sac had been forced until it was impossible to dissect it off. On the inner side the sac wall was wrapped about the external iliac vessel in such a manner that it could not be freed. To add to the complications the ureter was in close apposition to the aneurysm. This necessitated some modification of the Matas method. A flap made from the upper surface of the sac was turned down from the inside and sutured with fine chromicized catgut over the repaired artery wall. Another portion from the outside was sutured over this, thus supporting by two layers the line of suture. The cavity of the aneurysm was obliterated as far as possible by bringing its walls together, but this

amounted to so little that a handkerchief gauze was packed in. The walls of the sac were closed about this wick, leaving as small an opening as was consistent with early removal of the gauze. This wick was brought out at the outer angle of the transverse abdominal incision and the peritoneum sutured so that it was extraperitoneal.

The right rectus was sutured with interrupted No. 1 chromicized catgut sutures at the transverse abdominal incision and at the division at the pubes. The abdominal wall was closed by layers. Dry dressing and Dessault splint were employed to immobilize the pelvis and thigh.

The convalescence was practically uneventful. Immediately after the operation the circulation in the leg improved and the pulsation in the dorsalis pedis returned. There has been no return of the murmur nor of the pulsation in the right iliac fossa.

Following the removal of the gauze pack there was for a considerable time a profuse discharge, evidently from the sloughing blood clot and sac wall. The patient, as you see, has fairly good use of his leg and the circulation below the site of the aneurysm is normal.

TWO CASES OF EXTENSIVE THORACIC RESECTIONS: THE FIRST FOR ENCHONDROMA OF THE RIBS, THE SECOND FOR A CLOSED TUBERCULOUS EMPYEMA OF MANY YEARS' DURATION.

BY C. A. PORTER, M.D.

CASE I. J. H., aged thirty-one. Entered the Massachusetts General Hospital July 3, 1908. Past history negative. Present illness: Four years ago struck his elbow against the left side. For several weeks he noticed sharp, piercing pains, which were followed by an irregular lump which was neither sore nor tender. Operation performed April 19, 1906, by Dr. Flanders, of Lawrence. The growth recurred, and two "bony tumors" were removed by him in June, 1907. In November, 1907, another recurrence took place, with steady growth, dull pain and limitation of lateral motion in the spine. There has been no loss in weight; no cough.

On admission examination showed a well-developed and nourished man; somewhat pale. Heart irregular; pulmonic second much accentuated; occasional murmurs over the mitral area. Pulse 96, temperature normal, no leucocytosis; urine negative. A tumor, the size of the palm of the hand, is situated below and to the outer side of the left nipple, involving the costal arch and adjacent portions of the sixth, seventh, eighth and ninth ribs. The mass was very hard, irregularly nodulated, and its posterior surface could be felt depressing the diaphragm beneath the costal margin. There was no tenderness, no redness or edema, and the unattached skin showed the scars of the previous operations. Examination of lungs negative. Diagnosis: Chondo-sarcoma or enchondroma.

In view of the repeated recurrences and the steadily increasing size of the tumor, it seemed not unlikely that the disease was malignant. The conditions were explained to the patient, and a consultation held with Dr. Fred Murphy with reference to operation and the use of his positive pressure box. I operated on July 14, with Dr. Jones as assistant, and Dr. Garland undertook the etherizing, with the aid of Dr. Murphy and four helpers. During the induction of anesthesia the patient was apparently too deeply narcotized and became cyanotic. Pulse was of good quality. Right lateral decubitus; left arm held upwards and to the right. An incision was made along the left margin of the sternum, from a point opposite

the nipple, sweeping down along the costal margin, and backwards to the axillary line. This flap, containing skin and fat, was dissected upwards and backwards. The external, oblique origin was next divided two inches from the margin of the growth, the ribs were scraped bare of periosteum and the intercostal muscles severed, the vessels being picked up immediately. All of the ribs involved in the growth were then divided at this same level with rongeur forceps, and the tumor and ribs pulled inwards and slightly downwards. The pressure was raised to 8 cm. of water and the pleura freely divided. At once there was partial collapse of the lung, and a recession of the heart and mediastinum to the right. Increased pressure rapidly inflated the lung, which bobbed to and fro in the thoracic cavity. Upon further retraction there appeared, growing freely into the pleural cavity, a large mass, the size of a grapefruit, uncovered by pleura and showing the characteristic translucent whiteness of cartilage. The lung was not adherent. On careful examination it was found that the diaphragm was extensively involved in the growth, necessitating the removal of as much as four and a half inches of its anterior convexity. Both pulse and respiration were immediately and seriously deranged, in spite of increased inflation of the lungs, the heart and respiration becoming very rapid and irregular. The diaphragm was grasped firmly by large "T" forceps, and strong traction made downwards and to the left in order to steady the heart and mediastinum. In a few moments compensation occurred, and deep cyanosis was replaced by normal color. The pulse, though rapid, became regular. The costal cartilages were then quickly divided, close to the sternum, and the tumor removed. Using two pairs of double hooks to hold down the diaphragm, its anterior border was sutured with chromic catgut to the transversalis fascia and muscle, sewing from the median line outwards. The tension was extreme and the suture difficult. The lung was then grasped with blunt hooks and held close to the intercostal muscle below the sixth rib, and several sutures passed on round, curved needles, thus fixing the lung. There was slight bleeding; no emphysema. By this time the patient's condition was not good; the pulse had become rapid, and the breathing slightly labored. The remaining muscles were rapidly opposed, so far as possible, in their normal positions, with interrupted chromic catgut stitches. Next an endeavor was made to make an airtight wound by careful subcutaneous, interrupted and continuous sutures of catgut, and then a most accurate skin suture with silk-worm and continuous horsehair. A small gauze wick was placed over the diaphragm, the lung was fully inflated, and the last stitches tied; the object being to leave an inflated lung, which could not collapse from leakage of air into the pleural cavity. A small piece of gauze was placed over the wound, which was then covered with a large square of rubber dam, and a voluminous dressing and swathe. The box was quickly removed and the patient left lying on the table for half an hour, at which time he was recovering from the ether, with only a trace of cyanosis and a fair pulse of 140.

From the point of view of the operator, Dr. Murphy and Dr. Garland were able to regulate at will the expansion of the lung. Only when the diaphragm was severed was there serious disturbance of the pulse and respiration. On the seventh and eighth day, recovery seemed assured, but to my bitter disappointment, a bronchopneumonia of the right lung developed, and the patient died on the fourteenth day after the operation. On the evening of the day of

operation, the patient's temperature was 101, respiration 60, pulse 160. Oxygen had to be given for alarming occasional attacks of cyanosis, following extremely painful attempts at coughing. On the next day there developed a very severe purulent bronchitis, with a temperature of 103; leucocytosis of 21,000. Examination of the sputum showed pneumococci. In spite of carbonate of ammonia and oxygen, very little sputum could be raised, owing to the pain of coughing, and the mechanical difficulties present. With each cough the left side would bulge out, and thus let up intrathoracic tension. A carefully fitted and padded pasteboard splint helped conditions somewhat, but sufficient pressure to make the apparatus effective cramped the sound lung. No pneumothorax developed. By the fifth day coughing became more efficient and less painful, a considerable amount of tenacious purulent sputum being raised. On this day, also, there appeared effusion in the left chest, with well-marked succussion sounds on coughing. This was apparently caused by the constant friction of the parietes against the ends of the ribs, the skin over these pressure points becoming tender and sore, suggesting an internal decubitus. Over a pint of bloody fluid was aspirated through the flap, with marked improvement in respiration and pulse. Cultures from the fluid were sterile. There were two attacks of edema of the lungs during the first week, both of which for a time seemed critical. These were relieved by atropin and digalen. On the eighth day there was a free discharge of serum from the wound, and in spite of the excessive heat it seemed as if respiration and circulation had at last become adjusted and recovery would take place. At times a persistent hiccough was most troublesome. Throughout the convalescence there was glycosuria from one third to three per cent; for this I could offer no explanation. On July 22, the eighth day after the operation, though the temperature was only 100.5, pulse 130, and respiration 35, the white count was 47,200. On the thirteenth day after operation examination of the chest showed the left lung adherent anteriorly down to the fifth rib and to the ninth behind. Good respiratory sounds, except at the lower fifth behind, where a few râles were present. The right lung: pleural friction rub in the axillary line, with numerous diffuse râles. No dullness or bronchial breathing detected. Without especial rise in pulse or temperature, patient gradually failed on the fourteenth day, the pulse becoming weak and irregular, and cyanosis extreme. Death seemed to be due to exhaustion.

Post-mortem examination showed sound healing of the diaphragmatic wound, to which the lung was adherent. The lower half of the left lower lobe was atelectatic and leathery. Slight bronchitis of the remainder of the left lung. No infection of the wound. There was fibrinous pleurisy extending up to the third rib in front, and the fourth rib behind. On the right, there was fibrinous pleurisy over the whole lower lobe, which was filled with bronchopneumonic nodules. Similar nodules were present, though in smaller numbers, in the middle and lower part of the upper lobe. Cultures showed streptococci and influenza bacilli. Cultures from the heart and spleen were negative. The heart showed no macroscopic lesions.

I find it difficult to come to a definite conclusion with regard to the value of Dr. Murphy's positive pressure apparatus in this case. Considerable time was used up in making the apparatus airtight, and in spite of Dr. Garland's skill I

CASE I. X-ray J. H. Shadow of enchondroma left lower ribs.

CASE II. X-ray Lilly D. Black dot marks nipple. Whole left chest shows dark shadow. Note calcareous material close to vertebral column and the well-marked lateral curvature.

CASE I. View of tumor from inside. Note at the bottom of the photograph the peritoneum covering the diaphragm, through which cartilaginous nodules are growing. Photograph by L. S. Brown.

felt uncomfortable in regard to the depth of the anesthesia and oxygenation. There was no doubt that the expansion of the lung could be regulated at will, but I think it probable that the severe purulent bronchitis, coming on immediately after the operation, was caused by the concentrated ether vapor and deficient aëration in the box. Whether this operation could have been performed without positive pressure is a matter of opinion. There was no doubt that the patient's condition became worse as soon as the pressure was reduced, but was alarming only for a few moments after division of the diaphragm. I recall very clearly a case of my father's, Dr. C. B. Porter, where he removed a sarcoma which involved three ribs and a small part of the diaphragm. In spite of complete collapse of the left lung, there was little if any change in the pulse and respiration. Other similar cases are on record. My father's patient recovered without complications, except for slight late suppuration in the wound.

As a result of his experience in this case Dr. Murphy has suggested certain improvements in the apparatus which may do away with the objectionable features, or Dr. Robinson's new positive pressure apparatus may fulfill all indications.

While it is unfair to attribute a death, fourteen days after operation, to the method of etherizing, I cannot but feel that the patient's strength was seriously if not fatally undermined by the early bronchitis, which I think was probably due to the method of anesthesia. In the future, in the absence of special apparatus, I should consider: first, the advisability of producing an artificial pneumothorax by the injection of nitrogen gas, operation to be done at the end of a week, after the lungs and heart had become adjusted to the new conditions; second, operation without either negative or positive pressure, the lungs and mediastinum to be immediately steadied by traction, as soon as the chest was opened; third, preliminary suture of the lung, after removal of a rib, and secondary operation after adhesions were firm.

CASE II. Lillie D., aged fifty-one, entered hospital June 17, 1908. Family history: Father, one brother and two sisters died of consumption. Past history: "Eighteen years ago, lung trouble, when her heart was on the wrong side." She had pain through the left chest for a number of months, but was never tapped. Two years later, cough and hemoptysis. She has never been strong; a tendency to be short of breath, but has had no real illness until last December, when she had a hacking cough and pain in her left side on drawing a deep breath. Eight weeks ago, clothing too tight. Six weeks ago, lump in left side. Examination: small, pale, rather well-nourished woman. Heart: left border not determinable on account of flatness of left chest; right border three quarters of an inch from the sternal line; impulse palpable in costal angle; soft systolic murmur at apex. Lungs: left, dull down to third rib, flat below in front. Distant bronchial breathing, with diminished vocal fremitus; behind, flat at apex down to spine of scapula. Bronchial breathing with increased fremitus

and numerous râles at apex. Bronchovesicular breathing below. At base, almost complete absence of breath sounds and fremitus. Few scattered râles through left back. Right chest normal save for a few râles. Over lower three ribs in left axillary line is a tumor three by four inches, which is slightly boggy; no distinct fluctuation; slight tenderness and redness. X-ray shows no necrosis of the ribs; the whole left cavity in shadow; diaphragm depressed. Temperature and pulse normal, urine negative, no leucocytosis. Diagnosis: Chronic tuberculous empyema probably of many years' duration, now gradually pointing between the ribs.

Operation under ether, June 20, 1908. Incision into the tumor showed the muscles and subcutaneous tissues honeycombed and filled with a puriform fluid, which escaped by a tortuous opening between the eighth and ninth ribs. Five ribs, for a distance of six inches, were resected, revealing the following condition: The whole cavity from apex to diaphragm was filled with a soft, semi-fluid material resembling ripe Camembert cheese. The lung was completely collapsed against the vertebral column, firm and leathery, non-crepitant. It was covered with calcareous masses which were intimately adherent to the visceral pleura. Across the center of the cavity ran a cord about the size of a lead pencil, which after division spurted blood, evidently a patent branch of the pulmonary artery. There was no lung tissue whatever in the apex. The parietal pleura was about 1 cm. in thickness, tough, bluish-red in color, showing no evidence of tuberculosis or granulation tissue. It was impossible to irrigate away the greasy contents of this large cavity, so it was rapidly wiped as clean as possible and packed with six large handkerchiefs of gauze; the margins of the skin were protected with rubber dam, over which a large dressing with swathe was applied.

Examination of the muscles and subcutaneous tissue showed no evidence of tuberculosis. Cultures were sterile. The wound was dressed with the greatest care in order to avoid secondary infection. One by one the large pieces of gauze were removed, and in spite of some fever and the drain from this large wound, the patient left the hospital in six weeks in satisfactory general condition. On Sept. 20 she returned for further operation. The wound was granulating healthily; the calcareous masses had entirely separated from the remains of the lung. No fever, no leucocytosis; hemoglobin, 80%.

Operation, Sept. 24, under ether. An incision was carried upwards two inches from the vertebral column to the level of the spine of the scapula and down to the ribs. All the muscles were scraped outward to the mid-axillary line; there was free hemorrhage. The ribs were rapidly divided with rongeur forceps up to the third, and removed with the underlying thickened pleura. No attempt was made to free the obsolete lung, and, owing to the patient's condition it seemed inadvisable at this operation to remove any more of the parietal pleura. The cavity was again packed with gauze. Convalescence was satisfactory, and the patient left the hospital on Nov. 23, 1908, in the following condition: Patient has a good appetite, sleeps well; there has been no fever for ten days. The amount of discharge is steadily diminishing. The ends of the divided ribs are covered with healthy granulations. The anterior flap has become adherent to the diaphragm and is rapidly rolling in to obliterate what was once the lower half of the chest. Further operation will be necessary, when the patient has recovered her strength, to insure the closure of the upper part of the cavity, kept open by the second, third and fourth ribs.

This case is interesting because it showed so many auscultatory signs, where no lung was present, and also on account of the long duration of a sterile empyema.

X-RAY PLATES OF INTERLOBAR EXUDATES.

BY A. K. STONE, M.D.

THE clinician often meets with serious difficulty in determining the location of an exudate in the pleural cavity. If one side of the chest is full of fluid and the lung compressed it is a perfectly simple matter to insert a needle, preferably at the points of election, as described by Drs. Shattuck, Sears and others, and draw off the fluid.

There are times, however, when the exudate lurks in the deep recesses of the pleural cavity and seems to take special delight in eluding the exploring needle. At the same time the daily rise and fall of the temperature and the general condition of the patient makes the anxious physician feel that the exudate must be found and removed, as pus is probably present. With this in view

long and often sizable needles or trocars are thrust through the chest wall, so "as to get through the thickened pleura."

Although my experience has been comparatively small, I think that the pathologists will agree with me that it is very rare to have a pleura thickened beyond a quarter to a half inch, and such rare conditions are found almost always in old long-standing cases. Therefore I feel sure that large and long needles should be discarded and that the lung should not be subjected to violence by them.

It is not uncommon to have evidences of fluid in the chest and yet be perplexed as to its where-

abouts. Dullness even to flatness is usually present; the fremitus is lost; the breath sounds come to the ear faintly but still distinctly in all parts of the chest, or at times friction is heard distinctly over the whole of the dull area. Marked displacement of the heart is not apt to be found.

In cases where the physical signs are as described above the exudate will be found almost always between the lobes of the lungs.

Fluoroscopic examination of the patient in the sitting position, supplemented by an x-ray plate taken during full inspiration, may give the desired information regarding the seat of the exudate, and the picture conveyed to the examiner is very different in the cases of interlobar exudates and in the exudates localized up the chest wall.

The two plates that I have to show are both of interlobar exudates of the left side. The dense shadow is noted in each case as running from the region of the heart's apex upwards and outwards towards the axilla. In each case there is comparatively clear lung above this shadow and to a lesser extent below. It is this moderate shadow in the lower portion of the chest that makes it necessary to have the plate taken during full inspiration, so that as much light as possible shall come through the chest above the depressed diaphragm.

In the former of these cases there was profound constitutional disturbance, and on physical examination there was found a bulging of the anterior wall of the thorax over the third and fourth ribs; there was also dullness and well-marked friction sounds everywhere over the left side of the thorax, but no displacement of the heart to the right. The diagonal shadow on the plate was very dense and pus was suspected both from shadow and from the constitutional symptoms. When the chest was opened in the axillary line a large amount of clear fluid was evacuated. The examination of this exudate and the subsequent course of the disease showed this to be a case of tuberculous pleurisy.

In the second case (see plate) the shadow was no denser than in the previous plate but the fluid was found to be pus. Here the removal of the sixth rib in the anterior axillary line enabled Dr. Cobb to enter a well-defined cavity between the lobes of the lung.

I have no plate to show of a right-sided interlobar pleurisy, but from a study of the position of the lobes I should expect a similar appearance of the plate. The one patient with such a condition that I have had a chance to examine both with a fluoroscope and shadow plate, unfortunately for me, cleared up and the fluid absorbed, as proved by a later plate, the temperature also becoming normal. Therefore there was no operation to demonstrate the accuracy of the examination.

Looking at an anatomical plate of the chest with the lungs represented in place, it will be seen that the sulcus between the two lobes of the left lung crosses the anterior axillary line at about the fifth space or beneath the sixth rib, and a resection of this rib allows the circumscribed exudate