Original Articles.

VENTRAL HERNIA.¹

BY JOHN W. KEEFE, M.D., PROVIDENCE, R. I.,
Surgeon to the Rhode Island and the St. Joseph's Hospitals.

VENTRAL hernia is a subject about which the textbooks of the present day say but little, and in consequence the practitioner is prone to advise an abdominal support and say that nothing further is of value. The good results obtained by surgeons during the last decade in operations performed for the radical cure of hernia have led me to address you on this subject and to present cases in illustration of this distressing condition and its treatment by operation.

Umbilical hernia is most frequently found in middle-aged women, childbearing and obesity being most commonly the causes. One of the unfortunate results occasionally attending the many celiotomies of the present day is a hernia at the site of the abdominal incision, most frequently observed in operations for appendicitis where the wound has been allowed to heal by granulation. Bull reports fifty cases following operations for appendicitis which presented themselves at the Hospital for Ruptured and Crippled during a period of two years. The length of the incision, cutting off the nerve supply of the muscles, dividing muscular fibres, and the use of gauze drainage are factors worthy of careful consideration while operating for appendicitis.

The diagnosis of a ventral hernia can usually be readily made by inspection and palpation. The coverings of the hernia are skin, subcutaneous tissues and peritoneum. The contents are omentum, intestine, or both.

The hernia may cause no inconvenience for years and for this reason many are permitted to attain a great size and then may be attended with distress, difficulty of locomotion, pain, nausea, vomiting, strangulation of the intestine and peritonitis. Constipation and other digestive disturbances may be due to hernia.

Spontaneous cure has never been known to occur in a case of ventral hernia in an adult. Abdominal binders and trusses may prevent increase in the size of the hernia and give support, thus giving a feeling of comfort to the patient but never a cure. Proper surgical treatment, in selected cases, is the best method to employ for radical cure of a ventral hernia. In many cases the operation is simple and attended with little risk to the patient and a complete cure of the hernia. In herniae of long standing, the contents being irreducible, those of large size, those where there is atrophy of the muscles surrounding the hernia, and in old subjects, an operation may be contraindicated.

One may meet with one of the most difficult operations the surgeon has to deal with, in a hernia containing adherent and gangrenous omentum and intestine. The high percentage of mortality attending operations for the cure of strangulated hernia is a beacon warning us to operate early, when the hernia is of small size, and the patient in good physical condition, and not debilitated by pain and vomiting, perhaps stercoraceous in character. Bull and Coley report a mortality of less than one per cent. in four hundred operations for radical cure of hernia.

The patient may be prepared for the operation by

¹ Read at the meeting of the Rhode Island Medical Society, December 1, 1896.

daily baths, cathartics and enemas, with frequent doses of benzo-beta-naphthol, which I have found of great value in producing a collapsed condition of the intestines. A bichloride poultice applied over the abdomen some hours previous to the operation, followed by scrubbing and the use of the numerous antiseptics, — mercury, permanganate of potash, oxalic acid and a liberal washing with normal salt solution, — renders the site of the operation as nearly sterile as is necessary.

The incision may be over the most prominent part of the hernia or, preferably, elliptical, including the hernia sac. The sac in the vicinity of the ring should be dissected free and the muscle exposed on either side. When we have a reducible hernia to deal with the contents should be returned to the abdominal cavity before opening the sac. Where the contents are adherent, the omentum, intestine, or both should be carefully dissected from the sac and the superfluous or gangrenous omentum excised, a chain catgut ligature being used to control hemorrhage. The sac, including the fibrous ring, should be removed and the peritoneum united with catgut. My preference for bringing the fascia together is a buried silver-wire mattress suture or a machine-stitch of silk-worm-gut, the ends of which are brought out through the skin at either end of the wound, thus enabling one to remove them at the end of four weeks. Schede, of Bonn, Barker, of London, and Halsted, of Baltimore, prefer the buried silver wire, while Coley and others always use kangaroo tendon. The adipose tissue and skin may be united with catgut, or silk-worm-gut may be used in the integument. Sterilized gauze, cotton and a well-fitting binder constitute the dressing.

The after treatment consists of hot water for the first twenty-four hours, with an eighth of morphia if necessary, followed by easily digested nourishment.

CASE I. Female, age sixty-six years, entered St. Joseph's Hospital May 25, 1896. Married forty years. Had ten children; last child twenty-one years old. Noticed an umbilical hernia four years ago; has worn a truss for one year. About seven days ago she was unable to reduce the hernia and complained of pain in her stomach and vomited. Bowels have not moved well since, but an enema caused a slight movement.

May 25, 1896, 4 p.m. Operation under ether. Hernial sac very tense. An incision was made into the sac, which was found to contain omentum and a strangulated knuckle of intestine. The ring of the hernia was incised and the hernial contents returned to the abdominal cavity. The sac and redundant portions of skin were removed. The peritoneum was closed with catgut and fascia with a double row of mattress silk-worm-gut sutures. Interrupted silk-worm-gut sutures united the edges of the skin.

May 27th. Bowels moved freely.

June 3, 1896. The wound was dressed for the first time to-day and the superficial sutures were removed. The wound healed by first intention. The patient returned to her home three weeks from the day upon which she had her operation performed.

CASE II. Female, age thirty-eight years, entered the Rhode Island Hospital March 21, 1897. She has had an umbilical hernia for sixteen years, which during the past four years has caused her considerable trouble. She has been unable to reduce the contents of the hernia for the past three days and complains of a great deal of local pain. Examination reveals an

irreducible umbilical hernia about the size of a large orange, which feels quite hard.

March 21, 1897. Operation under ether. An elliptical incision was made in the skin about the hernia. The hernial sac was dissected free and opened. It was found to contain omentum, several sections of which were tied off with catgut and the distal portions of omentum removed. The sac was excised and the peritoneum united with a continuous catgut suture. The fascia was dissected free and united with buried silver-wire mattress sutures. Silkworm-gut was used to bring the edges of the wound together.

An aseptic dressing was placed over the wound and held in position with a wide abdominal binder.

April 6th. First dressing. The silkworm-gut sutures were removed. Healing has taken place by first intention.

April 12th. The patient is sitting up.

April 20th. Discharged well.

CASE III. Female, age forty-seven years, was admitted to the Rhode Island Hospital March 28, 1897. About twenty years ago she noticed a small protrusion at the umbilicus which gradually increased in size until two years ago when it reached the size of one's head. Binders and trusses were used without giving relief. She then commenced treatment with

one of the advertising hernia-cure specialists in town. She was treated by rest in bed and injections into the sac for a period covering one year. The hernia decreased in size but never disappeared. Three weeks ago the hernia became strangulated and was attended with pain, vomiting and fainting spells. These symptoms disappeared in a few days. On examination, an umbilical hernia was found about the size of a cocoanut with contents irreducible. An operation for radical cure was advised.

March 31, 1897. Operation under ether. An elliptical incision was made through the skin about the hernia. The sac was then dissected free to the muscular aponeurosis forming the ring. The sac was opened and the interior found trabeculated and filled with adherent omentum. This omentum was ligated with a chain-stitch of catgut and about two-thirds of the mass in the hernial sac was removed, the remainder being returned to the abdominal cavity. The edges of the ring were approximated with a buried silver wire. The muscular aponeurosis was dissected free on either side of ring and united with buried mattress sutures of silver wire. The skin wound was closed with silkworm-gut and an aseptic dressing applied.

April 10th. First dressing. Silkworm-gut removed. Healing by primary union.

May 5th. Discharged well, with advice to wear an abdominal band. One and one-half years later no return of the hernia.

CASE IV. Female, age forty-five years, entered the Rhode Island Hospital January 24, 1898. She was operated upon December 15, 1895, by one of the surgeons of the hospital for radical cure of a small umbilical hernia. The ring was stitched together with catgut, and over this the adjoining fascia was drawn with silkworm-gut. The skin was united with silkworm-gut. The patient was discharged cured.

Examination to-day shows a ventral hernia about as large as an English walnut occupying a position near the centre of the scar resulting from the previous operation. The preparatory treatment consisted of easily digested food, ten-grain doses of benzo-beta-naphthol every four hours and frequent doses of magnesium sulphate. Patient has a cough with a slight amount of expectoration.

January 26, 1898. Operation, chloroform and oxygen being used instead of ether on account of the chest symptoms. An elliptical incision was made in the skin about the hernia. The hernial sac was dissected out and opened. It was found to contain a mass of omentum which was easily reduced. The sac, when removed, was found to be much larger than it appeared before the operation. The peritoneum was united with catgut. The fascia, which was dissected free from the ring of the hernia, was held in apposition by two silkworm-gut sutures, used as a cobbler's stitch, the ends of the sutures then being brought out through the skin at either end of the wound. Buried interrupted kangaroo-tendon sutures were also used to hold the divided fascia together. Silkworm-gut sutures were employed to close the wound in the skin and subcutaneous tissues. An aseptic dressing was placed on the wound and covered with a large flannel binder. During the first twenty-four hours after the operation drachm doses of hot water were given every fifteen minutes if the patient was awake.

January 27th. Patient has had very little pain, no

nausea or vomiting. Cough annoying, so was ordered a codeia cough mixture.

January 29th. Patient very comfortable. One-tenth of a grain of calomel was given every ten minutes until ten doses were taken, then two drachms of magnesium sulphate repeated in three hours. About the time a movement might be expected a soap-suds enema was given. The bowels moved at least once a day while patient remained in the hospital.

February 4th. Wound dressed for the first time and superficial stitches removed. Primary union.

February 16th. An unsuccessful attempt was made to remove the cobbler's stitch of silkworm-gut.

February 18th. The cobbler's stitch was to-day removed without any difficulty.

February 22d. Patient's subsequent history uneventful. Discharged to-day feeling well.

CASE V. Female, age forty-four years, entered the Rhode Island Hospital January 28, 1898. This patient noticed a swelling at the navel about three months ago. This swelling appeared after she had lifted a heavy stove. Examination reveals an umbilical hernia about the size of an egg, but the patient says that at times the swelling is considerably larger. The patient was put to bed and given easily digested nourishment. Five grains of benzo-beta-naphthol was administered every four hours. Sulphate of magnesia was given every morning and followed by an enema.

January 31, 1898. Operation under chloroform and oxygen. An incision in the skin about the hernia was made and the sac dissected free down to the hernial ring. The sac was found much larger than it appeared to be by the external examination. The sac was opened and found to contain omentum which was adherent in many places. A large mass of omentum was tied off with interlocking catgut ligatures and removed. The hernial sac was excised, and the edges of the wound in the peritoneum united with a continuous catgut suture. The edges of the fascia were approximated with buried silver-wire mattress sutures and the edges of the skin with silkworm-gut. An aseptic dressing was applied. During the first twenty-four hours nothing but drachm doses of hot water were given every fifteen minutes when the patient was awake. The patient rallied well from the operation, no stimulation being necessary, and complained of only a little pain.

February 3d. One-tenth of a grain of calomel was given every ten minutes until ten doses were taken, followed by two doses of magnesium sulphate, three hours apart, and followed in two hours by a soap-suds enema. The bowels moved freely.

February 6th. Similar cathartics were given.

February 8th. First dressing. Silkworm-gut removed. Primary union.

February 26th. Discharged well.

November 21, 1898. Examination shows no evidence of return of hernia.

CASE VI. Female, age 38 years, entered the Rhode Island Hospital February 28, 1898. Two years ago she was operated upon for appendicitis with the formation of an abscess. Incision and drainage of the abscess were the methods employed, with the gradual closing of the wound by granulation. Two weeks after the wound had healed she noticed a small hernia at the site of the incision. This hernia gradually grew larger and caused so much inconvenience that she consulted an advertising sure-cure-for-hernia

specialist, who treated her by injections into the hernial sac twice a week during a period of about seven months. The alleged cure failing to cure, she consulted a surgeon, who operated for the cure of the hernia eleven months ago. Three days after the operation the hernia reappeared and is now as large as two fists. It is irregular in shape, the ring admitting four fingers. This cannot be replaced in the abdominal cavity, only a portion disappearing when the patient is in a recumbent position. She was again treated by the hernia-cure specialist twice a week for a period of four months. Patient was given ten-grain doses of benzo-beta-naphthol every four hours and daily doses of salines followed by an enema.

March 5, 1898. Operation under ether. An elliptical incision about eight inches long was made, includ-

ing the old cicatrix. The sac was dissected out with difficulty on account of the depth of the wound, there being about three inches of fat in the abdominal wall; the irregularity of the sac; and a diverticulum extending for some distance between the external and internal oblique muscles. The intestine and omentum were adherent to the sac and it required careful dissection to free them. Portions of omentum were tied off with catgut and excised. The peritoneum was united with a continuous catgut suture. Several buried wire mattress sutures brought together the transversalis and internal oblique muscles, and a continuous silver-wire suture approximated the severed edges of the external oblique muscle. The wound in the skin was closed with silkworm-gut and an aseptic dressing of iodo-

form gauze, sterilized gauze, cotton, and a binder applied. Drachm doses of hot water were given during the first twenty-four hours, with one-eighth grain of morphia for pain.

March 5th. Calomel, a saline and an enema caused a free evacuation from the bowels.

March 9th. Wound dressed and a small amount of pus was found in the lower angle of the wound.

March 10th. Superficial stitches removed.

April 24th. Lower part of wound healed by granulation.

April 27th. Discharged. No evidence of reappearance of hernia.

November 29, 1898. Examination shows no evidence of return of the hernia.

CASE VII. Female, age forty-nine years, entered St. Joseph's Hospital October 2, 1898. She has borne four children and had one miscarriage at three months. About fifteen years ago she noticed a small swelling at the navel, which has never wholly disappeared. The swelling has gradually increased in size until now it is as large as a child's head. Examination reveals a large umbilical hernia whose contents are irreducible. The skin covering the mass is ulcerated and has the appearance of an epithelioma. At frequent intervals during the past year she had had vomiting spells lasting from a day to ten days. Lately they are scarcely two weeks apart. The abdomen is very large, the wall containing a thick layer of fat. Rest in bed, benzo-beta-naphthol, calomel, salines, enemas and easily digested nourishment constituted the preparatory treatment. After a bath and thorough scrubbing with soap and water a bichloride poultice was worn during the night previous to the operation.

October 5, 1898. Operation under ether. The abdomen was scrubbed with ethereal soap solution, potassium permanganate, oxalic acid, bichloride of mercury, ether and sterilized salt solution. An elliptical incision was made about the hernia, the sac dissected free and opened. A large mass of omentum and several coils of intestine were found adherent to sac, which was trabeculated. The intestine and omentum were carefully dissected free, portions of the omentum being removed after ligating with a chain-catgut suture. The remainder of the contents of the sac were returned to the abdominal cavity and the sac excised. The peritoneum was dissected from the fascia and united with a continuous catgut suture. Several buried silver-wire mattress sutures brought the fascia firmly together. The skin was united with silkworm-gut. Owing to the thick layer of fat three deep silkworm-gut sutures were passed through the skin, fat and fascia at some distance from the wound on either side. An aseptic dressing was applied. The operation was long and tedious, owing to the trabeculated condition of the sac and the firmly adherent intestine and omentum; yet at the conclusion of the operation the patient's pulse was good, no stimulant having been given. Saline enemas given and drachm doses of hot water with coffee during the first twenty-four hours.

October 15th. Ten days after the operation the three tension silkworm-gut sutures removed. Slight amount of pus about one of the stitches.

October 19th. The silkworm-gut suture in the skin was removed. Primary union.

November 3d. Patient sitting up.

November 5th. Discharged well. There is firm union of the wound.

CASE VIII. The following case whose photographs I present with this history illustrates the enormous size an umbilical hernia may attain. One is better able to judge of the size of the hernia by comparing it with other portions of the body than if measurements were taken. At the time of my examination of the hernia I judged that fully three-fourths of the intestine and omentum occupied the hernial sac, rendering operation out of the question, as the present abdominal cavity was not large enough to hold all of the intestine contained in the sac. Another interesting feature of this case is the unusual hypertrophy of the mammary glands, one reaching to the umbilicus, as shown in the photograph. The patient did not consider the large size unusual or of any special importance, having suffered no inconvenience from them. This case emphasizes the value of early operation.

This patient, age sixty-two years, a hardworking woman, mother of five children, noticed twenty-eight years ago, while carrying her fourth child, a small swelling in the umbilicus. During the following ten years the swelling increased in size, but she thinks it has remained stationary during the past eighteen years. When she was married her weight was only one hundred pounds. After reaching her thirtieth year her weight has gradually increased until at present she weighs two hundred and sixty-two pounds. She had the hernia ten years before she wore a support of any kind. Trusses she found of no value, so she contrived an ingenious band to fasten about the body with a bag attached in front to receive and support the hernia. During the past five years she has had almost constant diarrhea. She has had only two attacks that appeared to be due to strangulation of the hernial contents, one two years ago and the other, for which I was consulted, one and one-half years ago. At present she runs a sewing machine and does the greater part of her own housework including washing.

A CASE OF MEDIASTINAL SARCOMA.¹

BY GEORGE G. SEARS, M. D., BOSTON.

JACOB D., a Russian, twenty-five years old, a carpenter by trade, was admitted to my service at the City Hospital in August, 1898. He knew nothing of his family history, and had paid so little attention to his own symptoms that only a very meagre account of them could be obtained. Last December he noticed that he was growing short of breath and that his face and neck became swollen when stooping or working hard. Two months ago he began to cough and occasionally spit up a little dark blood, and three weeks ago he noticed that his chest had enlarged. There has been no pain, but he has had a feeling of soreness across the lower portion of his chest, and has lately lost considerable flesh.

Physical examination showed a fairly nourished but anemic man with cyanotic extremities. The jugulars and the larger veins of the arms and in the axillæ, especially the right, were dilated and prominent, and at the level of the insertion of the diaphragm in front was a zone of dilated capillaries, while the veins of

¹ Read at a meeting of the Clinical Section of the Suffolk District Medical Society, November 16, 1898.