CASE 3.—The most interesting results were obtained from C. A., man, aged 29, admitted to Cook County Hospital, Dec. 18, 1913, and assigned to the care of one of us (W. W. H.). Two weeks previously he had been at Mercy Hospital (Chicago) for excessive thirst, general weakness, and frequent urination. He had been sick seven weeks, and during that time had lost 20 pounds. He was hungry all the time and had been eating large amounts of food since leaving Mercy Hospital. He was in the last stages of a severe diabetes mellitus. In spite of a restricted carbohydrate and large doses of sodium bicarbonate, he was growing progressively weaker. His ketonuria was marked; and his dextrose to nitrogen ratio was 1:35.

Feb. 11, 1914 (first trial), the empty stomach possessed good tone and showed moderate hunger contractions. Five days later (Feb. 16, 1914) the empty stomach showed two moderate and one powerful period of hunger contractions in less than three hours. This is rather unusual; for after a twelve hours' fast, hunger periods recur about every two hours. During one of the three periods mentioned above, all but a few cubic centimeters of the chloroform was expelled from the manometer, so that the graphic record is not complete. The fact is significant that the patient became restless and irritable toward the end of the experiment, and continually asked for food. Because of this increasing restlessness and irritability, the experiment was discontinued for obvious reasons.

The patient went into diabetic coma five days later, and died the following evening (Feb. 22, 1914).

It is regretted that other duties prevented us from recording observations between February 16 and February 21. Our experiments surely indicate that the contractions of the empty stomach are responsible for the sensations of hunger experienced by diabetics. We do not expect to find

that other patients so near death from other causes (save in special easily explainable instances) would speak of eating as often and as freely and exhibit hunger contractions of the same severe type as shown by this patient.

Incomplete as the results are, they nevertheless confirm the work on animals suffering from a fatal pancreatic diabetes when the polyphagia was attributed to the marked hunger contractions resulting from the inanition which is so prominent a feature of the disease.¹ For it was shown⁵ that the blood of an animal showing the polyphagia of pancreatic diabetes similar to the blood of a starved animal possessed the property of increasing the vigor of stomach contractions when such blood was injected intravenously into a normal animal.

SUMMARY

The method employed in studying the movements of the normal empty stomach (hunger contractions) is feasible for use in the study of movements of the stomach in pathologic states which point either directly or indirectly to an involvement of the stomach. Patients experienced no difficulty in swallowing the balloon; the apparatus necessary is simple and inexpensive; the

results obtained are of great interest and value both to the clinician and physiologist.

The results obtained so far warrant the following statements:

1. The hunger contractions obtained from a neurosthenic conformed to the normal, though interpreted by him as painful. They were neither more vigorous nor more frequently repeated than those of any normal person. His prompt "cure" justified our conclusion that we were dealing with a patient not suffering from any serious organic gastric disorder.

2. Hunger contractions were obtained in a case of cancer of the stomach without marked pyloric obstruction. The contractions were moderately powerful. The patient was quite emaciated (in a state of inanition), but took interest in food and experienced hunger during the contractions.

3. When, however, an acute gastritis is brought on by dietary indiscretions, there results a cessation of contractions of the empty stomach, and this is accompanied subjectively by a lack of interest in food and an absence of hunger (anorexia).

4. Most cachectic states are accompanied by an anorexia. Further work will no doubt show that this is due to the absence of well defined hunger contractions. The cachexia of diabetes mellitus, however, is accompanied by a more or less well marked polyphagia. We find that the latter results from frequent and vigorous contractions of the empty stomach (hunger contractions) which are responsible for the polyphagia exhibited by patients suffering with this disease.

Fig. 4.—Record of the stomach contractions in Case 3. As can be seen, the period ends in tetany at the close of which (b) a powerful contraction forced a considerable amount of chloroform out of the manometer.

CHYLOUS ASCITES AND CHYLOTHORAX

DUE TO CARCINOMA OF THE STOMACH

JOHN H. OUTLAND, M.D.

Surgeon to the Swedish and Bethany Hospitals

AND

LOGAN CLENDENING, M.D.

KANSAS CITY, MO.

Chylous ascites—the occurrence of ascitic fluid which is of a milky character, and the color of which is due to the presence of chyle or of fat globules (pseudochylous)—is a sufficiently rare condition to make a case of extreme interest, both clinically and physiologically. Several cases in which the condition resulted from a traumatic rupture of the receptaculum chyli have been studied by physiologists, who, assuming that they were dealing with pure chyle, have been interested in watching the effect of various diets on the composition of the chyle.

In 1910 Wallis and Scholberg¹ reported three cases and reviewed the literature, collecting reports of 173 cases; but they missed the cases reported by Hall and Morton,² Nichols,³ Batty Shaw⁴ and Edwards.⁵

1. Wallis and Scholberg: *Quart. Jour. Med.*, iii, 301; iv, 153.

2. Hall and Morton: *Northwestern Univ. Bull.*, June, 1910.

3. Nichols: *Med. News*, New York, Nov. 11, 1905.

4. Shaw, H. B.: *Jour. Path. and Bacteriol.*, 1900, vi, 339.

5. Edwards: *Practice of Medicine*, 1909.

5. Luckhardt, A. B., and Carlson, A. J.: On the Chemical Control of the Gastric Hunger Mechanism, *Am. Jour. Physiol.*, 1914, xxxvi, 37.

Since their article, cases have been reported by Gandin,⁶ Calwell,⁷ Marrack,⁸ Lediard,⁹ Collins,¹⁰ and Huber and Silver.¹¹

The article by Wallis and Scholberg is quite exhaustive, giving an abstract of all the 173 cases of chylous and chyloform ascites and the sixty cases of chylothorax, with complete analyses of the fluid obtained in their own cases and an extended discussion of the conditions.

In the cases abstracted by Wallis and Scholberg, there are seventeen in which the etiology is given as carcinoma of the stomach. Of these, in three—the cases of Zawadski (1891), of Lydhecker (1893) and of Menetier and Gauchler (1902)—there also was chylous fluid in the thorax. To these three we add here a fourth:

REPORT OF CASE

History.—W. T. B., man, aged 46, horse buyer, Anglo-Saxon, referred by Dr. Riley of Benedict, Kan., had a father, mother and one brother living and well. No brothers or sisters were dead. He possibly had a syphilitic infection fifteen or twenty years before. Some antisyphilitic treatment was carried on for a year, but not vigorously. He used to

Fig. 1.—Gross specimen: A, A, liver; B, gallbladder; C, mesentery of the intestine filled with glands; D, stomach, with tumor in the pyloric region.

use whisky too much, but took a cure about five years before, and had not used it since. He had never been in tropical countries, other than Texas, where he has been within two years.

He felt well up to six months before examination (April 16, 1915). The onset of his illness was sudden. One day in the afternoon he felt dizzy and vomited. He did not go to bed, but was weak and dizzy and continued to feel much the same way for two months. He vomited three or four times in that period. He has never vomited blood. In the last three months he has been troubled with indigestion, pain after meals, belching and bloating. The bowels are constipated.

March 27, 1915, the abdomen began to swell very suddenly, and in a few days was enormously distended. It was tapped, April 1, and about a gallon of milky fluid was removed.

April 8, it was again tapped, and 1½ gallons of ascitic fluid of the same character were removed.

An edema of the scrotum and penis was observed at about the time of the appearance of the ascites.

April 16, he was removed to the Swedish Hospital from his home in Benedict, Kan.

Physical Examination.—At this time the temperature was 99; pulse, 110; blood pressure, 135. Urine: specific gravity, 1.020; acid; a trace of albumin; no sugar; no casts. Blood: hemoglobin, 105; reds, 4,000,000; whites, 17,000. Differential count of whites showed normal relationship. The Wassermann reaction was negative.

Fig. 2.—Duodenum, showing lacteal vessels engorged and lumpy in appearance over its surface.

The patient looked old for his age. His hair was white. The veins on the cheeks were dilated. Heart: The apex was just outside the nipple line. There were no murmurs. A diffuse and soft swelling was on the left side of the neck just below the clavicle. The lungs were clear with normal vesicular murmur everywhere except in back of left chest, where there was dullness with no breath sounds and evidence of fluid. There was edema of the scrotum and penis, and enlargement of the glands in both inguinal regions. The skin was slightly edematous everywhere, not more so in the legs than over the chest and in the skin of the abdomen. The abdomen was enlarged, with evidences of fluid. No filariae were found in the blood.

Course.—April 17, the abdomen was tapped, and over a gallon of milky fluid withdrawn. The chest was aspirated at the same time, three and a half quarts being removed. This chest fluid was clear straw colored. The fluid removed from the abdomen was milky in appearance, with a tinge of pink. It contained 1.9 per cent. of albumin and fat, and did not react to Fehling's solution. Microscopically it showed small fat globules, and small specks in constant motion. Leukocytes and erythrocytes were numerous.

Fig. 3.—Small intestine, near end of jejunum; lymphatic vessels are stained with silver nitrate; note thickness of mesentery.

April 23, the abdomen and the chest were again tapped. Chylous fluid was found this time in the chest as well as the abdomen. A swelling in the left side of the neck the size of a small orange was observed. It receded in size when fluid from the abdomen was removed. After the fluid had been removed, a tumor extending across the upper abdomen, above the umbilicus and more marked on the left side, could be outlined on palpation. It was recorded that it felt like a tumor of the transverse colon. The liver was two fingers below the costal arch.

6. Gandin: *Ergebn. d. inn. Med. u. Kinderh.*, 1913, xii, 219.

7. Calwell: *Brit. Med. Jour.*, Feb. 21, 1914.

8. Marrack: *Quart. Jour. Med.*, 1912, vi, 462.

9. Lediard: *Clin. Jour.*, 1914, xliii, No. 9.

10. Collins: *Ann. Surg.*, 1913, lvi, No. 4.

11. Huber, Francis, and Silver, H. M.: A Case of Acute Chylous Ascites (Non-Fatty, Pseudochyloous, Latent or Milky Type) in a Boy Eight Years Old, *Am. Jour. Dis. Child.*, July, 1914, p. 50.

May 1, the abdomen and the chest were tapped. A large quantity of milky fluid was obtained from each.

May 12, the abdomen was tapped; 1½ gallons of chylous fluid were removed.

May 13, the patient died.

Necropsy.—A partial postmortem examination was made, May 13, at 8:30 p. m. The body was that of a middle-aged

Fig. 4.—Section of tumor in the stomach.

man. There was no rigor mortis. There was edema of the penis and scrotum. There was a swelling on the left side of the neck. When the abdomen was opened, about 3 quarts of milky fluid escaped. The intestines were distended and covered with a lymphatic exudate. The cecum and appendix were enormously distended. Small lymphatic channels over the surface of the intestine were greatly distended. All the glands of the mesentery were enlarged. The entire mesentery was a thick mass of glands which were soft in consistency.

The liver was enlarged and edematous. The gallbladder was large. There were no gallstones.

A tumor was present along the lesser curvature of the stomach, extending from the duodenum to well within the pylorus. The glands of the gastrohepatic mesentery were enlarged. No growth was found in the rectum.

Milky fluid was found in the left side of the chest. The lungs everywhere were clear and crepitant. There was no enlargement in the mediastinum. There was no aneurysm. The thoracic duct could not be located. The entrance of the thoracic duct into the large veins could not be made out.

The entire small intestine, stomach, liver and mesentery, with the covering of the abdominal aorta, were removed in one piece.

COMMENT

The case was interesting. The shifting of symptoms from day to day was alone sufficient to make it such, and besides, the diagnosis was always in doubt, ranging from filariasis to thoracic aneurysm. The case of Nichols,³ of which we had a report, and in which the ascites was due to cirrhosis of the liver, influenced us to cling to that diagnosis most constantly in view of our patient's alcoholic history. After he had been a week in the hospital the tumor, felt in the upper abdomen, made the diagnosis of carcinosis of the abdomen seem very reasonable.

Of especial interest was the swelling on the left side of the neck just above the clavicle. Was this due to an enlarged or tense thoracic duct? It was

noticed that it receded in size after tapplings of the abdomen or chest. It extended gradually up the neck, day by day, between tapplings, until it reached half way up to the sternomastoid muscle.

Was there an obstruction of the receptaculum chyli, either a thrombus or a metastatic plug? We are unable to state. The specimen removed from the body was taken with an especial view to a dissection of the great lymphatic duct, but while it was attempted no part of it was found. The demonstration of the receptaculum chyli or of the thoracic duct is one of extreme difficulty, and rarely satisfies any one but the dissector himself. After attempting it in this case we are regretfully inclined to take with a grain of salt the very glib reports of its condition made by other reporters of cases of chylous ascites.

As we did not find the receptaculum in the abdomen or the thoracic duct in the thorax, we feel inclined to believe that they were not affected in this case, as a metastasis of the tumor would have been seen in either of them. We believe that the chyle escaped into the abdomen through the small lymph vessels of the intestine and mesentery. We know, in support of this belief, that they were all very much dilated, that the lymph nodes all through the mesentery were enlarged, and that metastases were found in the small lymph vessels of the intestine and in the nodes from every portion of the mesentery (Figs. 2, 3, 4 and 5).

The fluid removed from the abdomen and the chest gave the evidence of being true chyle; it did not putrefy on standing, it did not clot, and it contained no macroscopic particles of fat. A great deal is made of the differences between chylous and pseudo-chylous and chyloform fluids by other authors. Huber and Silver make the following distinctions:

Fig. 5.—Section of mesenteric gland, showing metastasis.

1. True chylous effusion is due to the presence of chyle.
2. Chyloform or fatty, though resembling the first, owes its milky appearance to emulsified fat, the result of a fatty degeneration of cellular elements.
3. In the third variety "the opalescence is not due to the presence of fatty emulsion but is caused by the

presence of opalescent substances, the exact nature of which is still undetermined."

Wallis and Scholberg have a much simpler conception of the different nature of the fluids. "Chylous effusions would include, as before, those due to trauma or to the escape of chyle in diseases affecting the lacteal and lymphatic systems." Chyliform are all other kinds.

CASES IN THE LITERATURE

The article by Wallis and Scholberg, as we have stated above, is the most complete monograph in English on the subject. We have not attempted to confirm their references. The cases which we have found in the literature not included in their article, or reported since, are as follows:

Hall and Morton² report a case due to trauma; a wagon passed over the abdomen of a young boy. Many tapings were made, and true chyle obtained. The case was studied chemically by Hall in relation to diets.

Nichols³ reports a case in a man, aged 74, due to cirrhosis of the liver.

Two tapings were made, and 5,700 and 6,200 c.c. of fluid, respectively, obtained. The man died, and there was a necropsy.

Calwell⁷ reports a case due to nephritis (?). The patient was tapped once, and 3 pints were drawn off.

Marrack⁸ reports two cases, one due to appendicitis and one to nephritis.

Lediard⁹ reports four cases, one due to tubercular peritonitis, one to Hodgkin's disease, one to heart and liver disease, and one to a malignant cyst from the pelvis.

Collins¹⁰ reports a case due to trauma.

Huber and Silver¹¹ report a case in a child, aged 8, due to no determinable cause. An operation was performed and an opportunity given to examine the intestines, when the lacteal vessels were seen engorged, as in our case. A subcutaneous drainage of the peritoneum resulted in a cure. They quote Gandin, a reference to which we do not have access.

Lead Poisoning in Pottery Trades.—The causes of lead poisoning in the pottery trades are the lead ingredients, of which the glazes are partly composed. The most dangerous process is that of mixing the glaze, applying it to the ware and then removing the excess of the glaze from the ware, handling the ware while the glaze is still wet, decorating and painting the ware and also sweeping the rooms in which the glaze has been handled. Much of the glaze is without lead and is harmless; but in a large percentage of the various pottery wares, lead glazes are used. In only a few of the factories investigated by Dr. Hamilton did the percentage of lead exceed 20. In certain tile works, the percentage in the glaze ranges from 40 to 60. The workmen, besides being exposed to lead poisoning, also inhale much mineral dust and suffer from the effect of great heat in the firing of the enamels.—C. M. Price, *Modern Factory*.

DEVELOPMENT OF THE STATE HOSPITAL AT KANKAKEE (1880-1890) *

RICHARD DEWEY, A.M., M.D.,

WAUWATOSA, WIS.

In the latter half of the nineteenth century, many problems connected with the public care of the insane presented themselves for solution which developed conflicting views. Some of these questions were: the advisability of separate provision for the acute and chronic classes; the extent to which mechanical restraint was necessary or admissible; the need of establishing laboratories for psychopathologic research, of instituting training schools for attendants, of broadening and diversifying industries among the inmates and of increasing their privileges, and even of maintaining in some instances "open" instead of locked doors. Among all the questions, however, that which for a time took precedence of others concerned the introduction of a new type and style of building—the substitution of modern "segregated" or "detached" buildings, domestic in style, for the antiquated "congregate" structures previously in universal use. This paper is intended, among other things, to describe the new departure in architecture which was ushered in at Kankakee between 1880 and 1890.

Today, when this transformation has long been *un fait accompli*, it seems sufficiently simple; but in its inception it involved much controversy and debate which was at times tinged with acrimony.

After the close of the Civil War, a great mass of insane pauperism seemed to become suddenly a public burden. This incubus had been growing all along, but the people did not become

aware of it until after the war and its all-absorbing issues had been disposed of. Then the necessity for asylums became apparent, and every state in the Union, in its turn, found itself engaged in a struggle with this unwelcome and ever increasing burden. In the several states one asylum after another was provided, but in nearly every instance, long before the doors were open, more than enough hapless defectives had accumulated in the almshouses and jails of the state to fill it to its utmost capacity. A grim struggle with this increasing mass of human driftwood has been constantly in progress up to the present time. The struggle seems absurdly unequal; for the forces of the state continually lag behind, balking at the enormous expenditures involved. The situation may be likened to a "forlorn hope" assailing an inaccessible height. It

Fig. 6.—Section of intestinal wall, showing metastatic plugs in the lymph channels.

* This article is a revision of the writer's history of Kankakee, one of the forthcoming series of histories of "The Institutional Care of the Insane in the United States and Canada," to be published this year by the American Medico-Psychological Association, under the editorship of Dr. Henry M. Hurd. Volume I has already appeared from the Johns Hopkins Press.