not been destroyed by the disease. If, during the mastoid operation, the stapes was displaced or removed, the labyrinth should be opened. If the operator could not be sure that this accident had happened he should watch for labyrinth symptoms and operate on the labyrinth the moment they appeared. He thought that Mr. West's double vestibulotomy was a good operation and might prove to be useful at times. The Viennese method of removing the whole of the posterior wall of the petrous, however, was necessary in suspected disease of the posterior fossa. Simple vestibulotomy would not, of course, reveal a latent abscess in the posterior fossa.

Mr. C. E. WEST, in reply, said, regarding the Vienna mode of exposing the dura of the posterior fossa, that meningitis was really only a remote contingency of labyrinth operations, and if it did occur he drained through the internal meatus. He did not favour a wide exposure of the dura. Labyrinth operation was, he thought, indicated in general infective labyrinthitis, and when vertigo after the radical mastoid caused continued incapacity for work. In reply to Mr. Hunter Tod and Dr. Dundas Grant, he held that labyrinthitis could not be excluded unless a histological examination was made, and in this Mr. Sidney Scott agreed. In order to expose the promontory, he had found that free removal of the posterior meatal wall was quite sufficient.

The PRESIDENT, in conclusion, said that the names of Jansen, Bárány, Scott, and West would be linked together as the pioneers of labyrinth surgery, just as Wilde, Schwartze, Macewen, Stacke, and Ballance were the protagonists of the surgery of the mastoid.

(To be continued.)

BRITISH MEDICAL ASSOCIATION.

Meeting in London, Wednesday, July 27, 1910.

SECTION OF LARYNGOLOGY.

Mr. H. TILLEY, *President, in the Chair.*

Abstract Report by Drs. DAN MCKENZIE and HAROLD KISCH.

The PRESIDENT, in his opening remarks, made a sympathetic reference to the loss laryngology had sustained in the death of Mr. Cresswell Baber. Among the visitors to the Section he welcomed Drs. von Eicken, Luc, Bryan of New York, and H. S. Birket of Montreal.

Passing to more technical matters, the speaker first of all gave a brief *resumé* of the progress laryngology had made since its elevation to the ranks of a speciality, then he mentioned one or two of the more recent developments of the science, alluding more particularly to the subjects of bronchoscopy and œsophagoscopy,

and of tonsil enucleation. Attention was drawn, in a few brief and pregnant sentences, to the growing feeling of conservatism in the matter of operating in suppuration of the nasal sinuses, and especially of the frontal sinus. Finally, the President drew attention to various problems connected with the work in which further research was necessary, such as the place of vaccine therapy in nasal suppuration, and the factors which influenced the transformation of latent bacterial infections into active virulent disease.

DISCUSSION ON THE TECHNIQUE OF THE DIRECT EXAMINATION OF THE
ESOPHAGUS AND LOWER AIR-PASSAGES.

Introduced by PROF. DR. VON EICKEN (Freiburg), and
DR. D. R. PATERSON (Cardiff).

Dr. von EICKEN dealt with the direct examination of the air-passages. With reference to the method of illumination, the speaker thought that most people preferred an outer light, because an inner light encroached upon the lumen of the tube, and was liable to be dulled by blood and other secretions or discharges. The fixed lamp was the most convenient for many reasons, and it was especially suitable for beginners, but there were many cases in which the Kirstein light was to be preferred, and among these children were to be included.

The passing of the tube was sometimes a matter of some difficulty, especially in children. It was important to press down the tongue in order to isolate the epiglottis before attempting to pass the beak of the tube over the epiglottis. In passing through the glottis the end of the tube should be carried into the posterior commissure. Another plan, which was especially serviceable in the case of children, was to bring the tube at right angles to the glottic opening. Once the tube had reached the trachea care should be taken to keep the lumen of the trachea always in view. When the bronchus was reached the tube should be carried along its lateral wall, and scraping the end of the tube against the median wall of the bronchus should be avoided. At the bifurcation the right bronchus seemed like a continuation of the trachea.

The most difficult cases to examine were cases of bronchiectasis, for in that disease the tube was liable to be flooded with the copious foetid secretions. This could be minimised by adopting the following procedure: For one hour before the examination the patient is made to assume the lateral horizontal position with the healthy lung down, in order that the bronchiectatic cavities may

be emptied of their contents; then the patient is examined in the sitting position.

Difficulties and Dangers.—In children the passage of the tube through the narrow glottis was apt to be followed by acute laryngeal stenosis for which tracheotomy might be required. For that reason, therefore, the smallest possible tubes should be employed.

In adults with prominent incisor teeth and a short muscular neck the difficulty of inserting the endoscope might be overcome by carrying the tube into the angle of the mouth. Brüning's counter-pressure autoscope was especially useful for these cases, as by pushing back the larynx it brought the anterior commissure into view. For the same purpose Killian's angle speculum was of great service.

In order to reach the right bronchus the tube should be carried into the left angle of the mouth, and *vice versa*. It was at times impossible to change the tube from one angle of the mouth to the other without withdrawing it.

The co-operation of the patient should be enlisted in order to restrain coughing, etc., and it was advisable to instruct him to make a pre-arranged signal if he felt pain or serious discomfort.

With regard to general anaesthesia chloroform was preferable to ether, because it did not irritate the mucous membrane and stimulate secretion. Even when a general anaesthetic was used cocaine and adrenalin should be applied, especially in cases where a foreign body was lodged in the air-passages, because in these cases there was swelling and engorgement of the area surrounding the foreign body. In young children cocaine and other sedative drugs were very dangerous, much more dangerous than chloroform.

Examination of the air-passages was contra-indicated in arteriosclerosis and heart disease, and in aneurysm it was improper to use this method for the sole purpose of determining the exact site of the disease.

For these reasons a general clinical examination of the patient should precede the examination, and in cases of suspected tumour or abscess an X-ray examination should also be made. In children death has occurred as a result of merely extending the head in cases of abscess of the lower cervical vertebræ compressing the trachea against the manubrium sterni. This could be avoided by performing the extension cautiously, noting meantime whether any already existing dyspnoea was exaggerated during the movement.

Tracheotomy was called for before attempting to pass the tube when there was severe dyspnoea. If tracheotomy relieved the

dyspnoea an indication was thereby afforded that the obstruction was situated in the larynx. This operation was also called for in many cases of foreign body in the air-passages. When, for example, the foreign body was large no attempt should be made to withdraw it through the larynx, because it was liable to lodge below the glottis. Such foreign bodies as beans also necessitated tracheotomy because they rapidly swelled and became soft, so that when they were seized they crumbled into pieces, any one of which might remain behind or be aspirated into the bronchi.

The difficulties attendant upon the removal of foreign bodies such as bones which were frequently found buried in granulations had been overcome since Brünings had invented his toothed forceps. In like manner the same surgeon had supplied forceps for removing collar-studs and an expanding instrument which ensured the removal of hollow foreign bodies. As a result most of these could be removed by upper bronchoscopy, but there were still a certain number of cases in which it was impossible to remove the body save through a tracheotomy opening. For the case of foreign body lodged behind a bronchial stricture Brünings had invented a dilating bougie by which the stricture could be dilated and the foreign body removed.

The speaker related a case recently under the care of Prof. Killian in which a tack had been removed by an instrument which was made so as to enclose the head. In this case there was a stricture which was subsequently dilated with perfect success.

Dr. von Eicken concluded by giving a demonstration of the various instruments employed in the direct examination of the air-passages.

Dr. D. R. PATERSON, who confined his remarks to the subject of the direct examination of the œsophagus, said that œsophagoscopy differed from tracheoscopy in several ways. The trachea was an open tube with rigid walls, while the œsophagus was a collapsible tube with soft walls. At its upper end the œsophagus was closed by a sphincter which opposed the easy passing of the endoscope, and once the instrument had reached the œsophagus there was more tendency for its point to wander. This contrast between the two passages necessitated a corresponding difference in manipulation. As in tracheoscopy so in œsophagoscopy, all cases should be examined by the X-rays as well as by the ordinary clinical methods before direct examination was attempted.

With reference to illumination the speaker had found that in an emergency sufficient light could be obtained from an ordinary good

incandescent lamp if a large tube was used. He was in the habit of using an internal lamp, a forehead light, or a fixed light according to circumstances. Each had its uses. The drawback to the fixed light, a drawback which even the newest inventions had failed to remove, was that the light was in the way when instruments had to be introduced. He himself used the Kirstein lamp most frequently because it permitted more freedom in operating.

The œsophagus should never be examined blindly and the mandrin pilot had only a very restricted use. In this matter he entirely differed from Guisez, who had recently stated that the passage of a flexible bougie should always precede direct inspection. This procedure exposed the patient to a double passage of instruments and increased his discomfort very considerably.

Save in children and nervous adults local anæsthesia was sufficient. For a general anæsthetic chloroform was commonly employed. He had also found that ether by the open method was free from disadvantages.

The sitting position was only possible for five or ten minutes. After that time the mucus secreted by the œsophagus found its way into the larynx and set up coughing. This did not occur when the patient was lying down on his side or back.

In introducing the tube the point should be kept strictly in the middle line and passed over the interarytænoid region in this position. When the lip of the introitus projected the patient's head should be moved, he should be asked to swallow, or the tube should be rotated. It was at this point that the pilot sometimes proved useful. Once in the œsophagus, the deviation of the organ to the left should be remembered and the tube carried to the right angle of the mouth. "Sticking" was usually due to faulty direction.

In cases with ulceration, the most scrupulous care and delicacy were called for, because even with the ocular method perforation of the wall of the gullet had occurred.

Beaked tubes were less useful for the gullet than for the larynx. He preferred the old rounded tube, although it was more difficult to introduce. For the œsophagus, also, telescopic tubes were less adapted than for the air-passages, where the dimensions became reduced as we passed downwards.

Œsophageal forceps should be larger and stronger than bronchial forceps.

Dr. WM. HILL said that elongated tubes or cylinders of the same calibre throughout their entire length were almost universally employed

for endoscopy of the œsophagus, trachea, and bronchi; and extension tubes also were constructed on the cylindrical principle. He had found, however, that for the larynx and gullet elongated funnels possessed marked advantages over tubes of uniform calibre. Owing to the fact that the hand-lamp was attached centrally to the shaft of the endoscope there was no room for the passage of swabs and instruments by the side of the lamp down the endoscopic tube without displacing the hood and thus shutting off the light; moreover, the proximal end of the tubular endoscope being of the same narrow calibre as the distal end, the handle of the forceps or other instrument lay in the line of vision and obstructed the view in the narrow tube. In his own funnel-like endoscope the lumen of the proximal end of the instrument had a cross sectional area more than double that of the distal end, and by an eccentric attachment of the hand-lamp there was ample space for the passage of the instruments, sponge-holders, etc., by the side of the hood of the hand-lamp without cutting off the light. Forceps, etc., could be passed down close to one side of the funnel, and the inserted instruments, except at their distal ends, were not in the centre of vision. The immense gain in simplifying technique could only be fully appreciated after actually employing a funnel-like endoscope. He had shown before this Section at the Belfast Meeting last year a new pattern operating laryngoscope. That instrument was not a complete funnel, but had a lateral slit to give facility in the manipulation of large forceps, etc. Following on this he had designed elongated funnels on the same principle for employment in the gullet. These were at first complete funnels, but in order to facilitate the removal of the œsophagoscope after placing a radium apparatus in malignant strictures, he resorted to the principle of the slit-funnel, as in the laryngoscope. This pattern of endoscope also facilitated many other endo-œsophageal procedures, such as the dilating of strictures, the removal of portions of new growths, and especially the extraction of foreign bodies. For example, instead of using a special apparatus for the closing of an open safety-pin in the gullet before attempting extraction, with his slit œsophagoscope one could draw the pin-point of the safety-pin within the lumen of the endoscope with forceps, where it could not catch the mucosa; the jaws of the forceps were then transferred to the other limb of the safety-pin, traction on which partially shut it, and it could be drawn up through the endoscope with safety. The distal end of his endoscopes was bevelled, so that it was possible to pass the instrument throughout its whole course under vision, as in Brüning's tube-spatula. This was especially necessary in traversing the deep pharynx and cervical œsophagus, which was the dangerous area where perforations were occasionally made. With a bevelled-ended instrument passed under vision this calamity should not happen, more especially if the patient was directed to swallow when the post-cricoid narrowing was passed, and to cough so as to open the lumen when the mouth of the œsophagus was reached, and whilst traversing its collapsed cervical portion. If difficulty was experienced in passing the instrument on account of stricture in the post-cricoidal pharynx and upper œsophagus, the instrument should not be passed onward until the lumen has been dilated up by graduated bougies. The passage of the œsophagoscope had been described as an easy procedure, but it was also easy to perforate the gullet, and in some instances it was probable that neglect of caution in the region of the mouth of the œsophagus had led to these regrettable incidents. He had heard of six cases of œsophagosopic perforation of the gullet in this country alone with three deaths, so that a word of caution was not out of

place. The cross-section of his œsophagoscope was distinctly elliptical at the proximal end and nearly circular at the distal end, and the reason for this and for the long diameter being transverse was obvious. Extension tubes he considered to be a needless complication in œsophagoscopy technique; moreover, they are liable to break at the spring and to clog in the groove after frequent hospital use and sterilisation. Brüning's two smallest bronchial spatulæ with extension apparatus were certainly useful in dealing with the lower air-passages of infants and small children; but for older patients he preferred the slit funnels. His endoscopes were made of steel. These new pattern œsophagoscopes had been in daily use at St. Mary's Hospital for nearly a year, and those of his colleagues who had employed them found that they greatly simplified endoscopic surgical technique in the gullet. Dr. Hill concluded by detailing the dimensions of the instruments he had devised.

Dr. STCLAIR THOMSON said that when the indirect method could be used it should be resorted to in preference to the direct method for the removal of laryngeal neoplasms, the application of the galvano-cautery, etc. He had found the direct method disappointing in trying to remove minute growths from the anterior commissure. With regard to instruments he preferred the old fish-tail end to the beaked end.

Mr. E. WAGGETT remarked that the results when using the endoscope were better under cocaine than under a general anæsthetic. He advised that the patient should be instructed to make a signal when the examination became unbearable, and in order to enlist the patient's confidence the surgeon should promise to respect his wishes.

Mr. STUART-LOW had found that warming the cocaine improved its efficacy.

Mr. SCANES SPICER thought there was a tendency to use cocaine carelessly, and recommended that excess of cocaine solution should always be mopped up.

Mr. PERMEWAN agreed with Dr. StClair Thomson that it would be a misfortune if, as a result of the new method, the skill of laryngologists in the old method were to decline.

Dr. N. C. HARING had found the hypodermic administration of $\frac{1}{150}$ gr. atropin before chloroform was given useful in preventing an excessive secretion of mucus. He used a solution of 20 per cent. cocaine in alcohol.

Dr. IRWIN MOORE complimented Mr. William Hill upon his instruments.

Dr. BROWN KELLY had found that in children the larynx was most easily reached by the tube when the patient was lying on the side with the chin flexed. Referring to cardio-spasm, he said that the condition was generally diagnosed as malignant stricture during life. He had recently seen a case in which all attempt to pass the tube into the stomach had failed. After death it was found that the patient had been suffering from malignant disease of the lesser curvature of the stomach, and that the growth had pushed the mucous membrane of the viscus into the lower end of the œsophagus in such a way as to block it.

Dr. ANDREW WYLIE had been able to get rid of the trouble of excessive laryngeal secretion by small doses of potass. and sod. bromid. for a few days preceding examination.

Dr. BRYAN (New York) related a case of a child into whose trachea a grain of corn had slipped. Owing to swelling about the subglottis the tracheoscope could not be inserted, and tracheotomy had to be performed. Further interference was, however, rendered unnecessary by the spontaneous expulsion of the foreign body through the tracheotomy wound.

Dr. KIRKLAND had seen at Vienna, in a case of asthma, the contraction of the bronchi.

Dr. WALKER DOWNIE, regarding anæsthetics, had got the best results from cocaine preceded by bromides.

Mr. T. GUTHRIE had found the Kirstein lamp more serviceable than the Brüning light in cases of papillomata in childhood.

Dr. DONELAN agreed with the recommendation to give bromides. He also advised the retention of the old method in the removing of laryngeal growths.

Dr. PEGLER referred to the old-fashioned plan of spraying the larynx with bromide.

The PRESIDENT had had a large experience of foreign bodies removed by tracheo-bronchoscopy, many of which were on exhibition in the museum. He related the case of a child with papilloma of the larynx in which, on admission to hospital, a diagnosis of diphtheria had been made and antitoxin administered before he was asked to see it. In one of his cases, a baby, aged four days, a teat used as a "comforter" had stuck in the œsophagus. In children as young as this the larynx and trachea are so small that the presence of the tube in the gullet with the head extended is sufficient to obstruct the air-way completely. In examining the larynx of young children the patient should be turned on one side, the head flexed, while an assistant pushes up the larynx. He preferred a Kirstein lamp for many purposes. In an emergency half a dozen wax candles or a bicycle lamp proved to be a sufficient source of illumination. In addition to bromides he recommended the sucking of ice for a few minutes before the tube was passed.

Dr. von EICKEN agreed with Dr. StClair Thomson that for laryngeal manipulations the indirect method should be tried first. He had often been able to see a foreign body in the bronchus by means of the ordinary laryngeal mirror. As to anæsthetics, he preferred chloroform to ether because it was less irritating to the mucous membrane. Bromides were useful if there was time to give them, but in the case of foreign bodies, when promptitude was an important factor, it was, of course, impossible to delay until the bromide took effect. He repeated his warning of the danger of cocaine in childhood. The head should not be pushed too far back in the case of children. The diagnosis between cardio-spasm and cancer of the lesser curvature of the stomach was difficult, but in such cases he employed a special instrument in order to remove a portion of the diseased tissue for pathological examination. An important point in the diagnosis, however, lay in the fact that in cardio-spasm there was a history of dysphagia of some years' duration. With reference to Dr. Hill's split tubes he thought he would prefer a tube spatula with a Kirstein light. Acute cases of asthma could be treated with bronchoscopy and the injection of adrenalin.

Dr. D. R. PATERSON, in reply, drew attention to the importance of rapidity in examining and operating on the bronchial passages.

Dr. WM. HILL, in reply, said he was sceptical of what was called cardio-spasm. In any case, if the obstruction was due to spasm it would disappear under an anæsthetic. He thought that what was taken for spasm was tumefaction. He had seen thirty-seven cases of malignant stricture but had never yet come across a case of spasm.

CONGENITAL STENOSIS OF THE CHOANÆ.

BY MR. J. SMITH FRASER (Edinburgh).

Of 75 reported cases of congenital atresia of the choanæ 31 were males and 44 females. In 40 the atresia was bilateral.

Post-mortem specimens of the condition were rare, but he had succeeded in getting one.

The case was that of a female with suppuration in the left ear. The right choana was closed by a septum. The patient was not a mouth-breather and yet the palate was higher on the side of the atresia. Thus choanal atresia could not be cited in support of the usual theory as to the production of Gothic palate. The girl had died of otitic meningitis consequent upon syringing the attic.

The PRESIDENT remarked that several of these cases which had been successfully operated upon had been shown at the Laryngological Section of the Royal Society of Medicine.

Dr. DAN MCKENZIE said that one of the cases had been shown by him. He agreed with Mr. Fraser that no argument could be based upon the condition of the palate found in atresia of the choana in support of the open-mouth theory of deformed palate.

Dr. BROWN-KELLY had also had a similar case under his care.

Mr. W. STUART-LOW remarked that in these cases it was important to straighten the septum before operating on the atresia.

The PRESIDENT said that the rule was to remove not only the web, but also the posterior end of the vomer.

(To be continued.)

PROCEEDINGS OF THE AMERICAN LARYNGOLOGICAL, RHINOLOGICAL, AND OTOLOGICAL SOCIETY.

Meeting, 1910.

CONSIDERATION OF THE END-RESULTS OF THE OPERATION FOR SUBMUCOUS RESECTION OF THE NASAL SEPTUM.

BY DR. FREDERICK C. COBB (of Boston).

The paper showed the results of the submucous operation after the lapse of several years. The cases were operated by men sometimes experienced in the procedure, sometimes almost new to it; as a result the number of faults in technique were at first large, and perforation quite common. Fifty cases were