average as high in courage as others, other things being equal. They are not infrequently cited for gallantry. Some do heroic acts in a kind of impulsive manner like that in which they sometimes commit criminal actions. Some epileptics display symptoms resembling those of neurasthenia, and some who have been trephined present, in addition to all this, attacks of jacksonian epilepsy. In treatment, a quiet environment, sodium cacodylate and a milk-vegetable diet were the main reliance. In the great majority of the cases the epilepsy dated from early life but had been aggravated by campaign conditions.

Singer, H. D. MENTAL HYGIENE. [J. A. M. A., July 20, 1918.]

Singer says that mental hygiene is too much a closed book to the general practitioner and that training in this special line is neglected. The usual notion, he thinks, is that a healthy body is all that is necessary for healthy mentality. The fallacy of this view has been strongly emphasized by the results of examinations and experience in the formation of our new army. While it is not probable that obvious defects have been overlooked in the accepted cases, over 1 per cent. of those passed have been later found unfit because of some mental or nervous deficiency. Military training is not altogether physical, and he asks, is there not even more need of training in mental adjustments. The morale and *esprit de corps* of a good soldier and the general need of discipline and self control may be even more essential. Singer says: "Let us consider for a moment the factors involved in the consideration of mental adjustments. These may be divided under the two heads of (1) quantity, (2) quality. The former is structural; the latter functional." The best that we can do at present to meet the former is try to ascertain the brain tissue available and train it in a right direction. The tests are being constantly evolved and their application is simple. The quality, on the other hand, represents the use which is made of the available tools, and how far this is due to hereditary causes and to subsequent training and education is unknown. Whatever may be the true relation between these two causative factors, training and habits are far more readily open to interference than heredity. But we must first answer certain questions, namely: "1. What constitutes a healthy or satisfactory adjustment? 2. What are the indications of a liability to fail? 3. What measures must be taken to remedy or minimize these faulty tendencies?" These questions cannot be answered with absolute certainty. The first requires the consideration of the following: How far has the individual learned to observe the rules for the regulation of conduct which have been gradually evolved in civilized society to the restraint if necessary of his biologic desires. Singer brings in as an illustration the development of a war neurosis in the patient as the result of finding himself in an intolerable position through his war environment, a condition he would not have to face in ordinary civilized life. Proper hygienic study of such cases may lead to their prevention. As an answer to the third question, the methods to be followed, the author says, are similar in kind

to those adopted in the ordinary methods used today for the treatment of war neuroses. They concern education, the development of interests and habits, and have nothing to do with punishment. The necessity of the medical practitioner giving his attention to these matters cannot be questioned, while expert knowledge and direction are also essential. The state should provide the means for this training of the physician, and the coöperation of local physicians is also essential. Our medical colleges have so far only partially met the needs of mental hygiene. Their courses on its subjects are brief and insufficient. If this war brings prominently to the fore the importance of mental health to national efficiency it will have accomplished an inestimable benefit.

Uhlman. MOBILIZATION PSYCHOSES IN THE SWISS ARMY. [Med. Rec., June 22, 1918. Ed. from *Correspond. bl. f. Sch. Aer.*]

This arthur refers to the well-known fact that at the outbreak of a war there is a general alteration in the psyche of the entire community and an increase of psychic morbidity in both the military and civilian population. A contingent of psychoses also follows mobilization of an army even in peace times. The Swiss troops were mobilized at the outbreak of the present war and the author by communicating with the heads of the insane asylums obtained data as to the number of psychoses during and for several months following mobilization which were the direct result of the latter. There is of course no one cause for war psychoses but the content tends to have a military coloring and the type of the mental alteration is severe. Fear effects are common—a man fears that he will be thought a spy or traitor or deserter and expects to be shot.

In the first ten days of August about 250,000 troops were called out and 32 were evacuated to insane asylums. From August 1 to December 31 the total number of such evacuations was 151. In other words, the reaction due to mobilization was soon followed by equilibrium. Monthly figures show a steady decline and a curve for the five months' period shows a sharp rise which culminates August 10 and is followed by a short sudden and a long gradual fall. Among the mobilization psychoses those due to alcohol are always well represented and of the 151 cases 35 were of this origin. The proportion is especially high in the older men of the landwehr and landsturm—12 cases out of 39. The proportion of alcohol psychoses in the marching troops was 20 per cent. It would, however, be unfair to regard all such cases as examples of delirium tremens for subjects with endogenous psychoses may drink to excess. It is almost certain that some of these cases should be credited to dementia præcox. All of these alcohol psychoses were chronic. The author has no single classification of dementia præcox and follows the asylum diagnoses, which evidently are not patterned throughout on Kraepelin's classification. We find 61 cases of dementia variously termed katatonic, hallucinatory, paranoid, and hebephrenic, which doubtless jointly answer to dementia præcox, and adding the 35 alcoholics