dently squeezed outwards by the pressure of a solid substance behind them. On opening the chest, it was seen that its cavity was encroached upon by the diaphragm having been pushed up. There were slight adhesions at the apex of the left lung, and one or two old tubercles, in a quiescent state, in the corresponding part of the right lung. The lungs and heart presented nothing else worth notice. The spleen was of very large size, weighing three pounds eight ounces; its texture was gristly and hard, but there was no distinct or circumscribed morbid deposit in it. The great omentum was firmly adherent to this enlarged spleen, and puckered up so as to form a rather short band of adhesion between it and the transverse colon. The liver was also very large, weighing six pounds one ounce and a quarter; its texture appeared healthy. The large intestine and the lower part of the ileum contained bloody fluid in many parts, but no breach of surface could be found, although the condition of the gut above was diligently examined. The right kidney appeared healthy; the left contained a large mass of crude tubercular matter of gristle-like consistence in its substance. The blood examined under the microscope showed an unmistakable increase in the proportionate number of the white corpuscles. The pancreas was healthy.

GUYS' HOSPITAL.

LEUCOCYTHÆMIA SPLENICA; THE PATIENT RELIEVED BY TREATMENT.

(Under the care of Dr. ADDISON.)

In the diagnosis of this important disease some physicians might refrain from making an examination of the blood, with the view of not abstracting any of that fluid by venesection from an already impoverished body. This can readily be obviated by a slight puncture of the finger with a needle, when a sufficiency is at once obtained to determine its condition. In the notes of the following case, taken by Mr. F. M. Cann, Dr. Addison's clinical clerk, it is not mentioned whether the essential and predominating feature of leucocythæmia were present; but all doubts on this point are set at rest by information which we have derived from Dr. Wilks, who says that there was an excess of white corpuscles over the red. The spleen is enormously enlarged in this patient, but whilst under treatment it became somewhat diminished, and for the present she has left the hospital slightly relieved. She has never been the subject of ague.

Sarah B—, aged twenty-four, unmarried, a cook, admitted into Lydia ward, Oct. 12th, 1859, under Dr. Addison. Up to her present illness was always healthy. About six months ago, after lifting a heavy saucepan and carrying it a short distance, she set it down, and then felt something "snap," as she describes it, in her left side beneath the mamma. She continued very unwell during the whole day, and when in bed felt unable to lie in the horizontal position, and consequently passed nearly the whole night sitting up in bed, suffering pain in her side, which persisted three or four days. She then became better, though feeling very weak. A fortnight later the pain returned in her side, and then descended to the left iliac region; this attack lasted eight or nine days, and during it she noticed that her abdomen swelled somewhat at night after undressing, but resumed nearly its natural size by the morning. When between twenty-one and twenty-two years of age (nearly three years ago) she continually suffered from flushing of her face and cold feet and legs; and has within the last few months suffered also in the same way. She has never had any distinct shivering fits followed by sweating. She states that for the last six years she has noticed herself become stout in the winter, and has got thin again in the summer. Commenced menstruating at fourteen years of age, and occasionally suffers severe pain during that period. When sixteen years old, she ceased to menstruate for eight months. Four months ago the catamenia stopped for eight weeks; the discharge then commenced with severe pain in the loins; it was dark-coloured, thick, and foetid, and lasted five days. Sixteen days afterwards she suffered in the same way, and has seen nothing since. Has had leucorrhœa, at times, for years.

On admission, she presented an emaciated and anæmic appearance; eyes sunken; lungs and heart healthy, but the sounds of the latter audible over the whole anterior part of it. On deep inspiration, she felt a dragging pain over the seventh and eighth ribs at their junction with the cartilages. A large swelling

occupied the greater part of the abdomen, and was dull on percussion, extending from the left hypochondrium across the epigastrium to the right hypochondrium; also in the umbilical, left iliac and lumbar regions, and downwards into the hypogastric region almost to the pubis. On loosening her dress at night, she feels pain in the epigastrium; and whilst walking, suffers uneasiness over the whole swelling, which appears to increase in size, as it were, whilst she is moving about. Bowels regular; passes very much urine in the morning, and but little during the day, is always high coloured, and deposits a large sediment on cooling, which is redissolved by heat, and contains no albumen; tongue dry, and feels thirsty.

Oct. 12th.—Put on four grains of quinine every four hours.

14th.—Half a drachm of strong mercurial ointment to be rubbed every night over the region of the spleen.

16th.—During the last six months, particularly when in bed, she suddenly awakes, and finds herself perspiring profusely, her night-dress being occasionally saturated, and immediately afterwards feels very cold. She does not always perspire, but sometimes suffers from "dry heat" all over her body, which prevents her sleeping. She is now dressed, and her skin is perspiring unnaturally. Has lately suffered from lowness of spirits, and feels inadequate to any exertion, even to talk. Complains of loss of memory, and feels as though nothing made an impression on her.

17th.—Soon after taking a meal she feels a sinking sensation in her stomach. The tumour is certainly decreasing.

18th.—A blister ordered to be applied to the splenic region.

20th.—Sulcus of spleen is in a line with the umbilicus.

24th.—Pain at times felt over the whole swelling. The quinine appears to be the cause of pain in the head. A blister to the splenic region.

26th.—To have eight grains of quinine at night, besides her day doses as usual. Specific gravity of urine 1015; no albumen.

28th.—Not much improved. Three grains of quinine, one grain of extract of henbane, and a quarter of a grain of blue pill to be taken three times a day.

Next day the swelling was rather larger, and she felt more pain on the left side than she had ever previously suffered.

Nov. 1st.—Ordered one grain of bromide of potassium, with two grains of extract of hyoscyamus, twice a day.

7th.—Coughs a good deal. Severe pain on pressure over the tumour, which has somewhat diminished; its sulcus was on a level with the umbilicus. She afterwards had a slight attack of bronchitis, and on the 18th the urine was high coloured and loaded with lithates. She was put on four grains of quinine every four hours.

Next day, four ounces of wine were ordered, as she was weak and low.

26th.—Great pain in swelling, which is much indurated. Has leucorrhœa, but her catamenia have been absent four months and a half.

Dec. 16th.—She left the hospital, slightly relieved.

KING'S COLLEGE HOSPITAL.

LEUCOCYTHÆMIA SPLENICA.

THE PATIENT SUBJECT TO AGUE TWELVE YEARS BEFORE;
RELIEF BY TREATMENT.

* (Under the care of Dr. GEORGE JOHNSON.)

A THIRD example of this disease is afforded by the subjoined case, differing in its history, however, from the other two in that the patient had been subject to ague some twelve years ago, with its accompanying congestion of the spleen. It would be irrational to infer that his present illness was in any way connected with, or consequent upon, his old ague, for the period of intermission has been sufficiently long to permit of the enjoyment of comparatively good health.

It is seldom, indeed, that an opportunity is afforded us of being enabled to place upon record three such well-marked examples of this interesting affection, in each of which the splenic hypertrophy has been so well observed, with an absence of all other disease. We did not perceive any of the extreme pallor of anæmia lymphatica, nor was there any lymphatic glandular enlargement in these cases. We might have added a few other examples, but the present are, in every way, sufficient to fully illustrate the malady.

The notes of the following case were taken by Mr. M. F. Anderson, clinical clerk:—

Meyer B—, aged twenty-nine, a pedlar, a native of